

UCLA

UCLA Previously Published Works

Title

Drinking patterns among Korean adults: results of the 2009 Korean community health survey.

Permalink

<https://escholarship.org/uc/item/2zh7010f>

Journal

Journal of preventive medicine and public health = Yebang Uihakhoe chi, 46(4)

ISSN

1975-8375

Authors

Ryu, So Yeon
Crespi, Catherine M
Maxwell, Annette E

Publication Date

2013-07-01

DOI

10.3961/jpmph.2013.46.4.183

Peer reviewed

Drinking Patterns Among Korean Adults: Results of the 2009 Korean Community Health Survey

So Yeon Ryu¹, Catherine M. Crespi², Annette E. Maxwell²

¹Department of Preventive Medicine, Chosun University Medical School, Gwangju, Korea; ²Fielding School of Public Health and Jonsson Comprehensive Cancer Center, University of California Los Angeles, Los Angeles, CA, USA

Objectives: In Korea, the proportion of deaths due to alcohol is estimated at 8.9%, far exceeding the global estimate of 3.8%. Therefore, this study was performed to examine the factors associated with low-risk, moderate-risk, and high-risk drinking patterns in Korean adults and to identify target populations for prevention and control of alcohol-related diseases and deaths.

Methods: We analyzed data from 230 715 Korean adults aged 19 years and older who participated in the 2009 Korean Community Health Survey. Multinomial logistic regression analysis was used to examine associations between socio-demographic and health-related factors and patterns of alcohol use.

Results: A substantially larger proportion of men than women engaged in high risk (21.2% vs. 3.4%) and moderate-risk alcohol use (15.5% vs. 8.2%). In both sexes, moderate- and high-risk uses were associated with younger age, higher income, being currently employed, smoking, being overweight/ obese, and good self-rated health.

Conclusions: Given the large proportion of the population that is engaging in moderate- and high-risk drinking and given the social norms that support this behavior, public health policies and campaigns to reduce alcohol consumption targeting the entire population are indicated.

Key words: Alcohol drinking, Epidemiology, High-risk drinking, Risk factors, Republic of Korea

INTRODUCTION

Alcohol use has been recognized as one of the major risk factors worldwide of preventable mortality and morbidity [1]. Alcohol use is the third greatest contributor to the global burden of disease, and is estimated to cause 3.8% of all deaths

and to result in 4.6% of all disability-adjusted life years lost and a disproportionate number of fatal injuries [2]. In Korea, alcohol use is estimated to cause 8.9% of all deaths [3]. More than half of these deaths are associated with binge drinking [4].

Alcohol use is associated with a variety of health outcomes that vary with the level of consumption [5]. In some studies, alcohol use in light to moderate quantities has been associated with better self-perceived health status, improved cardiovascular health, lower risk of osteoporosis in women, and lower rates of hospitalization [6,7]. In contrast, heavy drinking and binge drinking have been associated with unintentional and/or intentional injuries, interpersonal violence, human immunodeficiency virus infections, sexually transmitted infections, neurological damage, poor control of diabetes, hepatitis, hy-

Received: February 25, 2012 Accepted: May 31, 2012

Corresponding author: So Yeon Ryu, MD, PhD

309 Pilmun-daero, Dong-gu, Gwangju 501-759, Korea

Tel: +82-62-230-6483, Fax: +82-62-225-8293

E-mail: canrsy@chosun.ac.kr

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

pertension, gastrointestinal and heart problem, liver cirrhosis, cancers such as oral, rectal, and liver cancer, stroke, and alcohol dependence [4,5,8-10].

In a recent large nationally representative survey in Korea, the Korea National Health and Nutrition Examination Survey (KNHANES)-IV, monthly alcohol drinking of adults was reported to be 59.0%, and the prevalence of high-risk alcohol drinking was 18.0%. These rates show an increasing trend compared with that of the previous year [11]. Besides this statistical report, many epidemiologic studies in Korea have focused on the health effects of alcohol drinking [12-16].

It is important to understand and monitor alcohol use patterns to assess the public health influences of this behavior, to identify target groups for public health programs addressing alcohol use, and to plan and evaluate evidence-based strategies to reduce risky use. Although several previous studies in Korea, to some extent, have explained alcohol consumption, they did not successfully describe overall drinking patterns in Korea because of a small-sized population, the use of old data or some restricted sample such as older adults, adolescents, or college students [17-21]. To our knowledge, there is still a lack of evidence on alcohol use patterns in Korea, especially at the national level.

Therefore, this analysis was conducted to investigate alcohol use patterns based on the level of alcohol drinking, specifically, non-use, low-risk use, moderate-risk use, and high-risk use, in a nationally representative sample of Korean adults from the 2009 Community Health Survey, and to examine comprehensively the multiple factors, including socio-demographic characteristics, health behaviors, and health status, associated with alcohol use patterns relative to alcohol non-use.

METHODS

Subjects

This study used data from the public use files of the 2009 Korea Community Health Survey (KCHS) conducted by the Korea Centers for Disease Control and Prevention. The KCHS is an annual nationwide health survey conducted since 2008 to provide population-based estimates of health indicators to be used for the development and assessment of public health policies and programs. The 2009 KCHS used a multistage sampling design to obtain a representative sample of adults aged 19 years or older. Within each of 253 communities, 90 primary sampling units (PSUs) corresponding to smaller geographic

entities were randomly selected, followed by the random selection of 5 to 8 households within each PSU and an in-person interview of all adults in the household. Households were sampled from a registry of residents. To guarantee an adequate sample size in each community (about 900 persons in each community), when interviewers could not reach the sampled household after at least three visits, a replacement method was used, in which a randomly substituted household was selected [22]. The 2009 KCHS interviewed a total of 230 715 individuals.

Study Variables

The dependent variable in this study was the category of alcohol use pattern. Persons who had ever drunk any kind of alcoholic beverage during the past 12 months were classified as current drinkers and were asked more questions on the quantity consumed in a typical day and the drinking frequency in one time. We created categories of alcohol use patterns using the official definitions of KCHS to identify subgroups of individuals who used alcohol and whose drinking pattern may have been putting them at greater risk of harm relative to individuals in other groups [11]. We classified respondents into four categories: non-use, high-risk use (seven or more drinks in men or five or more drinks in women on the same occasion on at least 2 days within the past 7 days), moderate-risk use (seven or more drinks in men or five or more drinks in women on the same occasion on at least 1 day within the past 30 days), and low-risk use (some alcohol consumption but less than moderate-risk use).

For the comprehensive analysis of multiple factors associated with alcohol use, we selected socio-demographic variables, health behaviors, and health status including self-rated and mental health as independent variables. These variables have been associated with alcohol use in prior studies [6,20,23-26]. Socio-demographic variables included sex, age (19-44, 45-64, ≥ 65 years), marital status (never married, married/live with partner, formerly married [divorced/separated/widowed]), educational attainment (no formal education, elementary school, middle school, high school, college and above), monthly household income (1.49 million Korean won [KRW] or less, 1.50-2.99 million KRW, 3 million KRW or more), and current employment status (employed, unemployed). Health behavior variables were smoking status (current, former, non-smoking) and physical activity (participated in moderate physical activity for 5 days or more per week and for 30 minutes or more per activ-

ity or in vigorous activity for 3 days or more per week and for 20 minutes or more per activity). Health status-related characteristics included obese status (normal, body mass index [BMI] <25.0 kg/m²; overweight/obesity, BMI ≥25.0 kg/m²), self-rated health (good, fair, poor), and depressive symptoms evaluated by the Center for Epidemiological Studies Depression Scale (CES-D) (normal, CES-D score <21; high depressive symptoms, CES-D score ≥21).

Statistical Analysis

SAS version 9.2 (SAS Inc., Cary, NC, USA) was used for all statistical analysis; the analyses used sampling weights to account for the complex sampling design of the KCHS. The level of significance was set to 0.05. To control for potential sex-related differences in risk of alcohol use, sex-stratified analyses were conducted. We assessed the association of the alcohol use pattern with socio-demographic and health-related characteristics using chi-square tests. Multinomial logistic regression analyses were used to identify correlates of low-risk, moderate-risk, and high-risk use relative to non-use with adjustment for all variables.

RESULTS

Table 1 shows the characteristics of the study population. Of the population, 49.4% were men and 50.6% were women. Overall, 69.3% of the respondents were current drinkers (82.2% in men, 56.8% in women). The men had statistically significantly higher prevalence rates of moderate-risk use (15.5% vs. 8.2%) and of high-risk use (21.2% vs. 3.4%) relative to women (Table 1).

In both sexes, significant associations between alcohol use patterns and age group, marital status, educational attainment, monthly income, and employment status were found. Any alcohol use was more prevalent among individuals with higher educational level, higher income level, and employed status in both sexes. A higher prevalence of high-risk use was evident among middle-aged (45 to 64 years) and married men; higher prevalence rates of current drinking, low-risk, and moderate use were evident among younger aged and unmarried men. In contrast to men, in women, higher prevalence rates of drinking in any use pattern were evident among younger aged and unmarried persons (Table 2).

In both sexes, certain health-related characteristics were significantly associated with alcohol use patterns. In men, the

prevalence rates of high-risk use were higher among current smokers, persons who participated in physical activity, persons who were overweight or obese, those with relatively good self-rated health, and those with depressive symptoms. In women, the prevalence rates of high-risk use were higher among current smokers, persons who were not overweight, those with relatively good self-rated health, and those with depressive symptoms (Table 3).

The adjusted odds ratios (ORs) for low-, moderate-, and high-risk alcohol use relative to non-use are shown in Table 4. In both sexes, the ORs of persons who were younger, had a higher monthly income, were currently employed, were current or former smokers, and had good or fair self-rated health relative to their reference groups were significantly higher in low-, moderate-, and high-risk use. Also, in both sexes, being overweight or obese was associated with moderate- and high-risk use.

There were sex-specific associations for some characteristics. For the men, higher educational attainment was significantly associated with low- and moderate-risk use. The ORs of the men who were married or formerly married relative to the never married were lower in moderate-risk use, but higher in high-risk use. The OR of the men who participated in physical activity was higher in high-risk use relative to that for physically inactive men, and those with high depressive symptoms had lower odds of low- and moderate-risk use. For women, the ORs of low-, moderate-, and high-risk drinking of persons who were married or formerly married were significantly lower compared to those of never-married women. The ORs of low-, moderate-, and high-risk use of women who had graduated from middle and high school were higher than those of uneducated women. The ORs of moderate- and high-risk drinking of women with high depressive symptoms were higher than those of women without depressive symptoms.

DISCUSSION

Using a representative population-based sample, we examined the rates of alcohol use and factors associated with low-, moderate-, and high-risk use. Overall, 69.3% of the population (82.2% of men, 56.8% of women) reported alcohol use during the previous year, and the rate of high-risk drinking was 12.0% (21.2% of men, 3.4% of women). These results are to some degree different from those of some restricted area studies and similar national surveys. For example, in the 2009 Korea Na-

Table 1. Characteristics of the study population

Variables	Total		Men		Women		p-value
	Sample size	% (SE)	Sample size	% (SE)	Sample size	% (SE)	
Age (y)							
19-44	95 611	53.1 (0.1)	46 068	55.0 (0.2)	49 543	51.2 (0.2)	<0.001
45-64	83 419	33.4 (0.1)	39 988	33.8 (0.2)	43 431	33.0 (0.2)	
≥ 65	51 685	13.5 (0.1)	21 024	11.2 (0.1)	30 661	15.8 (0.1)	
Marital status							
Never married	36 271	22.6 (0.1)	20 892	27.1 (0.2)	15 379	18.2 (0.2)	<0.001
Married	153 806	63.8 (0.1)	75 961	65.4 (0.2)	77 845	62.2 (0.2)	
Formerly married	40 430	13.6 (0.1)	10 120	7.5 (0.1)	30 310	19.5 (0.1)	
Educational level							
Uneducated	28 535	6.6 (0.1)	6214	3.1 (0.1)	22 321	10.1 (0.1)	<0.001
Elementary school	37 289	10.3 (0.1)	14 756	7.9 (0.1)	22 533	12.8 (0.1)	
Middle school	27 395	10.1 (0.1)	13 406	9.6 (0.1)	13 989	10.6 (0.1)	
High school	80 110	39.9 (0.1)	41 340	42.4 (0.2)	41 340	37.5 (0.2)	
College and over	57 179	33.0 (0.2)	31 259	37.0 (0.2)	31 259	29.1 (0.2)	
Monthly income (×10 ⁶ KRW)							
≤ 1.49	67 486	20.2 (0.1)	28 063	17.9 (0.2)	39 423	22.5 (0.2)	<0.001
1.50-2.99	70 129	32.7 (0.2)	34 772	34.0 (0.2)	36 257	31.5 (0.2)	
≥ 3.00	83 900	47.1 (0.2)	40 412	48.1 (0.2)	43 488	46.0 (0.2)	
Employment status							
Unemployed	103 528	42.6 (0.1)	26 052	24.7 (0.2)	77 476	60.0 (0.2)	<0.001
Employed	126 094	57.4 (0.1)	80 341	75.3 (0.2)	45 753	40.0 (0.2)	
Smoking status							
Current smoking	54 779	25.7 (0.1)	50 144	48.1 (0.2)	4635	3.7 (0.1)	<0.001
Ex-smoking	31 168	13.0 (0.1)	29 135	24.6 (0.2)	2033	1.6 (0.0)	
Non-smoking	144 588	61.3 (0.1)	27 675	27.3 (0.2)	116 913	94.6 (0.1)	
Physical activity							
No	178 892	79.0 (0.1)	77 885	74.5 (0.2)	101 007	83.5 (0.1)	<0.001
Yes	51 603	21.0 (0.1)	29 093	25.5 (0.2)	22 510	16.5 (0.1)	
Body weight							
Normal	172 831	77.7 (0.1)	77 107	72.4 (0.2)	95 724	82.9 (0.1)	<0.001
Overweight/obese	50 698	22.3 (0.1)	28 331	27.6 (0.2)	22 367	17.1 (0.1)	
Self-rated health							
Good	96 580	46.4 (0.2)	50 205	50.8 (0.2)	46 375	42.1 (0.2)	<0.001
Fair	87 462	39.3 (0.1)	39 585	37.8 (0.2)	47 877	40.7 (0.2)	
Poor	46 594	14.3 (0.1)	17 248	11.3 (0.1)	29 346	17.2 (0.1)	
Depressive symptoms ¹							
Normal	215 257	93.9 (0.1)	102 191	94.6 (0.1)	113 066	90.3 (0.1)	<0.001
High	14 821	6.1 (0.1)	4589	5.4 (0.1)	10 232	9.7 (0.1)	
Alcohol use pattern							
Non-use	87 263	30.6 (0.1)	24 589	17.8 (0.1)	62 674	43.2 (0.2)	<0.001
Low-risk use	97 269	45.3 (0.1)	47 793	45.5 (0.2)	49 476	45.2 (0.2)	
Moderate-risk use	21 257	11.8 (0.1)	13 332	15.5 (0.1)	7925	8.2 (0.1)	
High-risk use	24 863	12.2 (0.1)	21 347	21.2 (0.2)	3516	3.4 (0.1)	

All percentages were weighted to represent the total population of the 2009 Community Health Survey.

SE, standard error; KRW, Korean won.

¹High depressive symptoms defined as a Center for Epidemiological Studies Depression Scale score of 21 or higher.

Table 2. Association between alcohol use patterns and socio-demographic characteristics according to sex

Variables	Men					Women				
	Non-use	Low-risk	Moderate-risk	High-risk	p-value	Non-use	Low-risk	Moderate-risk	High-risk	p-value
Age (y)										
19-44	11.6 (0.2)	45.6 (0.3)	20.8 (0.2)	22.0 (0.2)	<0.001	29.1 (0.2)	53.1 (0.3)	12.9 (0.2)	4.9 (0.1)	<0.001
45-64	19.7 (0.2)	45.5 (0.3)	10.8 (0.2)	24.0 (0.3)		49.1 (0.3)	44.0 (0.3)	4.5 (0.1)	2.4 (0.1)	
≥65	42.4 (0.5)	44.8 (0.3)	3.5 (0.2)	9.3 (0.3)		76.7 (0.3)	22.3 (0.3)	0.5 (0.1)	0.5 (0.1)	
Marital status										
Never married	13.8 (0.3)	46.3 (0.4)	22.9 (0.4)	17.0 (0.3)	<0.001	24.7 (0.4)	50.2 (0.5)	18.6 (0.4)	6.5 (0.2)	<0.001
Married	18.8 (0.2)	45.6 (0.2)	12.8 (0.2)	22.8 (0.2)		43.8 (0.2)	47.4 (0.2)	6.2 (0.1)	2.6 (0.1)	
Formerly married	24.0 (0.5)	41.5 (0.6)	11.8 (0.4)	22.7 (0.5)		58.6 (0.4)	33.6 (0.4)	4.7 (0.2)	3.1 (0.2)	
Educational level										
Uneducated	37.5 (0.8)	40.5 (0.9)	6.3 (0.5)	15.7 (0.7)	<0.001	72.2 (0.4)	25.4 (0.4)	1.4 (0.1)	1.0 (0.1)	<0.001
Elementary school	34.2 (0.5)	42.1 (0.6)	6.2 (0.3)	17.5 (0.5)		63.8 (0.4)	32.2 (0.4)	2.5 (0.1)	1.5 (0.1)	
Middle school	24.1 (0.5)	44.4 (0.6)	9.2 (0.3)	22.3 (0.5)		48.1 (0.5)	42.3 (0.5)	5.7 (0.2)	3.9 (0.1)	
High school	15.5 (0.2)	45.6 (0.3)	16.8 (0.2)	22.1 (0.3)		34.5 (0.3)	49.4 (0.3)	11.1 (0.2)	5.0 (0.1)	
College and over	13.6 (0.2)	46.8 (0.3)	18.4 (0.3)	21.2 (0.3)		33.5 (0.3)	53.5 (0.4)	10.2 (0.2)	2.8 (0.1)	
Monthly income (×10 ⁶ KRW)										
≤1.49	31.1 (0.4)	42.5 (0.4)	9.9 (0.3)	16.4 (0.3)	<0.001	59.6 (0.4)	32.5 (0.3)	5.1 (0.2)	2.8 (0.1)	<0.001
1.50-2.99	17.0 (0.3)	45.1 (0.4)	16.1 (0.3)	21.8 (0.3)		41.6 (0.3)	45.5 (0.3)	9.0 (0.2)	3.9 (0.2)	
≥3.00	13.2 (0.2)	46.7 (0.3)	17.2 (0.2)	22.9 (0.3)		35.7 (0.3)	51.6 (0.3)	9.3 (0.2)	3.4 (0.1)	
Employment status										
Unemployed	28.1 (0.3)	45.2 (0.4)	14.2 (0.3)	12.5 (0.3)	<0.001	50.8 (0.2)	41.1 (0.2)	5.9 (0.1)	2.2 (0.1)	<0.001
Employed	14.4 (0.1)	45.6 (0.2)	15.9 (0.2)	24.1 (0.2)		31.8 (0.3)	51.3 (0.3)	11.8 (0.2)	5.1 (0.1)	

Values are presented as % (standard error). All percentages were weighted to represent the total population of the 2009 Community Health Survey. KRW, Korean won.

Table 3. Associations between alcohol use patterns and health-related characteristics according to sex

Variables	Men					Women				
	Non-use	Low-risk	Moderate-risk	High-risk	p-value	Non-use	Low-risk	Moderate-risk	High-risk	p-value
Smoking status										
Current smoking	11.0 (0.2)	42.6 (0.3)	17.7 (0.2)	28.7 (0.3)	<0.001	29.4 (0.8)	34.5 (0.9)	16.1 (0.7)	20.0 (0.9)	<0.001
Ex-smoking	20.9 (0.3)	47.1 (0.4)	12.7 (0.3)	19.3 (0.3)		37.2 (1.3)	39.3 (1.5)	12.6 (1.0)	10.9 (0.9)	
Non-smoking	26.9 (0.3)	49.3 (0.4)	14.1 (0.3)	9.7 (0.2)		43.8 (0.2)	45.8 (0.2)	7.8 (0.1)	2.6 (0.1)	
Physical activity										
No	18.2 (0.2)	45.5 (0.2)	15.6 (0.2)	20.7 (0.2)	<0.001	43.9 (0.2)	44.7 (0.2)	8.0 (0.1)	3.4 (0.1)	<0.001
Yes	16.6 (0.3)	45.4 (0.4)	15.2 (0.3)	22.8 (0.3)		39.5 (0.4)	47.8 (0.4)	9.1 (0.3)	3.6 (0.2)	
Body weight										
Normal	18.4 (0.2)	47.4 (0.2)	14.7 (0.2)	19.5 (0.2)	<0.001	41.3 (0.2)	46.6 (0.2)	8.6 (0.1)	3.5 (0.1)	<0.001
Overweight/obese	15.8 (0.3)	40.8 (0.4)	17.6 (0.3)	25.8 (0.3)		49.1 (0.4)	40.8 (0.4)	7.0 (0.2)	3.1 (0.2)	
Self-rated health										
Good	15.6 (0.2)	47.0 (0.3)	17.2 (0.2)	20.2 (0.2)	<0.001	38.0 (0.3)	48.4 (0.3)	10.1 (0.2)	3.5 (0.1)	<0.001
Fair	14.9 (0.2)	46.2 (0.3)	15.5 (0.2)	23.4 (0.3)		38.7 (0.3)	49.0 (0.3)	8.4 (0.2)	3.9 (0.1)	
Poor	37.6 (0.5)	36.5 (0.5)	7.6 (0.3)	18.3 (0.4)		66.7 (0.4)	28.5 (0.4)	3.0 (0.2)	1.8 (0.1)	
Depressive symptoms ¹										
Normal	17.3 (0.2)	45.8 (0.2)	15.7 (0.2)	21.0 (0.2)	<0.001	42.6 (0.2)	46.0 (0.2)	8.1 (0.1)	3.2 (0.1)	<0.001
High	28.6 (0.8)	38.2 (0.9)	10.1 (0.6)	23.1 (0.8)		49.2 (0.7)	35.7 (0.6)	9.5 (0.4)	5.5 (0.3)	

Values are presented as % (standard error). All percentages were weighted to represent the total population of the 2009 Community Health Survey. ¹High depressive symptoms defined as a Center for Epidemiological Studies Depression Scale score of 21 or higher.

Table 4. Multinomial logistic regression results for the relationship between alcohol use patterns and socio-demographic and health-related characteristics in Korean adults

Variables	Men			Women		
	Low-risk vs. non-use	Moderate-risk vs. non-use	High-risk vs. non-use	Low-risk vs. non-use	Moderate-risk vs. non-use	High-risk vs. non-use
Age (≥ 65 y)						
19-44	2.08 (1.91, 2.26)	7.99 (6.87, 9.30)	4.76 (4.25, 5.32)	3.53 (3.28, 3.79)	28.9 (21.6, 38.7)	15.6 (11.2, 21.9)
45-64	1.27 (1.19, 1.36)	3.08 (2.68, 3.55)	2.69 (2.44, 2.96)	2.05 (1.93, 2.17)	8.62 (6.50, 11.4)	4.94 (3.60, 6.77)
Marital status (/never married)						
Married	0.96 (0.89, 1.03)	0.69 (0.63, 0.75)	1.15 (1.06, 1.26)	0.84 (0.79, 0.88)	0.39 (0.36, 0.42)	0.47 (0.42, 0.53)
Formerly married	0.91 (0.82, 1.01)	0.79 (0.70, 0.91)	1.22 (1.08, 1.37)	0.85 (0.79, 0.91)	0.53 (0.48, 0.60)	0.64 (0.54, 0.76)
Educational level (/uneducated)						
Elementary school	1.11 (1.00, 1.23)	1.27 (1.00, 1.61)	1.02 (0.88, 1.18)	1.01 (0.95, 1.09)	0.93 (0.73, 1.19)	0.98 (0.70, 1.37)
Middle school	1.23 (1.11, 1.38)	1.44 (1.14, 1.81)	1.13 (0.97, 1.31)	1.19 (1.10, 1.29)	1.36 (1.08, 1.71)	1.75 (1.25, 2.46)
High school	1.44 (1.29, 1.60)	2.04 (1.64, 2.55)	1.26 (1.08, 1.45)	1.30 (1.21, 1.41)	1.59 (1.28, 1.98)	1.56 (1.12, 2.16)
College and over	1.45 (1.30, 1.63)	2.18 (1.74, 2.73)	1.14 (0.98, 1.32)	1.16 (1.06, 1.26)	0.99 (0.79, 1.25)	0.64 (0.46, 0.90)
Monthly income (≤ 1.49) ($\times 10^6$ KRW)						
1.50-2.99	1.23 (1.15, 1.30)	1.43 (1.30, 1.56)	1.27 (1.18, 1.38)	1.21 (1.15, 1.27)	1.31 (1.19, 1.44)	1.34 (1.17, 1.53)
≥ 3.00	1.58 (1.48, 1.69)	1.90 (1.73, 2.08)	1.74 (1.60, 1.89)	1.47 (1.40, 1.54)	1.50 (1.36, 1.65)	1.53 (1.33, 1.76)
Employment (/unemployed)						
Employed	1.32 (1.25, 1.39)	1.30 (1.20, 1.40)	1.89 (1.75, 2.03)	1.44 (1.39, 1.50)	1.97 (1.85, 2.11)	2.47 (2.25, 2.72)
Smoking status (/non-smoking)						
Current smoking	2.28 (2.16, 2.41)	3.60 (3.34, 3.87)	7.16 (6.65, 7.71)	1.57 (1.41, 1.75)	4.00 (3.45, 4.65)	12.8 (11.0, 14.9)
Ex-smoking	1.75 (1.65, 1.85)	2.34 (2.15, 2.54)	3.46 (3.19, 3.76)	1.45 (1.26, 1.68)	3.06 (2.44, 3.84)	6.89 (5.45, 8.72)
Physical activity (/no)						
Yes	1.02 (0.97, 1.07)	1.01 (0.94, 1.08)	1.15 (1.08, 1.22)	1.15 (1.10, 1.20)	1.32 (1.22, 1.43)	1.26 (1.11, 1.42)
Body weight(/normal)						
Overweight/obese	0.92 (0.87, 0.97)	1.28 (1.20, 1.36)	1.35 (1.27, 1.43)	0.99 (0.94, 1.03)	1.22 (1.12, 1.33)	1.13 (1.00, 1.28)
Self, rated health (/poor)						
Good	1.88 (1.76, 2.00)	2.01 (1.80, 2.26)	1.50 (1.37, 1.63)	1.31 (1.24, 1.38)	1.61 (1.41, 1.83)	1.42 (1.19, 1.70)
Fair	2.10 (1.97, 2.24)	2.33 (2.08, 2.61)	1.84 (1.69, 1.99)	1.61 (1.53, 1.69)	1.89 (1.66, 2.15)	1.89 (1.59, 2.24)
Depressive symptoms (/normal) ¹						
High	0.82 (0.74, 0.91)	0.72 (0.62, 0.84)	1.11 (0.98, 1.25)	1.04 (0.98, 1.11)	1.61 (1.43, 1.82)	1.79 (1.53, 2.09)

Values are presented as adjusted odds ratio (95% confidence interval).

KRW, Korean won.

¹High depressive symptoms defined as a Center for Epidemiological Studies Depression Scale score of 21 or higher.

tional Health and Nutrition Examination Survey, which shared the same definition of yearly drinking and high-risk drinking with this study, the overall prevalence of past-year alcohol use among adults was slightly higher, at 75.7% (86.0% of men, 68.4% of women), and the rates of high-risk alcohol use were 24.6% in men and 7.3% in women [12]. These discrepancies may be due to differences between age and sex distributions of samples and weighting methods between the KCHS and KNHANES.

However, these rates of alcohol use in Korea are still higher than those in other countries, including China, the US, and

Brazil [6,24,25,27,28], and reflect cultural norms about drinking in Korea [29,30]. In Korea, alcohol use is widely considered to be an important means of social relationships, and it is accepted, or even encouraged, to drink during social interactions, accompanied by a meal, in order to enhance relations with friends or business partners, and to establish a happy and congenial mood for adults, especially men [29]. In this study, intake of alcohol at any level, including low-, moderate-, and high-risk drinking, was more prevalent among men than among women.

Furthermore, the Korean culture of drinking may explain the

association between alcohol drinking and socio-demographic factors including age, marital status, educational level, monthly income, and employment status. These findings are largely consistent with previous studies [7,23-27,31,32]. In men, the risk of high-risk drinking was higher at relatively younger ages, among the married or the formerly married, among those with higher monthly income, and among the employed. In Korea, persons with the above-mentioned factors are more likely to become involved in social relationships, which could increase the likelihood of high-risk drinking. One additional reason why high-risk drinking was associated with these factors among men is that alcohol drinking, even if it is high-risk drinking, has not been recognized widely among Koreans to be a health risk [4,27].

Unlike the other factors, marital status showed different results for the two levels of moderate- and high-risk drinking in this study. Married or formerly married men were less likely than never-married men to engage in moderate-risk drinking; however, they were more likely to engage in high-risk drinking. In contrast to men, women who were married were less likely to report high-risk drinking. The explanation of this result might be both social and biological. The married women's drinking may be discouraged because of the women's expected roles as housewives or mothers, including child delivery and rearing and maintaining relationships with their husband and family [33]. On the other hand, employment increased alcohol consumption among the women.

The associations between high-risk drinking and health behaviors including smoking and obesity found in this study are consistent with those found in previous studies [8,9,25]. Even though causal relationships between these factors cannot be determined, these factors are highly interrelated, and health education and campaigns should be used to publicize these problems more comprehensively [25].

Contrary to expectations, high-risk drinking was associated with some positive health behaviors and health status including physical activity and good perception of self-rated health. Interestingly, this result is consistent with previous studies [25,34], which showed that heavy drinkers perceived lower health risks related to drinking than persons who drank less than them.

In men, depressive symptoms were negatively associated with low- and moderate-risk drinking, but not high-risk drinking; in women, depressive symptoms were positively associated with moderate- and high-risk drinking. Several previous

studies have found that alcohol drinking was associated with poor mental health, including higher perceived stress and depressive symptoms [32-37]. These studies have suggested that high levels of stress or depressive symptoms were associated with alcohol use because alcohol was perceived to reduce tension in tension-producing circumstances [38]. In our study, the results for women were consistent with previous studies. However, the men with high levels of depressive symptoms were more likely to report non-use of alcohol than men with lower levels of depressive symptoms. Further research is needed to explain these gender differences and the associations between alcohol use and mental health among Koreans.

This study has several limitations. First, the data were self-reported. Health behaviors including alcohol drinking, smoking, and obesity may be underreported in surveys because of recall bias and social desirability bias [39,40]. Second, to ensure an adequate sample size for the KCHS, the replacement method was used when sampled households declined to participate in this survey, and this was not considered in the weight calculation for ease of computation. Third, the KCHS did not collect information from persons living in institutional settings, and so the data might not be representative of those populations. Fourth, the data were cross-sectional, which limits the ability to infer causal relations between alcohol drinking and the associated factors.

In conclusion, alcohol use, and especially high-risk drinking, is prevalent among Korean adults, which is a major public health concern. Culturally appropriate programs for specific at-risk groups (e.g., currently smoking employed men, never-married women who are employed) may be useful to reduce high-risk behaviors. However, given the large proportion of the population that is engaging in moderate-risk and high-risk drinking and given the social norms that support this behavior, public health policies and campaigns to reduce alcohol consumption targeting the entire population are indicated.

ACKNOWLEDGEMENTS

This study was supported by the research fund of Chosun University, 2012.

CONFLICT OF INTEREST

The authors have no conflicts of interest related to the material presented in this paper.

REFERENCES

1. World Health Organization. Public health problems caused by harmful use of alcohol, 2005 [cited 2013 July 23]. Available from; http://www.who.int/substance_abuse/activities/public_health_alcohol/en/index.html.
2. Rehm J, Mathers C, Popova S, Thavorncharoensap M, Teerawattananon Y, Patra J. Global burden of disease and injury and economic cost attributable to alcohol use and alcohol-use disorders. *Lancet* 2009;373(9682):2223-2233.
3. Korean Statistical Information Service. 2009 Results of Korean causes of death; 2010 [cited 2012 Aug 10]. Available from: http://kosis.kr/ups/ups_01List01.jsp?grp_no=&pubcode=YD&type=F (Korean).
4. Brewer RD, Swahn MH. Binge drinking and violence. *JAMA* 2005;294(5):616-618.
5. Rehm J, Room R, Graham K, Monteiro M, Gmel G, Sempos CT. The relationship of average volume of alcohol consumption and patterns of drinking to burden of disease: an overview. *Addiction* 2003;98(9):1209-1228.
6. Blazer DG, Wu LT. The epidemiology of at-risk and binge drinking among middle-aged and elderly community adults: National Survey on Drug Use and Health. *Am J Psychiatry* 2009; 166(10):1162-1169.
7. Balsa AI, Homer JF, Fleming MF, French MT. Alcohol consumption and health among elders. *Gerontologist* 2008;48(5):622-636.
8. Chen CM, Yi H, Falk DE, Stinson FS, Dawson DA. Alcohol use and alcohol use disorders in the United States: main findings from the 2001-2002 National Epidemiologic Survey on Alcohol and Related Conditions. Bethesda: National Institute of Health; 2006, p. 22-23.
9. Green CA, Polen MR. The health and health behaviors of people who do not drink alcohol. *Am J Prev Med* 2001;21(4):298-305.
10. Ruidavets JB, Ducimetière P, Evans A, Montaye M, Haas B, Bingham A, et al. Patterns of alcohol consumption and ischaemic heart disease in culturally divergent countries: the Prospective Epidemiological Study of Myocardial Infarction (PRIME). *BMJ* 2010;341:c6077.
11. Ministry of Health and Welfare; Korea Centers for Disease Control and Prevention. Korea health statistics 2009: Korea National Health and Nutrition Examination Survey (KNHANES IV-3). Seoul: Ministry of Health and Welfare; 2010, p. 22-23.
12. Hahm BJ, Cho MJ. Prevalence of alcohol use disorder in a South Korean community: changes in the pattern of prevalence over the past 15 years. *Soc Psychiatry Psychiatr Epidemiol* 2005;40(2):114-119.
13. Chun S, Reid EA, Yun M. The association of alcohol drinking pattern and self-inflicted intentional injury in Korea: a cross-sectional WHO collaborative emergency room study. *BMJ Open* 2013;3(4):e002469.
14. Jung SK, Kim MK, Shin J, Choi BY. A cross-sectional analysis of the relationship between daily alcohol consumption and serum adiponectin levels among adults aged 40 years or more in a rural area of Korea. *Eur J Clin Nutr* 2013. doi: 10.1038/ejcn.2013.74.
15. Jang H, Jang WM, Park JH, Oh J, Oh MK, Hwang SH, et al. Alcohol consumption and the risk of type 2 diabetes mellitus: effect modification by hypercholesterolemia: the Third Korea National Health and Nutrition Examination Survey (2005). *Asia Pac J Clin Nutr* 2012;21(4):588-593.
16. Kim MK, Ko MJ, Han JT. Alcohol consumption and mortality from all-cause and cancers among 1.34 million Koreans: the results from the Korea national health insurance corporation's health examinee cohort in 2000. *Cancer Causes Control* 2010; 21(12):2295-2302.
17. Kwon C, Nam CH, Shin JK, Kim GH, Lee MK, Ha EP. Drinking aspect of community residents and its related factors. *J Korean Alcohol Sci* 2002;3(1):36-52 (Korean).
18. Kim HR. Prevalence of alcohol consumption by socio-demographic factors in the Seoul metropolitan area. *J Korean Alcohol Sci* 2002;3(1):87-101 (Korean).
19. Park JT, Kim BG, Jhun HJ. Alcohol consumption and the CAGE questionnaire in Korean adults: results from the Second Korea National Health and Nutrition Examination Survey. *J Korean Med Sci* 2008;23(2):199-206.
20. Hwang TY, Lee JJ, Lee KS, Kang PS, Kim SK. Drinking and its relationship with smoking among university students. *J Agric Med Community Health* 2007;32(2):97-105 (Korean).
21. Baek DH, Hwang BD, Moon HJ, Yoon HJ, Lee SK. A study on health status of the elderly in rural community according to drinking pattern. *J Agric Med Community Health* 2006;31(3): 263-273 (Korean).
22. Kim YT, Choi BY, Lee KO, Kim H, Chun JH, Kim SY, et al. Overview of Korean community health survey. *J Korean Med Assoc* 2012;55(1):74-83 (Korean).
23. Naimi TS, Brewer RD, Mokdad A, Denny C, Serdula MK, Marks JS. Binge drinking among US adults. *JAMA* 2003;289(1):70-75.
24. Li Y, Jiang Y, Zhang M, Yin P, Wu F, Zhao W. Drinking behaviour

- among men and women in China: the 2007 China Chronic Disease and Risk Factor Surveillance. *Addiction* 2011;106(11):1946-1956.
25. Paul LA, Grubaugh AL, Frueh BC, Ellis C, Egede LE. Associations between binge and heavy drinking and health behaviors in a nationally representative sample. *Addict Behav* 2011;36(12):1240-1245.
26. Bryant AN, Kim G. Racial/ethnic differences in prevalence and correlates of binge drinking among older adults. *Aging Ment Health* 2012;16(2):208-217.
27. Kanny D, Liu Y, Brewer RD; Centers for Disease Control and Prevention (CDC). Binge drinking: United States, 2009. *MMWR Surveill Summ* 2011;60 Suppl:101-104.
28. Wolle CC, Sanches M, Zilberman ML, Caetano R, Zaleski M, Laranjeira RR, et al. Differences in drinking patterns between men and women in Brazil. *Rev Bras Psiquiatr* 2011;33(4):367-373.
29. Weatherspoon AJ, Park JY, Johnson RC. A family study of homeland Korean alcohol use. *Addict Behav* 2001;26(1):101-113.
30. Helzer JE, Canino GJ, Yeh EK, Bland RC, Lee CK, Hwu HG, et al. Alcoholism: North America and Asia. A comparison of population surveys with the Diagnostic Interview Schedule. *Arch Gen Psychiatry* 1990;47(4):313-319.
31. Serdula MK, Brewer RD, Gillespie C, Denny CH, Mokdad A. Trends in alcohol use and binge drinking, 1985-1999: results of a multi-state survey. *Am J Prev Med* 2004;26(4):294-298.
32. Satre DD, Gordon NP, Weisner C. Alcohol consumption, medical conditions, and health behavior in older adults. *Am J Health Behav* 2007;31(3):238-248.
33. Wilsnack RW, Vogeltanz ND, Wilsnack SC, Harris TR, Ahlström S, Bondy S, et al. Gender differences in alcohol consumption and adverse drinking consequences: cross-cultural patterns. *Addiction* 2000;95(2):251-265.
34. Slater MD, Basil MD, Maibach EW. A cluster analysis of alcohol-related attitudes and behaviors in the general population. *J Stud Alcohol* 1999;60(5):667-674.
35. Cooper ML, Frone MR, Russell M, Mudar P. Drinking to regulate positive and negative emotions: a motivational model of alcohol use. *J Pers Soc Psychol* 1995;69(5):990-1005.
36. McCreary DR, Sadava SW. Stress, alcohol use and alcohol-related problems: the influence of negative and positive affect in two cohorts of young adults. *J Stud Alcohol* 2000;61(3):466-474.
37. Sebena R, El Ansari W, Stock C, Orosova O, Mikolajczyk RT. Are perceived stress, depressive symptoms and religiosity associated with alcohol consumption? A survey of freshmen university students across five European countries. *Subst Abuse Treat Prev Policy* 2012;7(1):21.
38. Jones-Webb R, Jacobs DR Jr, Flack JM, Liu K. Relationships between depressive symptoms, anxiety, alcohol consumption, and blood pressure: results from the CARDIA Study. *Coronary Artery Risk Development in Young Adults Study. Alcohol Clin Exp Res* 1996;20(3):420-427.
39. Patrick DL, Cheadle A, Thompson DC, Diehr P, Koepsell T, Kinne S. The validity of self-reported smoking: a review and meta-analysis. *Am J Public Health* 1994;84(7):1086-1093.
40. Stockwell T, Donath S, Cooper-Stanbury M, Chikritzhs T, Catalano P, Mateo C. Under-reporting of alcohol consumption in household surveys: a comparison of quantity-frequency, graduated-frequency and recent recall. *Addiction* 2004;99(8):1024-1033.