

UC Irvine

UC Irvine Previously Published Works

Title

Too much of a good thing: immunodeficiency due to hyperactive PI3K signaling

Permalink

<https://escholarship.org/uc/item/3c77p8zz>

Journal

Journal of Clinical Investigation, 124(9)

ISSN

0021-9738

Authors

Walsh, Craig M

Fruman, David A

Publication Date

2014-09-02

DOI

10.1172/jci77198

Peer reviewed

Too much of a good thing: immunodeficiency due to hyperactive PI3K signaling

Craig M. Walsh and David A. Fruman

Institute for Immunology and Department of Molecular Biology and Biochemistry, University of California, Irvine, Irvine, California, USA.

Primary immune deficiency diseases arise due to heritable defects that often involve signaling molecules required for immune cell function. Typically, these genetic defects cause loss of gene function, resulting in primary immune deficiencies such as severe combined immune deficiency (SCID) and X-linked agammaglobulinemia (XLA); however, gain-of-function mutations may also promote immune deficiency. In this issue of the *JCI*, Deau et al. establish that gain-of-function mutations in *PIK3R1*, which encodes the p85 α regulatory subunit of class IA PI3Ks, lead to immunodeficiency. These observations are consistent with previous reports that hyperactivating mutations in *PIK3CD*, which encodes the p110 δ catalytic subunit, are capable of promoting immune deficiency. Mutations that reduce PI3K activity also result in defective lymphocyte development and function; therefore, these findings support the notion that too little or too much PI3K activity leads to immunodeficiency.

Hyperactive PI3K signaling and immune deficiency

PI3Ks were discovered by Cantley and colleagues in 1985 (1) and have since been found to play numerous roles in cells (2, 3). Through recruitment to growth factor and cytokine receptor intracellular signaling complexes, class I PI3Ks promote the phosphorylation of phosphatidylinositol-(4,5)-bisphosphate (PIP₂) to phosphatidylinositol-(3,4,5)-triphosphate (PIP₃), which activates several downstream signaling molecules via membrane recruitment to signaling complexes. Class IA PI3Ks are heterodimeric and composed of a catalytic subunit (p110 α , p110 β , and p110 δ) bound to a regulatory subunit (p85 α /p55 α /p50 α , p85 β , and p55 γ). Although p110 α and p110 β are ubiquitously expressed, p110 δ is restricted to immune cells (4). Indeed, loss of p110 δ through gene targeting substantially affects adaptive immunity, impairing B cell development and resulting in a profound loss of T and B cell activation and function (4). Deletion of

Pik3r1, which encodes PI3K p85 α , p55 α , and p50 α regulatory subunits, leads to defects in B cell development and function (5, 6), likely due to *Pik3r1*-independent class IA PI3K activity in T cells. PI3K is activated in both T and B cells via antigen receptor signaling, cytokine and chemokine binding, and surface integrin activation (3, 4). These extracellular stimuli depend on PI3K for coordination of downstream signaling enzymes, including the serine/threonine kinases PDK1, AKT, and mTOR. By influencing these pathways, PI3K affects a wide range of cellular functions, modulating cellular metabolism, gene expression, and posttranslational regulation of protein and organelle function. The myriad roles that PI3K plays in the immune system, especially in T and B lymphocytes, are widely documented (3, 7, 8).

In this issue, Deau et al. used whole-exome sequencing and identified two distinct *PIK3R1* heterozygous splice site mutations in four immunodeficient patients. The

two different mutations affected the same splice site, leading to aberrant splicing of *PIK3R1* exon 10 and resulting truncation of the p85 α protein (9). Similarly aberrant *PIK3R1* splicing has been observed in tumor cells and found to increase PI3K enzyme activity through loss of an inhibitory contact between the regulatory and catalytic subunits (10, 11). Consistent with this, Deau et al. found elevated PI3K activity in patient lymphocytes, with enhanced phosphorylation of AKT on serine 473 and threonine 308 (9). The patients did not present with lymphoproliferative symptoms, but instead exhibited enhanced lymphocyte activation-induced cell death (AICD). While overall lymphocyte counts were in the normal range, the naive fraction of CD4⁺ and CD8⁺ T cells was dramatically diminished. Some of the patients had elevated levels of transitional B cells, while others presented with a paucity of IgM IgD⁺ CD27⁺ memory B cells. Furthermore, these patients lacked IgA production, and most of them had much higher titers of IgM compared with IgG. These results suggest that these *PIK3R1* hyperactivating mutations lead to defects in lymphoid homeostasis and B cell Ig class switching.

The findings of Deau et al. (9) are reminiscent of previous observations made in patients bearing activating mutations in *PIK3CD*, which encodes p110 δ . In these individuals, the catalytic activity of p110 δ is dramatically increased as the result of an E1021K mutation (12). This activating mutation led to the development of a primary immune deficiency, termed “activated p110 δ syndrome” (APDS), a condition marked by susceptibility to respiratory tract infection, lymphopenia, and progressive airway damage. Upon closer inspection, Angulo et al. observed elevated serum IgM, an increased fraction of transitional B cells, and heightened AICD and AKT phosphorylation in APDS patient lymphocytes (12). Structural and biochemical evaluation of the E1021K-containing protein revealed an alteration in the C lobe of the kinase domain, which

► Related Article: p. 3923

Conflict of interest: The authors have declared that no conflict of interest exists.

Reference information: *J Clin Invest.* 2014;124(9):3688–3690. doi:10.1172/JCI77198.

interacts with lipid substrates in the cell membrane and is also involved in binding to the SRC homology 2 (SH2) domain of the regulatory subunits. Lucas et al. had similar findings in their characterization of 14 immune-deficient patients from seven families bearing distinct heterozygous germline gain-of-function mutations in *PIK3CD* (13). They found that three different gain-of-function *PIK3CD* mutations led to p110 δ hyperactivity, with heightened AKT phosphorylation and mTOR activity. Furthermore, individuals harboring gain-of-function *PIK3CD* mutations had an overrepresentation of senescent effector T cells, with enhanced glucose uptake and terminal effector differentiation. While not specifically addressed by Deau et al. (9), it is likely that the described *PIK3R1* mutation truncates the p85 α domain in a manner that alters its interaction with p110 δ , resulting in hyperactivation of the catalytic domain (11). Although the symptoms described in the APDS patients are somewhat distinct from those observed in the patients characterized by Deau and colleagues (9), it is possible that a greater concordance in immunodeficiency symptoms will become apparent with the identification of more patients bearing these activating mutations.

Conclusions and future directions

The implications of these findings are important in several respects. First, patients bearing gain-of-function mutations in *PIK3R1*, such as those described by Deau and colleagues (9), may benefit from inhibitors of the PI3K and mTOR pathways that are currently available in the clinic or under scrutiny in clinical trials (14). Indeed, the PI3K δ inhibitor IC87114 was effective in reducing the hyperactive AKT phosphorylation and heightened AICD in patient samples bearing the *PIK3R1* truncation mutation (9). PI3K δ inhibitors also affected similar parameters in patients bearing *PIK3CD* mutations (12, 13). Second, given that the identified mutations occur in the germline, it is somewhat surprising that such gain-of-function mutations did not result in an apparent increase in susceptibility to blood cancers. Previous work by Carrera and colleagues has established that a similar truncation in p85 α leads to the generation of the oncogene

“p65-PI3K,” a similarly dysregulated form of p85 α that promotes constitutive PI3K catalytic activity (10). Because PI3K and downstream pathways such as mTOR are vital for tumor cell growth and survival, PI3K hyperactivation might be expected to promote oncogenic transformation. Likewise, APDS patients do not appear to have higher susceptibility to cancer, although this may simply be due to the highly specific *PIK3CD* expression pattern (14). *PIK3R1* is more broadly expressed; therefore, it is possible that identification of additional patients bearing such mutations as those described by Deau et al. (9) may reveal a heightened susceptibility to oncogenic transformation in various tissues. Alternatively, the internal truncation caused by these splice site mutations may not increase PI3K catalytic activity as profoundly as was observed with p65-PI3K. It will be important to compare the catalytic activity of p110 catalytic subunits when paired with WT, p65-PI3K, and these exon 10 truncations of p85 α .

The work by Deau et al. lays the groundwork for future studies to evaluate the differential effects of PI3K signaling on lymphocyte biology and immune function. A key issue is how hyperactive PI3K signaling affects lymphocyte differentiation. In each study, hyperactive class IA PI3K signaling variably resulted in an enhancement of transitional B cells, with an increase in serum IgM and a consequent decrease in more mature IgG isoforms (9, 12, 13). These findings are intriguing in light of the observation that mice with B cells lacking PTEN, a phosphoinositide phosphatase that counteracts PI3K signaling, have impaired Ig class switch recombination and hyper-IgM syndrome (15). Treatment of these mice with p110 δ inhibitors rescued the Ig class switching phenotype. On the other hand, an immunodeficient patient lacking the p85 α subunit of PI3K had a profound defect in B cell development and Ig production (16). Together, these observations lend credence to the hypothesis that PI3K activity must be carefully tuned for proper Ig isotype class switching.

Hyperactive PI3K signaling also leads to enhanced cellular metabolism, which was observed in lymphocytes derived from mice bearing gain-of-function p110 δ (13). As lymphocytes are highly dependent on metabolic switching after activation (17), it may be that the defective

generation of more mature Ig isoforms and enhanced generation of transitional B cell populations associated with hyperactive PI3K is due to activating mutations in the class IA PI3K pathway reprogramming lymphocyte metabolism. Failure to properly control metabolic signaling can result in premature terminal differentiation and cell death (18); therefore, it may be that unrestricted activation of pathways downstream of class IA PI3K lead to the early death of lymphocytes in patients bearing *PIK3CD* and *PIK3R1* mutations. This may explain the observation of heightened AICD in T cells obtained from patients bearing the *PIK3CD* and *PIK3R1* mutations (9, 13). It will thus be important to carefully evaluate lymphocyte metabolic regulation in these patients. PI3K, mTOR, and metabolic regulation are all important for helper T cell differentiation (19, 20). Future studies should also characterize the effect of class IA PI3K gain-of-function mutations on distinct helper cell subsets, as deficiencies in distinct helper subsets could greatly affect immune function in these patients. It is also possible that hyperactive class IA PI3K signaling may lead to overactivation of immune self-tolerance mechanisms; thus, evaluation of regulatory T cells and antiinflammatory cytokine production will be interesting. Perhaps the salient message of these studies is that PI3K signaling does not act in a digital manner; context and signal levels clearly matter to the immune system (3).

Acknowledgments

C.M. Walsh is supported by grants from the National Multiple Sclerosis Society (CA 1058-A-8), the California Institute for Regenerative Medicine (CIRM RM1-01717 and TR3-05603), and the NIH (GM099040). D.A. Fruman is supported by grants from the NIH (CA158383 and AI099656).

Address correspondence to: Craig M. Walsh, 3215 McGaugh Hall, University of California, Irvine, Irvine, California 92697-3900, USA. Phone: 949.824.8487; E-mail: cwalsh@uci.edu. Or to: David A. Fruman, 3242 McGaugh Hall, University of California, Irvine, Irvine, California 92697-3900, USA. Phone: 949.824.1947; E-mail: dfruman@uci.edu.

- Whitman M, Kaplan DR, Schaffhausen B, Cantley L, Roberts TM. Association of phosphatidylinositol kinase activity with polyoma middle-T competent for transformation. *Nature*. 1985;315(6016):239–242.
- Deane JA, Fruman DA. Phosphoinositide 3-kinase: diverse roles in immune cell activation. *Annu Rev Immunol*. 2004;22:563–598.
- Fruman DA, Bismuth G. Fine tuning the immune response with PI3K. *Immunol Rev*. 2009;228(1):253–272.
- Okkenhaug K, Ali K, Vanhaesebroeck B. Antigen receptor signalling: a distinctive role for the p110delta isoform of PI3K. *Trends Immunol*. 2007;28(2):80–87.
- Fruman DA, et al. Impaired B cell development and proliferation in absence of phosphoinositide 3-kinase p85a. *Science*. 1999;283(5400):393–397.
- Suzuki H, et al. Xid-like immunodeficiency in mice with disruption of the p85a subunit of phosphoinositide 3-kinase. *Science*. 1999;283(5400):390–392.
- Okkenhaug K, Fruman DA. PI3Ks in lymphocyte signaling and development. *Curr Top Microbiol Immunol*. 2010;346:57–85.
- Vanhaesebroeck B, Guillermet-Guibert J, Graupera M, Bilanges B. The emerging mechanisms of isoform-specific PI3K signalling. *Nat Rev Mol Cell Biol*. 2010;11(5):329–341.
- Deau M-C, et al. A human immunodeficiency caused by mutations in the *PIK3R1* gene. *J Clin Invest*. 2014;124(9):3923–3928.
- Jimenez C, et al. Identification and characterization of a new oncogene derived from the regulatory subunit of phosphoinositide 3-kinase. *EMBO J*. 1998;17(3):743–753.
- Backer JM. The regulation of class IA PI 3-kinases by inter-subunit interactions. *Curr Top Microbiol Immunol*. 2010;346:87–114.
- Angulo I, et al. Phosphoinositide 3-kinase delta gene mutation predisposes to respiratory infection and airway damage. *Science*. 2013;342(6160):866–871.
- Lucas CL, et al. Dominant-activating germline mutations in the gene encoding the PI(3)K catalytic subunit p110delta result in T cell senescence and human immunodeficiency. *Nat Immunol*. 2014;15(1):88–97.
- Conley ME, Fruman DA. Genetics. Can cancer drugs treat immunodeficiency? *Science*. 2013;342(6160):814–815.
- Omori SA, et al. Regulation of class-switch recombination and plasma cell differentiation by phosphatidylinositol 3-kinase signaling. *Immunity*. 2006;25(4):545–557.
- Conley ME, et al. Agammaglobulinemia and absent B lineage cells in a patient lacking the p85a subunit of PI3K. *J Exp Med*. 2012;209(3):463–470.
- MacIver NJ, Michalek RD, Rathmell JC. Metabolic regulation of T lymphocytes. *Annu Rev Immunol*. 2013;31:259–283.
- Fracchia KM, Pai CY, Walsh CM. Modulation of T cell metabolism and function through calcium signaling. *Front Immunol*. 2013;4:324.
- Gerriets VA, Rathmell JC. Metabolic pathways in T cell fate and function. *Trends Immunol*. 2012;33(4):168–173.
- Waickman AT, Powell JD. mTOR, metabolism, and the regulation of T-cell differentiation and function. *Immunol Rev*. 2012;249(1):43–58.