

UCLA

UCLA Previously Published Works

Title

Molecular subclasses of clear cell ovarian carcinoma and their impact on disease behavior and outcomes

Permalink

<https://escholarship.org/uc/item/3m9399p3>

Journal

Clinical Cancer Research, 28(22)

ISSN

1078-0432

Authors

Bolton, Kelly L
Chen, Denise
de la Fuente, Rosario I Corona
et al.

Publication Date

2022-11-14

DOI

10.1158/1078-0432.ccr-21-3817

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Molecular Subclasses of Clear Cell Ovarian Carcinoma and Their Impact on Disease Behavior and Outcomes



Kelly L. Bolton¹, Denise Chen², Rosario Corona de la Fuente³, Zhuxuan Fu⁴, Rajmohan Murali⁵, Martin Köbel⁶, Yanis Tazi⁵, Julie M. Cunningham⁷, Irenaeus C.C. Chan¹, Brian J. Wiley¹, Lea A. Moukarzel⁵, Stacey J. Winham⁷, Sebastian M. Armasu⁷, Jenny Lester⁸, Esther Elishaev⁴, Angela Laslavic⁴, Catherine J. Kennedy^{9,10}, Anna Piskorz¹¹, Magdalena Sekowska¹¹, Alison H. Brand^{9,12}, Yoke-Eng Chiew^{9,10}, Paul Pharoah¹¹, Kevin M. Elias¹³, Ronny Drapkin¹⁴, Michael Churchman¹⁵, Charlie Gourley¹⁵, Anna DeFazio^{9,10,12,16}, Beth Karlan⁸, James D. Brenton¹¹, Britta Weigelt⁵, Michael S. Anglesio¹⁷, David Huntsman¹⁷, Simon Gayther⁵, Jason Konner⁵, Francesmary Modugno⁴, Kate Lawrenson³, Ellen L. Goode⁷, and Elli Papaemmanuil⁵

ABSTRACT

Purpose: To identify molecular subclasses of clear cell ovarian carcinoma (CCOC) and assess their impact on clinical presentation and outcomes.

Experimental Design: We profiled 421 primary CCOCs that passed quality control using a targeted deep sequencing panel of 163 putative CCOC driver genes and whole transcriptome sequencing of 211 of these tumors. Molecularly defined subgroups were identified and tested for association with clinical characteristics and overall survival.

Results: We detected a putative somatic driver mutation in at least one candidate gene in 95% (401/421) of CCOC tumors including *ARID1A* (in 49% of tumors), *PIK3CA* (49%), *TERT* (20%), and *TP53* (16%). Clustering of cancer driver mutations and RNA expression converged upon two distinct subclasses of CCOC. The first was dominated by *ARID1A*-mutated tumors with

enriched expression of canonical CCOC genes and markers of platinum resistance; the second was largely comprised of tumors with *TP53* mutations and enriched for the expression of genes involved in extracellular matrix organization and mesenchymal differentiation. Compared with the *ARID1A*-mutated group, women with *TP53*-mutated tumors were more likely to have advanced-stage disease, no antecedent history of endometriosis, and poorer survival, driven by their advanced stage at presentation. In women with *ARID1A*-mutated tumors, there was a trend toward a lower rate of response to first-line platinum-based therapy.

Conclusions: Our study suggests that CCOC consists of two distinct molecular subclasses with distinct clinical presentation and outcomes, with potential relevance to both traditional and experimental therapy responsiveness.

See related commentary by Lheureux, p. 4838

Introduction

Historically, tumor treatment approaches have been dictated by tissue site, but large-scale molecular profiling efforts have shown that remarkable heterogeneity exists in the landscape of cancer driver genes and pathways within tumor types and even within histologic subtypes. This has been well characterized for many common tumors through multi-omic profiling (1) and characterization of the genetic determinants of tumor behavior and outcome has led to the development of personalized therapeutic approaches. Indeed, for some cancers, prognosis and therapeutic strategies are based primarily on their presence

of genetic driver mutations identified in the tumor (2–7). For several rare cancer types such as ovarian clear cell carcinoma (CCOC), no strong associations between molecular profiles and clinical presentation or outcomes are known and broad-acting platinum-based chemotherapy remains the standard of care.

When diagnosed at an advanced stage, CCOC has a worse outcome than other invasive ovarian cancers including the more common high-grade serous ovarian carcinoma (HGSOC; median overall survival of 10 months; refs. 8, 9) presents at a younger age (10), and is less responsive to platinum-based therapy (11). Relatively small studies suggest that CCOC possesses several driver events that

¹Washington University School of Medicine, St. Louis, Missouri. ²Philadelphia College of Osteopathic Medicine, Philadelphia, Pennsylvania. ³Cedars-Sinai Medical Center, Los Angeles, California. ⁴University of Pittsburgh Graduate School of Public Health, Pittsburgh, Pennsylvania. ⁵Memorial Sloan-Kettering Cancer Center, New York, New York. ⁶The University of Calgary, Calgary, Alberta, Canada. ⁷Mayo Clinic, Rochester, Minnesota. ⁸David Geffen School of Medicine, Department of Obstetrics and Gynecology, University of California at Los Angeles, Los Angeles, California. ⁹Department of Gynaecological Oncology, Westmead Hospital, Sydney, New South Wales, Australia. ¹⁰Centre for Cancer Research, The Westmead Institute for Medical Research, Sydney, New South Wales, Australia. ¹¹University of Cambridge, Cambridge, United Kingdom. ¹²The University of Sydney, Sydney, New South Wales, Australia. ¹³Brigham and Women's Hospital, Boston, Massachusetts. ¹⁴University of Pennsylvania, Philadelphia, Pennsylvania. ¹⁵University of Edinburgh, Edinburgh, United Kingdom. ¹⁶The Daffodil Centre, The University of Sydney, a joint venture with Cancer Council NSW, Sydney, New

South Wales, Australia. ¹⁷University of British Columbia, Vancouver, British Columbia, Canada.

K.L. Bolton, D. Chen, R. Corona de la Fuente, Z. Fu, R. Murali, M. Köbel, Y. Tazi, J.M. Cunningham, L.A. Moukarzel, S. Gayther, J. Konner, F. Modugno, K. Lawrenson, E.L. Goode, and E. Papaemmanuil contributed equally to this article.

Corresponding Author: Kelly L. Bolton, Medicine, Washington University, 4444 Forest Park, St. Louis, MO 63108. E-mail: bolton@wustl.edu

Clin Cancer Res 2022;28:4947–56

doi: 10.1158/1078-0432.CCR-21-3817

This open access article is distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) license.

©2022 The Authors; Published by the American Association for Cancer Research

Translational Relevance

Clear cell ovarian cancer (CCOC) is the second most common subtype of epithelial ovarian cancer and when diagnosed at an advanced stage, has a poor prognosis. The relationship between molecular profiles and clinical presentation or outcomes is still unknown but could help guide the development of personalized therapeutic approaches for CCOC. Here, we profiled 421 primary CCOCs using deep targeted sequencing and whole transcriptome sequencing on a subset of 211. Clustering of cancer driver mutations and RNA expression converged upon two distinct subclasses of CCOC. The first was dominated by *ARID1A*-mutated tumors with enriched expression of canonical CCOC genes and markers of platinum resistance; the second was largely comprised of tumors with *TP53* mutations and enriched for the expression of genes involved in extracellular matrix organization and mesenchymal differentiation. These two distinct molecular subclasses showed distinct clinical presentation and outcomes, with potential relevance to therapeutic responsiveness.

are distinct from HGSOC. CCOC is thought to arise from endometriotic lesions with recurrent somatic mutations in *PIK3CA* and *ARID1A*, which are rare in HGSOC (12–15). In addition, the existing data suggests that CCOCs are commonly *TP53*-wild-type (whereas HGSOC ubiquitously harbors *TP53* mutations) and exhibits fewer structural rearrangements than HGSOC (13). However, it is not known whether clinically meaningful molecular subtypes of CCOC exist.

In the current study, we performed comprehensive targeted sequencing and transcriptomic profiling of a large, multi-ethnic cohort of 421 primary CCOCs to identify disease subclasses with distinct biology and clinical behavior, which in turn may provide avenues for personalized therapeutic approaches.

Materials and Methods

Study participants

Clinical data and therapy-naïve fresh frozen tumor material were utilized from women diagnosed with invasive CCOC and enrolled into research studies from the following sites: Memorial Sloan Kettering Cancer Center Gynecology Tissue Bank (MSK), Mayo Clinic (MAY), Addenbrooks Hospital (ADD), Cedars-Sinai Medical Center (WCP; Los Angeles, CA), University of Pittsburgh (PIT; Pittsburgh, PA), Gynaecological Oncology Biobank (GynBiobank) at Westmead Hospital (WMH, Sydney, Australia), University of Edinburgh (SCOT; Scotland), Canadian Ovarian Experimental Unified Resource (COEUR; multiple sites), Brigham and Women's Hospital (BWH; Boston, MA), and University of Pennsylvania (UPA; Philadelphia, PA). Participants provided written informed consent. The studies were conducted in accordance with recognized ethical guidelines (e.g., Declaration of Helsinki, CIOMS, Belmont Report, U.S. Common Rule), and approved by local institutional review boards. Extraction of DNA/RNA was performed centrally at MSK (for cases from MSK, WCP, PIT, BWH, and UPA) or locally (for cases from MAY, ADD, WMH, and COEUR). For the cases which were extracted centrally at MSK, slides from frozen tissue sections were reviewed by a pathologist (R. Murali) and extraction of DNA/RNA was performed from tumor sections, selected based on high content (>80%) of clear cell carcinoma. In total, tumors from 447 women diagnosed with CCOC were

analyzed. Race and menstruation status (pre vs. postmenopausal) was obtained through participant self-report. History of endometriosis was also obtained through self-report except at MSK where endometriosis was only available if mentioned on the pathology report. Tumor characteristics and clinical outcomes were obtained through medical record review.

Targeted DNA sequencing and analysis

We performed targeted sequencing of 163 putative CCOC driver genes (Supplementary Table S1) in DNA samples from the 447 tumor and blood-derived DNA from 16 unmatched controls using a custom Nimblegen capture-based panel. Genes were selected based on a combined analysis of 105 clear cell somatic sequencing studies including: (i) whole genome sequencing of 31 CCOCs from Wang and colleagues (13); (ii) whole-exome sequencing of eight cases from Jones and colleagues (12); (iii) targeted sequencing of 26 CCOCs using a panel of 465 known cancer drivers (MSK-IMPACT; ref. 16); and targeted or whole exome sequencing of 40 CCOCs from project GENIE (17). Included in our panel were 119 genes where somatic mutations have been identified in two or more CCOCs; 41 established cancer driver genes based on the COSMIC Cancer Gene Census (18) mutated in one CCOC and three genes in the SWI/SNF complex (*SMARCB1*, *SMARCC1*, *SMARCC2*) (14) that have been implicated in CCOC biology (Supplementary Table S1; ref. 19). We also included on the sequencing panel highly polymorphic single nucleotide variants distributed every 3 MB throughout the genome to capture large copy number deletions/amplifications.

Of 447 tumor samples, 421 (94%) passed quality control. As a technical set of normal samples (panel of normals), we included DNA extracted from the blood of 10 healthy, cancer-free individuals. Two tumor samples failed due to low coverage, 12 due to sample contamination and 12 due to duplication. The median sequencing coverage per sample was 539x. Raw sequence data were aligned to the human genome (NCBI build 37) using BWA (20). Variant calling for single nucleotide variants was performed using Mutect2 (21), Strelka (22), and CaVEMan (23) and for insertions/deletions using Pindel (24), Mutect2 (21), and Strelka (22). We considered mutations to be true if they: (i) passed at least two variant callers; (ii) were present at a variant allele fraction of greater than 2%; (iii) were present in gNOMAD (25) whole-exome sequencing data with a maximum population frequency of less than 0.001; (iv) had a variant allele frequency (VAF) at least two times greater than the median VAF in a panel of normal samples; and (v) were present in none of the panel of normal samples at a VAF of 2% or greater. We further excluded mutations in low complexity regions [DUST (26) score >7]. Mutations in known cancer hotspots that met all other requirements but failed due to low complexity or to only being passed by one variant caller were retained for consideration. We calculated a microsatellite instability score for each tumor using MSI sensor (27).

We used Bayesian Dirichlet processes to establish classification rules that partitioned tumors into subgroups, minimizing overlap between categories. The Dirichlet process defines an infinite prior distribution for the number and proportions of clusters in a mixture model, fitted with the use of the Markov chain Monte Carlo method (28). Our method was based on an implementation of the Dirichlet process mixture model available at <https://github.com/nicolaroberts/hdp> using a non-hierarchical Dirichlet process. We used 5,000 burnin iterations and subsequently sampled 10,000 realizations at intervals of 20 iterations. From this collection of data, we computed the optimal number of clusters, requiring that 90% of the samples were assigned a cluster.

Whole transcriptome sequencing and analysis

RNA sequencing (RNA-Seq) libraries were prepared for 211 cases from total RNA derived from the same tumor section using poly(A) enrichment of the mRNA. One hundred bp paired-end libraries were sequenced on Illumina's HiSeq at a targeted depth of 40 million reads per sample. We performed alignment using STAR (version STAR_2.5.1b; ref. 29) against the reference genome hg38 (GENCODE v26). Reads were summarized using featureCounts (version 1.5.0-p1; ref. 30). RNA clusters were defined using hierarchical clustering using the top 500 most variable protein coding genes (clustering parameters: method = ward.D2, distance = canberra). Differentially expressed genes between RNA cluster 1 and RNA cluster 2 samples were obtained using the R package DESeq2 (version 1.28.1; ref. 31) with collection site and RNA cluster as part of the design formula. Pathway enrichment analysis was performed using Metascape (version 3.5; ref. 4), looking for enrichment of GO and KEGG terms, Hallmark, Reactome and BioCarta Gene Sets, and Canonical Pathways. The top 500 most overexpressed genes in RNA cluster 1 (\log_2 fold change <1 and FDR <0.05) and the top 500 most overexpressed genes in RNA cluster 2 were used as input for Metascape (32).

Outcome analyses

Survival data was available for 350 cases. Survival time was calculated from the date of diagnosis to last follow-up and allowed for left truncation for cases who were consented following diagnosis. We right censored at five years from diagnosis to reduce non-ovarian cancer related deaths. Race, age at diagnosis (continuous and quadratic, assigned as site median for three cases), tumor stage, extent of residual disease, and study site were considered as covariates using a Cox proportional hazards model. Proportionality of hazards was examined using Schoenfeld residuals. In addition, contingency analysis was done on tumor mutational status and tumor cluster with primary treatment response (complete response or partial response compared to stable or progressive disease) stratified by tumor stage and vital status up to five years using a χ^2 test.

Data availability

The somatic variant calls and normalized RNA-seq intensity data, code, and deidentified clinical data are available here: https://github.com/kbolton-lab/Bolton_OCCC. This will enable all the figures and tables to be re-generated and also provide data for others for future analyses. We will also make the BAMs/FASTQs available to researchers through contacting Kelly Bolton (bolton@wustl.edu).

Results

Clinical characteristics

Key characteristics, other than race, of the 421 participants included in the study did not vary between study sites (Table 1). Compared with clinical characteristics reported in the literature for women with HGSO (10, 33), women with CCOC in this cohort were more likely to be of Asian ancestry (12% of individuals with non-missing race), have a history of endometriosis (13%), and present with early-stage disease (69%).

Targeted DNA sequencing of candidate CCOC driver genes

In 163 candidate CCOC driver genes we identified 6,361 mutations. Of these, 1,488 mutations were classified as potentially pathogenic based upon annotation in OncoKB (34), frequency in COSMIC, frequency in previously published CCOC sequencing data (12, 13, 16), predicted pathogenicity based on PolyPhen (35) and SIFT (36), and prior evidence in the literature (Supplementary Table S2). At least one putative driver mutation was identified in 401 of 421 tumors (95%) (mean number of mutations 3, range 1–25; Fig. 1A and C). The most commonly mutated genes were *ARID1A* (49%, $N = 205$), *PIK3CA* (45%, $N = 188$), and the *TERT* promoter (20%, $N = 84$). The most frequently recurrent mutations were clonally dominant with a VAF $>35\%$ (e.g., *ARID1A* and *TP53*) suggesting that they represented early events while others (e.g., *CREBBP*) were more often subclonal, possibly representing secondary events (Fig. 1B). We detected a higher proportion (16%, $N = 71$) of tumors with *TP53* mutations than has been

Table 1. Clinical characteristics of CCOC cases sequenced using targeted panel.

	ADD (N = 28)	BWH (N = 9)	COEUR (N = 181)	MAY (N = 38)	MSK (N = 60)	PIT (N = 24)	SCOT (N = 22)	UPA (N = 7)	WCP (N = 28)	WMH (N = 24)	Overall (N = 421)
Age (y)											
<40	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
40–50	1 (3.6%)	0 (0%)	37 (20.4%)	3 (7.9%)	6 (10.0%)	4 (16.7%)	4 (18.2%)	2 (28.6%)	7 (25.0%)	5 (20.8%)	69 (16.4%)
50–60	8 (28.6%)	2 (22.2%)	81 (44.8%)	16 (42.1)	28 (46.7)	9 (37.5%)	7 (31.8%)	2 (28.6%)	14 (50.0)	10 (41.7)	177 (42.0)
60–70	13 (46.4)	7 (77.8%)	48 (26.5%)	9 (23.7%)	19 (31.7)	5 (20.8%)	9 (40.9%)	2 (28.6%)	4 (14.3%)	5 (20.8%)	121 (28.7)
70–80	6 (21.4%)	0 (0%)	10 (5.5%)	7 (18.4%)	6 (10.0%)	4 (16.7%)	1 (4.5%)	1 (14.3%)	1 (3.6%)	3 (12.5%)	39 (9.3%)
≥80	0 (0%)	0 (0%)	1 (0.6%)	2 (5.3%)	0 (0%)	2 (8.3%)	1 (4.5%)	0 (0%)	0 (0%)	0 (0%)	6 (1.4%)
Missing	0 (0%)	0 (0%)	4 (2.2%)	1 (2.6%)	1 (1.7%)	0 (0%)	0 (0%)	0 (0%)	2 (7.1%)	1 (4.2%)	9 (2.1%)
Race											
White	16 (57.1)	9 (100%)	0 (0%)	38 (100%)	44 (73.3)	23 (95.8)	0 (0%)	6 (85.7%)	23 (82.1)	10 (41.7)	169 (40.1)
Asian	2 (7.1%)	0 (0%)	0 (0%)	0 (0%)	13 (21.7)	0 (0%)	0 (0%)	0 (0%)	4 (14.3%)	4 (16.7%)	23 (5.5%)
Black	0 (0%)	0 (0%)	0 (0%)	0 (0%)	1 (1.7%)	1 (4.2%)	0 (0%)	1 (14.3%)	1 (3.6%)	0 (0%)	4 (1.0%)
Other	0 (0%)	0 (0%)	0 (0%)	0 (0%)	2 (3.3%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	2 (0.5%)
Unknown	10 (35.7)	0 (0%)	181 (100%)	0 (0%)	0 (0%)	0 (0%)	22 (100%)	0 (0%)	0 (0%)	10 (41.7)	223 (53.0)
Endometriosis											
Yes	0 (0%)	0 (0%)	13 (7.2%)	10 (26.3)	6 (10.0%)	0 (0%)	2 (9.1%)	2 (28.6%)	7 (25.0%)	3 (12.5%)	43 (10.2%)
No	0 (0%)	9 (100%)	168 (92.8)	26 (68.4)	49 (81.7)	0 (0%)	20 (90.9)	5 (71.4%)	0 (0%)	0 (0%)	277 (65.8)
Unknown	28 (100%)	0 (0%)	0 (0%)	2 (5.3%)	5 (8.3%)	24 (100%)	0 (0%)	0 (0%)	21 (75.0)	21 (87.5)	101 (24.0)
FIGO stage											
I/II	17 (60.7)	7 (77.8%)	128 (70.7)	25 (65.8)	42 (70.0)	16 (66.7)	14 (63.6)	2 (28.6%)	15 (53.6)	16 (66.7)	282 (67.0)
III/IV	5 (17.9%)	2 (22.2%)	46 (25.4%)	12 (31.6)	17 (28.3)	8 (33.3%)	7 (31.8%)	5 (71.4%)	13 (46.4)	7 (29.2%)	122 (29.0)
Missing	6 (21.4%)	0 (0%)	7 (3.9%)	1 (2.6%)	1 (1.7%)	0 (0%)	1 (4.5%)	0 (0%)	0 (0%)	1 (4.2%)	17 (4.0%)

cancer fusion partners but have not been reported in CCOC before (Supplementary Fig. S3).

We evaluated mutation status with respect to clinical and epidemiological factors including age, race, tumor, and history of endometriosis. Compared with *ARID1A*-mutated tumors, patients with *KRAS* mutations were older at presentation (median age 53 vs. 67, $P = 0.03$; Fig. 2A). Individuals with a history of endometriosis were more likely to have *ARID1A*-mutated tumors (72% and 47% of patients with and without endometriosis respectively, $P = 2 \times 10^{-4}$; Fig. 2B). Advanced stage tumors were more likely to harbor *TP53* mutations than early-stage tumors (27% vs. 11% respectively, $P = 2 \times 10^{-4}$; Fig. 2C). Among *TP53* mutant tumors, a similar proportion (50% and 51%, respectively) were advanced stage with or without co-occurring *ARID1A* mutations. There was a trend toward a higher frequency of *ARID1A*-mutated tumors in women of east Asian descent but this was not significant (Fig. 2D).

We next examined the relationship between mutational burden, cancer driver genes, and patterns of genetic cooccurrence. Several genes harbored recurrent mutations within the same tumor (Supplementary Fig. S4). This seen for both tumor suppressor genes (e.g., *ARID1A*) and specific oncogenes including *PIK3R1* and *PIK3CA*. Among tumors with multiple *PIK3CA* mutations, variants were more likely to occur in nonhotspot locations within the gene (Supplementary Fig. S5; ref. 44). MSI sensor score was higher among individuals more than 10 driver mutations ($N = 12$, 3%) and among those with *MSH2* and *MSH6* mutations (Supplementary Fig. S6). We observed a statistically significant co-occurrence between mutations in *ARID1A*, *PIK3CA*, *TP53* and *BRCA1/BRCA2*. Mutual exclusivity between somatic mutations of *ARID1A*, *TP53*, *PIK3CA* and *PIK3R1* (Supplementary Fig. S7) suggests that these may represent distinct pathways to oncogenesis. The exclusivity

between *TP53* and *ARID1A* mutation was stronger in the setting of multiple *ARID1A* mutations (OR = 0.21; 95% CI, 0.07–0.54; $P = 2 \times 10^{-4}$) compared with a single *ARID1A* mutations (OR = 0.68; 95% CI, 0.32–1.34; $P = 0.28$). “We observed 54 mutations in genes known to be relevant to high penetrance genetic predisposition to ovarian cancer including *PMS2*, *MSH6*, *MSH2*, *BRCA1*, and *BRCA2*. Overall, 52% of these mutations were present at a VAF in the tumor of $\geq 35\%$. In the absence of matched normal tissue sequencing, we were not able to distinguish these from germline variants. Thus, it is possible that up to 26 cases (6% of the cohort) harbored a germline pathogenic variant in a known cancer susceptibility gene.”

Because we observed clear patterns of exclusivity and cooccurrence between gene drivers, we used unsupervised clustering approaches to define nonoverlapping subgroups of CCOC based on their mutational spectrum. We defined seven subgroups (Supplementary Fig. S8) and compared the frequency of mutations between subgroups. Four clusters were characterized by having an *ARID1A* mutation; the first cluster (cluster A) was characterized by a single *ARID1A* mutation in combination with another disease defining mutation (e.g., *PIK3CA*, *TERT*, *TP53*, *KRAS*, *PTEN*, *PPP2R1A*, *PIK3R1*, *CREBBP*, or *SPOP*; $N = 86$); the second (cluster B) with a single *ARID1A* mutation alone or in combination with non-disease defining mutation ($N = 19$); the third (cluster C) with multiple *ARID1A* mutations combined with a *PIK3CA* mutation ($N = 81$); and a fourth (cluster D) with multiple *ARID1A* mutations and *PIK3CA* wild-type ($N = 25$). Two clusters were *ARID1A* wildtype: Cluster E was defined by a *TP53* mutation ($N = 50$); and cluster F by other non-*TP53* disease-defining mutations ($N = 104$). A final cluster (cluster G) was characterized by mutations in *SMARCA4* ($N = 13$); a mutation typically observed in small cell ovarian carcinoma (23). The remaining tumors were undefined ($N = 57$).




Figure 2.

Frequency of somatic mutations by clinical characteristics including age at diagnosis (A), endometriosis (B), stage (C), and race (D). Genes that were mutated in at least 20 individuals with nonmissing values for the clinical characteristic were included. Shown are q -values (FDR corrected P values) based on Fisher exact test. *, $q < 0.05$; **, $q < 0.01$.

Similar to the patterns we observed when studying the association between individual mutations and clinical features, the *TP53*-mutated, *ARID1A* wild-type cluster showed an enrichment of advanced stage disease while tumors belonging to the *ARID1A*-mutant clusters were more likely in individuals of Asian ancestry and those with a history of endometriosis (Supplementary Fig. S9). Individuals in cluster G (*SMARCA4*-mutant tumors) had a nonsignificant trend towards a younger age at diagnosis ($P = 0.32$).

Transcriptomic profiling of CCOC

Transcriptomic profiles were generated for 212 CCOC tumors in which targeted sequencing was also performed. Using unsupervised clustering informed by expression of the 500 most variable genes, we identified two main RNA clusters (Supplementary Fig. S10): Expression cluster 1 showed higher expression of genes previously reported as highly expressed in CCOC including *ANXA4* and *GPX3*, both of which are linked to platinum resistance (45, 46). Among the most highly expressed genes in cluster 1 compared with 2 also included *GPX3* (47), which is known to be overexpressed in endometriosis compared to normal endometrial tissue, and *EEF1A2*, known to be overexpressed in CCOC associated endometriosis but not benign endometriosis (48). Genes that characterized this cluster were enriched in metabolic pathways including flavonoid glucuronidation ($P = 10^{-15}$) and monocarboxylic acid metabolism ($P = 10^{-13}$). Expression cluster 2 showed enriched expression of genes involved in extracellular matrix (ECM) organization ($P = 10^{-22}$) and mesenchymal differentiation, including genes such as *ADGR2* and *PDCH19* (Supplementary Fig. S10 and Fig. 3B). Compared to cluster 1, expression cluster 2 also showed higher expression of *WT1* and lower expression of CCOC marker *HNF1B*, which are features classically associated with high-grade serous ovarian cancer (Fig. 3B; ref. 9). Expression cluster 2 was enriched with *TP53*-mutant tumors (55% of cases in cluster 2 compared with 10% in cluster 1). When comparing RNA expression and mutation clusters, cluster 2 was largely comprised of tumors belonging to mutation cluster E, that is, *TP53*-mutant *ARID1A*-wild type tumors (45% of cluster 2) and the undefined mutation cluster (33% of cluster 2; Fig. 3A).

Clinical outcomes

There was no statistically significant association between overall survival and CCOC mutations when examined on a per-gene level in Cox proportional hazards models stratified by study site (Supplementary Table S3). We observed a nonsignificant trend toward improved survival for patients with *ARID1A* (HR = 0.82; 95% CI, 0.58–1.15; $P = 0.24$) and *PTEN* (HR = 0.52; 95% CI, 0.24–1.12; $P = 0.10$) mutant tumors. Because of the similarity of the *ARID1A*-mutant clusters in regards to clinical presentation and outcome, we combined these clusters for the purpose of survival analysis. Women with *TP53*-mutant, *ARID1A*-wild type tumors had worse overall survival compared to those with *ARID1A*-mutant tumors (HR = 1.72; 95% CI, 1.06–2.81; $P = 0.03$; Fig. 4A). Similarly, RNA-seq cluster 2 showed an increased risk of death compared with RNA-seq cluster 1 (Fig. 4B, Tumor Cluster 2 vs. Tumor Cluster 1 HR 2.8; 95% CI, 1.66–4.84; $P = 1 \times 10^{-4}$). Covariate adjustment for age, race, stage, and residual disease attenuated the estimated mutation and cluster-associated risk (Supplementary Table S4). To explore how these subgroups might influence therapy outcome, we studied the relationship between mutation status and response to first line therapy with platinum/taxane combination therapy. We limited this to women with advanced stage disease who successfully underwent debulking surgery followed by combination platinum/taxol therapy ($N = 36$). Women

with *ARID1A* wild-type, *TP53*-mutant tumors were more likely to have a complete response 75% ($N = 11$) compared to *ARID1A*-mutant tumors (55%), although this was not statistically significant ($P = 0.33$) in this small sample size.

Discussion

Our results have several clinical implications. First, the results of both genomic and transcriptomic cluster associations with clinical presentation and outcome converged, suggesting two main subgroups of CCOC: The first subtype included *ARID1A*-mutant tumors (particularly double-mutant tumors) and other common CCOC mutations (e.g., *PIK3CA*, *TERT*, etc.) that showed enriched expression of metabolic pathways, presented with early-stage disease and were more likely to have a history of endometriosis. We denote this group as “classic-CCOC”, which represented 83% of our cohort. The second CCOC subtype was dominated by *TP53*-mutant tumors that showed enriched expression of genes involved in extracellular matrix organization, mesenchymal differentiation and immune-related pathways. These cases presented with advanced disease and had worse survival. Interestingly, *TP53* mutations either in the presence or absence of cooccurring *ARID1A* mutations were associated with a higher degree of genomic instability and aggressive, advanced stage tumors. The worse survival for tumors in this “HGSOC-like” subgroup was largely explained by advanced stage and higher burdens of residual disease.

Within both the “classic-CCOC” and “HGSOC-like” subgroups we noted a subset of individuals had tumor with mutations in genes known to be both somatic drivers of ovarian cancer and germline susceptibility genes including *PMS2*, *MSH6*, *MSH2*, *BRCA1*, and *BRCA2*. Due to the absence of matched normal samples, we were unable to fully distinguish whether these represented somatic or germline events and is a limitation of our study. Future studies estimating the frequency of CCOC cases that arise in women with strong hereditary predisposition and who may be considered for risk reducing bilateral salpingo-oophorectomy should be prioritized (49).

There is increasing recognition that other histologic types of ovarian carcinoma, including HGSOC and endometrioid carcinoma, can contain areas with clear cell change complicating the histologic diagnosis (50). While a subset of cases in the “HGSOC-like” cluster are misclassified HGSOC, and is a weakness of our study, it is unlikely that this alone explains our findings. Firstly, all of our cases were morphologically diagnosed by expert gynecological pathologists and at some centers, this morphologic review was supplemented by immunohistochemistry for histotype-specific markers. Secondly, in a subset of *TP53*-mutant cases, we reconfirmed the diagnosis of CCOC using a combination of morphologic and immunohistochemical features. Thus, our results suggest that a subset of *bona fide* CCOCs with HGSOC-like features exist. Our results also emphasize that expert histologic review of CCOC cases, particularly those who present with *TP53*-mutant, *ARID1A*-wild type tumors, is warranted given similarities to the biology and behavior of HGSOC.

Gene expression profiles of the “classic-CCOC” and “HGSOC-like” CCOC subtypes we observed are similar to those reported by Tan and colleagues (51) which also reported two clusters, the first enriched for genes in metabolic pathways and the second, a less common mesenchymal-like subgroup associated with late-stage disease. However, unlikely Tan and colleagues, we observed differences in the frequency of *TP53*-mutated tumors across clusters. The source of this discrepancy is unclear and may include differences in sequencing technology (Tan and colleagues performed targeted sequencing using Ion Torrent) and patient characteristics (Tan and colleagues, included only





Figure 3. The transcriptome of clear cell ovarian cancer samples. **A**, Sankey plot showing the correspondence of the sample annotations RNA clusters and DNA clusters. **B**, Heatmap showing the normalized gene expression of the top 50 most differentially expressed genes between RNA cluster 1 and RNA cluster 2.

women of Asian ancestry which trend towards lower frequencies of *TP53*-mutated tumors in our analysis and which are known to have lower frequencies of endometrial ovarian cancer). The overlap between genes highly expressed in our “classic-CCOC” subgroup and those

enriched in endometriosis provide further support for the likely transition from endometriosis to carcinoma in CCOC.

The greatest translational impact from these molecular CCOC subtypes is expected to lie in the development of therapeutic

**Figure 4.**

Association between CCOC molecular subgroups and all-cause mortality. Shown are the Kaplan–Meier plots for the survival probability over 5 years following CCOC diagnosis stratified by mutational clusters defined by ARID1A/TP53 mutation status (**A**) and RNA-seq expression clusters (**B**).

approaches tailored to the vulnerabilities of each group. Interestingly, despite being aggressive on presentation, a trend was seen towards the “HGSOC-like” CCOC subgroup having higher response rates to first line platinum-based chemotherapy. Future studies are warranted to further explore whether genomic subtypes of CCOC predict response to platinum-based and other therapies as treatment data were limited here. The “classic-CCOC” subgroup dominated by mutations in the SWI/SNF pathway and markers linked to chemo-resistance may be of particular relevance to target for investigational first-line therapies. Recent data suggests that the SWI/SNF pathway plays a novel role in the regulation of antitumor immunity, and that SWI/SNF deficiency can be therapeutically targeted by immune checkpoint blockade (19). Several studies are currently evaluating the role of immune check point inhibitors in CCOC including NCT03405454, NCT03425565. While a limitation of our study was that we were unable to assess MMR functional status, we did note a rare subset of tumors (3%) with higher mutational burden (>10 drivers) and MSI sensor score. The extent to which the subset of CCOCs with higher total mutation and with MMR deficiency show improved responsiveness to immune checkpoint blockade in ongoing clinical trials will be an important avenue of investigation. Additional targeted therapeutic strategies have been explored in preclinical settings including epigenetic synthetic lethality, some of which are entering into clinical trials. The PI3K inhibitor, alpelisib, is now FDA approved for HR-positive breast cancer and ongoing trials in additional *PIK3CA*-mutated cancers including CCOC are underway. Double *PIK3CA* mutations appear to hyperactivate PI3K signaling and enhance tumor growth and may confer increased responsiveness to PI3K inhibitors than those with a single mutation (52). Thus, for CCOC cases harboring multiple *PIK3CA* mutations, PI3K inhibitors either alone or in combination with other agents may represent a promising approach.

The strengths of this study include the large sample size, use of multiple study sites, inclusion of women of European and non-European ancestry, and integration of genetic and transcriptomic markers of disease behavior and outcome. While this is the most extensive genomic study of CCOC to date, greater sample size with

additional follow-up data will allow improved assessment and validation of these clinically relevant subtypes. Although future analyses would benefit from larger patient collections, our current results suggest that genomic classification may inform the future development of targeted therapeutics in CCOC.

Authors' Disclosures

C.J. Kennedy reports grants from National Health and Medical Research Council and Cancer Institute New South Wales during the conduct of the study. Y.-E. Chiew reports grants from National Health and Medical Research Council of Australia and The Cancer Institute New South Wales during the conduct of the study. P. Pharoah reports grants from Cancer Research UK during the conduct of the study. R. Drapkin reports personal fees from Repare Therapeutics and Cedilla Therapeutics and other support from VOC Health outside the submitted work. C. Gourley reports grants and personal fees from AstraZeneca, MSD, GSK, and Nucana; personal fees from Clovis, Foundation One, Chugai, Cor2Ed, and Takeda; and grants from Novartis, Aprea, BerGenBio, and Medannexin outside the submitted work. A. DeFazio reports grants from National Health and Medical Research Council of Australia and The Cancer Institute NSW during the conduct of the study, as well as grants and other support from AstraZeneca outside the submitted work. B. Karlan reports grants from American Cancer Society during the conduct of the study as well as relationships with AstraZeneca (investigational therapeutic), Merck (investigational therapeutic), and Amgen (investigational therapeutic). J.D. Brenton reports grants from Cancer Research UK during the conduct of the study as well as personal fees from GSK and AstraZeneca and other support from Tailor Bio outside the submitted work. B. Weigelt reports personal fees from Repare Therapeutics outside the submitted work. D. Huntsman reports being Founder and CMO for Canexia Health. J. Konner reports personal fees from AstraZeneca, Clovis, and Tesaro outside the submitted work. F. Modugno reports grants from University of Pittsburgh during the conduct of the study. E. Papaemmanuil reports other support from Isabl Inc outside the submitted work. No disclosures were reported by the other authors.

Authors' Contributions

K.L. Bolton: Conceptualization, resources, data curation, formal analysis, funding acquisition, writing—original draft, writing—review and editing. **D. Chen:** Resources, data curation, investigation, writing—original draft, project administration, writing—review and editing. **R. Corona de la Fuente:** Formal analysis, writing—original draft, writing—review and editing. **Z. Fu:** Data curation, writing—original draft, writing—review and editing. **R. Murali:** Data curation, validation, investigation, writing—

original draft, writing–review and editing. **M. Köbel**: Data curation, investigation, methodology, writing–original draft, writing–review and editing. **Y. Tazi**: Formal analysis, writing–original draft, writing–review and editing. **J.M. Cunningham**: Writing–original draft, writing–review and editing. **I.C.C. Chan**: Formal analysis, writing–original draft, writing–review and editing. **B.J. Wiley**: Formal analysis, writing–original draft, writing–review and editing. **L.A. Moukarzel**: Data curation, writing–original draft, writing–review and editing. **S.J. Winham**: Formal analysis, writing–original draft. **S.M. Armasu**: Formal analysis, writing–original draft. **J. Lester**: Resources, data curation, writing–original draft. **E. Elishaev**: Resources, data curation, writing–original draft. **A. Laslavic**: Resources, writing–original draft. **C.J. Kennedy**: Resources, writing–original draft. **A. Piskorz**: Resources, writing–original draft. **M. Sekowska**: Data curation, writing–original draft. **A.H. Brand**: Data curation, writing–original draft. **Y.-E. Chiew**: Data curation, writing–original draft. **P. Pharoah**: Conceptualization, data curation, writing–original draft, writing–review and editing. **K.M. Elias**: Data curation, writing–original draft. **R. Drapkin**: Data curation, writing–original draft. **M. Churchman**: Data curation, writing–original draft. **C. Gourley**: Resources, data curation, writing–original draft. **A. DeFazio**: Resources, data curation, writing–original draft. **B. Karlan**: Resources, data curation, supervision, writing–original draft. **J.D. Brenton**: Resources, data curation, supervision, writing–original draft. **B. Weigelt**: Resources, data curation, supervision, investigation, writing–original draft. **M.S. Anglesio**: Resources, data curation, supervision, writing–original draft. **D. Huntsman**: Resources, data curation, writing–original draft. **S. Gayther**: Conceptualization, resources, data curation, supervision, investigation, writing–original draft, writing–review and editing. **J. Konner**: Conceptualization, resources, supervision, funding acquisition, writing–original draft, project administration, writing–review and editing. **F. Modugno**: Conceptualization, resources, data curation, supervision, writing–original draft, project administration, writing–review and editing. **K. Lawrenson**: Conceptualization, resources, data curation, supervision, visualization, methodology, writing–original draft, project administration, writing–review and editing. **E.L. Goode**: Conceptualization, resources, data curation, supervision, investigation, writing–original draft, project administration, writing–review and editing. **E. Papaemmanuil**: Conceptualization, resources, supervision, funding acquisition, methodology, writing–original draft, writing–review and editing.

Acknowledgments

Research reported in this publication was supported in part by a Cancer Center Support Grant of the NIH/NCI (Grant No. P30CA008748, MSK) and the Cycle for

Survival including the Fatma Fund. B. Weigelt is funded in part by Breast Cancer Research Foundation and NIH/NCI (P50 CA247749 01) grants. K.L. Bolton is funded by the Damon Runyon Cancer Research Foundation, the American Society of Hematology, the Evans MDS Foundation and the NCI (Grant 5K08CA241318). Additional support was provided by R21CA222867, R01CA248288, P30CA015083, and P50CA136393. M.S. Anglesio was funded through a Michael Smith Health Research BC Scholar Program award and the Janet D. Cottrelle Foundation Scholars Program (managed by the BC Cancer Foundation). This study used resources provided by the Canadian Ovarian Cancer Research Consortium's COEUR biobank funded by the Terry Fox Research Institute and managed and supervised by the Centre hospitalier de l'Université de Montréal. The Consortium acknowledges contributions of its COEUR biobank from Institutions across Canada (for a full list see <https://www.tfri.ca/coeur>). This work was supported by the Westmead Hospital Department of Gynaecological Oncology, Sydney Australia. The Gynaecological Oncology Biobank at Westmead (GynBiobank), a member of the Australasian Biospecimen Network-Oncology group, was funded by the National Health and Medical Research Council of Australia (Enabling Grants ID 310670 & ID 628903) and the Cancer Institute NSW (Grants 12/RIG/1–17 & 15/RIG/1–16). The Westmead GynBiobank acknowledges financial support from the Sydney West Translational Cancer Research Centre, funded by the Cancer Institute NSW. A. Piskorz, M. Sekowska, and J.D. Brenton were supported by Cancer Research UK grant 22905. Additional support was also provided by the National Institute of Health Research (NIHR) Cambridge Biomedical Research Centre (BRC-1215–20014). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, or the Department of Health and Social Care.

The publication costs of this article were defrayed in part by the payment of publication fees. Therefore, and solely to indicate this fact, this article is hereby marked “advertisement” in accordance with 18 USC section 1734.

Note

Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

Received October 25, 2021; revised February 24, 2022; accepted July 7, 2022; published first July 11, 2022.

References

- Bailey MH, Tokheim C, Porta-Pardo E, Sengupta S, Bertrand D, Weerasinghe A, et al. Comprehensive characterization of cancer driver genes and mutations. *Cell* 2018;173:371–85.
- Dong J, Li B, Lin D, Zhou Q, Huang D. Advances in targeted therapy and immunotherapy for non-small cell lung cancer based on accurate molecular typing. *Front Pharmacol* 2019;10:230.
- Quintás-Cardama A, Kantarjian H, Cortes J. Imatinib and beyond—exploring the full potential of targeted therapy for CML. *Nat Rev Clin Oncol* 2009;6:535–43.
- Ortega J, Vigil CE, Chodkiewicz C. Current progress in targeted therapy for colorectal cancer. *Cancer Control* 2010;17:7–15.
- Kayser S, Levis MJ. Advances in targeted therapy for acute myeloid leukaemia. *Br J Haematol* 2018;180:484–500.
- Tangutoori S, Baldwin P, Sridhar S. A new era of targeted therapy. *Maturitas* 2015;81:5–9.
- Vaishnavi A, Le AT, Doebele RC. TRKing down an old oncogene in a new era of targeted therapy. *Cancer Discov* 2015;5:25–34.
- Lee Y-Y, Kim T-J, Kim M-J, Kim H-J, Song T, Kim MK, et al. Prognosis of ovarian clear cell carcinoma compared to other histological subtypes: a meta-analysis. *Gynecol Oncol* 2011;122:541–7.
- Irodi A, Rye T, Herbert K, Churchman M, Bartos C, Mackean M, et al. Patterns of clinicopathological features and outcome in epithelial ovarian cancer patients: 35 years of prospectively collected data. *BJOG* 2020;127:1409–20.
- Wentzensen N, Poole EM, Trabert B, White E, Arslan AA, Patel AV, et al. Ovarian cancer risk factors by histologic subtype: an analysis from the ovarian cancer cohort consortium. *J Clin Oncol* 2016;34:2888–98.
- Iida Y, Okamoto A, Hollis RL, Gourley C, Herrington CS. Clear cell carcinoma of the ovary: a clinical and molecular perspective. *Int J Gynecol Cancer* 2021;31:605–16.
- Jones S, Wang T-L, Shih I-M, Mao T-L, Nakayama K, Roden R, et al. Frequent mutations of chromatin remodeling gene ARID1A in ovarian clear cell carcinoma. *Science* 2010;330:228–31.
- Wang YK, Bashashati A, Anglesio MS, Cochrane DR, Grewal DS, Ha G, et al. Genomic consequences of aberrant DNA repair mechanisms stratify ovarian cancer histotypes. *Nat Genet* 2017;49:856–65.
- Anglesio MS, Bashashati A, Wang YK, Senz J, Ha G, Yang W, et al. Multifocal endometriotic lesions associated with cancer are clonal and carry a high mutation burden. *J Pathol* 2015;236:201–9.
- Murakami R, Matsumura N, Brown JB, Higasa K, Tsutsumi T, Kamada M, et al. Exome sequencing landscape analysis in ovarian clear cell carcinoma shed light on key chromosomal regions and mutation gene networks. *Am J Pathol* 2017;187:2246–58.
- Cheng DT, Mitchell TN, Zehir A, Shah RH, Benayed R, Syed A, et al. Memorial Sloan Kettering-integrated mutation profiling of actionable cancer targets (MSK-IMPACT): A hybridization capture-based next-generation sequencing clinical assay for solid tumor molecular oncology. *J Mol Diagn* 2015;17:251–64.
- André F, Arnedos M, Baras AS, Baselga J, Bedard PL, Berger MF, et al. AACR Project GENIE: powering precision medicine through an international consortium. *Cancer Discov* 2017;7:818–31.
- Sondka Z, Bamford S, Cole CG, Ward SA, Dunham I, Forbes SA. The COSMIC cancer gene census: describing genetic dysfunction across all human cancers. *Nat Rev Cancer* 2018;18:696–705.

19. Fukumoto T, Magno E, Zhang R. SWI/SNF complexes in ovarian cancer: mechanistic insights and therapeutic implications. *Mol Cancer Res* 2018;16:1819–25.
20. Li H, Durbin R. Fast and accurate short read alignment with burrows-wheeler transform. *Bioinformatics* 2009;25:1754–60.
21. Benjamin D. Calling somatic SNVs and indels with Mutect2. *bioRxiv* 2019; 861054.
22. Saunders CT, Wong WSW, Swamy S, Becq J, Murray LJ, Cheatham RK, Strelka: accurate somatic small-variant calling from sequenced tumor-normal sample pairs. *Bioinformatics* 2012;28:1811–7.
23. Jones D, Raine KM, Davies H, Tarpey PS, Butler AP, Teague JW, et al. cgpCaVEManWrapper: simple execution of CaVEMan in order to detect somatic single nucleotide variants in NGS data. *Curr Protoc Bioinformatics* 2016;56:15.10.1–15.10.18.
24. Ye K, Schulz MH, Long Q, Apweiler R, Ning Z. Pindel: a pattern growth approach to detect break points of large deletions and medium sized insertions from paired-end short reads. *Bioinformatics* 2009;25:2865–71.
25. Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alfoldi J, Wang Q, et al. The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature* 2020;581:434–43.
26. Morgulis A, Gertz EM, Schäffer AA, Agarwala R. A fast and symmetric DUST implementation to mask low-complexity DNA sequences. *J Comput Biol* 2006; 13:1028–40.
27. Niu B, Ye K, Zhang Q, Lu C, Xie M, McLellan MD, et al. MSIsensor: microsatellite instability detection using paired tumor-normal sequence data. *Bioinformatics* 2014;30:1015–6.
28. Teh YW, Jordan MI, Beal MJ, Blei DM. Hierarchical Dirichlet Processes. *J Am Stat Assoc* 2006;101:1566–81.
29. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 2013;29:15–21.
30. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* 2014;30:923–30.
31. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014;15:550.
32. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al. Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nat Commun* 2019;10:1523.
33. Lisio M-A, Fu L, Goyeneche A, Gao Z-H, Telleria C. High-grade serous ovarian cancer: basic sciences, clinical and therapeutic standpoints. *Int J Mol Sci* 2019;20:952.
34. Chakravarty D, Gao J, Phillips S, Kundra R, Zhang H, Wang J, et al. OncoKB: a precision oncology knowledge base. *JCO Precis Oncol* 2017;2017:PO.17.00011.
35. Adzhubei I, Jordan DM, Sunyaev SR. Predicting functional effect of human missense mutations using PolyPhen-2. *Curr Protoc Hum Genet* 2013;Chapter 7: Unit7.20.
36. Ng PC. SIFT: predicting amino acid changes that affect protein function. *Nucleic Acids Res* 2003;31:3812–4.
37. Kuo K-T, Mao T-L, Jones S, Veras E, Ayhan A, Wang T-L, et al. Frequent activating mutations of PIK3CA in ovarian clear cell carcinoma. *Am J Pathol* 2009;174:1597–601.
38. Friedlander ML, Russell K, Millis S, Gatalica Z, Bender R, Voss A. Molecular profiling of clear cell ovarian cancers. *Int J Gynecol Cancer* 2016;26:648–54.
39. Köbel M, Rahimi K, Rambau PF, Naugler C, Le Page C, Meunier L, et al. An immunohistochemical algorithm for ovarian carcinoma typing. *Int J Gynecol Pathol* 2016;35:430–41.
40. Jelinic P, Mueller JJ, Olvera N, Dao F, Scott SN, Shah R, et al. Recurrent SMARCA4 mutations in small cell carcinoma of the ovary. *Nat Genet* 2014; 46:424–6.
41. Ramos P, Karnezis AN, Craig DW, Sekulic A, Russell ML, Hendricks WPD, et al. Small cell carcinoma of the ovary, hypercalcemic type, displays frequent inactivating germline and somatic mutations in SMARCA4. *Nat Genet* 2014; 46:427–9.
42. Witkowski L, Carrot-Zhang J, Albrecht S, Fahiminiya S, Hamel N, Tomiak E, et al. Germline and somatic SMARCA4 mutations characterize small cell carcinoma of the ovary, hypercalcemic type. *Nat Genet* 2014;46:438–43.
43. Earp MA, Raghavan R, Li Q, Dai J, Winham SJ, Cunningham JM, et al. Characterization of fusion genes in common and rare epithelial ovarian cancer histologic subtypes. *Oncotarget* 2017;8:46891–9.
44. Saito Y, Koya J, Araki M, Kogure Y, Shingaki S, Tabata M, et al. Landscape and function of multiple mutations within individual oncogenes. *Nature* 2020;582: 95–99.
45. Morimoto A, Serada S, Enomoto T, Kim A, Matsuzaki S, Takahashi T, et al. Annexin A4 induces platinum resistance in a chloride-and calcium-dependent manner. *Oncotarget* 2014;5:7776–87.
46. Saga Y, Ohwada M, Suzuki M, Konno R, Kigawa J, Ueno S, et al. Glutathione peroxidase 3 is a candidate mechanism of anticancer drug resistance of ovarian clear cell adenocarcinoma. *Oncol Rep* 2008;20:1299–303.
47. Tamaresis JS, Irwin JC, Goldfien GA, Rabban JT, Burney RO, Nezhat C, et al. Molecular classification of endometriosis and disease stage using high-dimensional genomic data. *Endocrinology* 2014;155:4986–99.
48. Worley MJ, Liu S, Hua Y, Kwok JS-L, Samuel A, Hou L, et al. Molecular changes in endometriosis-associated ovarian clear cell carcinoma. *Eur J Cancer* 2015;51: 1831–42.
49. Berek JS, Chalas E, Edelson M, Moore DH, Burke WM, Cliby WA, et al. Prophylactic and risk-reducing bilateral salpingo-oophorectomy: recommendations based on risk of ovarian cancer. *Obstet Gynecol* 2010;116:733–43.
50. Delair D, Han G, Irving JA, Leung S, Ewanowich CA, Longacre TA, et al. HNF-1 β in ovarian carcinomas with serous and clear cell change. *Int J Gynecol Pathol* 2013;32:541–6.
51. Tan TZ, Ye J, Yee CV, Lim D, Ngoi NYL, Tan DSP, et al. Analysis of gene expression signatures identifies prognostic and functionally distinct ovarian clear cell carcinoma subtypes. *EBioMedicine* 2019;50:203–10.
52. Vasan N, Razavi P, Johnson JL, Shao H, Shah H, Antoine A, et al. Double PIK3CA mutations in cis increase oncogenicity and sensitivity to PI3K α inhibitors. *Science* 2019;366:714–23.