

UC Irvine

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health

Title

(O-15) Factors Associated with Conversion to In-person Visit Among Patients Presenting for Pediatric Telehealth Encounters

Permalink

<https://escholarship.org/uc/item/3ng2w4nt>

Journal

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health, 25(1.1)

ISSN

1936-900X

Authors

Lim, C. Anthony
Tomlinson, Kirk
Stoffels, Guillaume
et al.

Publication Date

2024

DOI

10.5811/westjem.63022

Copyright Information

Copyright 2024 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

7 (O-I5) Factors Associated with Conversion to In-person Visit Among Patients Presenting for Pediatric Telehealth Encounters

Kirk Tomlinson, MD; Guillaume Stoffels, MA MS; Yvette C. Calderon, MD,MS

Oral Presenter: C. Anthony Lim, MD MS

Objectives: To identify patient- and illness-related factors associated with conversion to in-person visits among children presenting for pediatric telehealth encounters.

Background: Increases in telehealth utilization during the COVID-19 pandemic have been driven by limited availability of office visits due to infection prevention guidelines, parental hesitancy to enter healthcare facilities, and parity in reimbursement for providers. Due to increased demand and limited number of telehealth providers, it is important to determine which children may benefit the most from these encounters.

Methods: In this retrospective case-control study, patients 0-21 years old presenting for a telehealth encounter were evaluated. Children who had an in-person visit within seven days of the telehealth visit were identified as conversion cases and matched in a 1:3 ratio to controls by age in months up to 36 months and by year for those 4 and older. Patient demographics, past medical history, symptoms, and diagnoses were collected. A multivariable logistic regression model was developed including variables significantly associated with conversion on univariate analysis.

Results: From March–April 2020, there were 2,465 pediatric telehealth encounters. Of these, there were 67 (3%) conversions to in-person visits. 79% of these conversions originated from general pediatric telehealth encounters and the remaining from subspecialty telehealth visits. 69% of these conversions were to the ED and 31% in the clinic. Median days to in-person visit was 2 (1, 5). Median age was 25 months (1, 172), 66% were female, and 43% had a chronic medical condition. 55% were uninsured or on Medicaid, and the remaining were commercially insured. The most common symptoms reported included 42% respiratory, 22% fever, 19% pain, 18% vomiting/diarrhea, and 18% rash. After matching with controls based on age, a multivariable logistic regression model revealed that a history of cancer (OR 15.8, 95% CI 1.3-195.0), emesis (OR 4.6, 95% CI 1.1-18.9), pain (OR 4.5, 95% CI 1.5-13.3), non-COVID-19 related respiratory symptoms (OR 3.9, 95% CI 1.5-9.7), and telehealth visit with a specialist in allergy, endocrinology, gastroenterology, or pulmonary (OR 0.3, 95% CI 0.1-0.7) were associated with conversion, with an AUC of 0.82.

Conclusion: This introductory evaluation may suggest that certain patient- and illness-related factors are associated

with telehealth conversion to in-person encounters. To appropriately allocate telehealth and ambulatory resources, further study will determine whether some children will benefit if they are triaged directly to in-person visits.

8 (O-F6) A Mixed-methods Study of Barriers and Facilitators to Point-of-care Ultrasound Implementation for Emergency Department Providers at the Durham Veterans Affairs Healthcare System

Anna Tupetz, DPT, MScGH; Luna Ragsdale, MD, MPH; Padmaja Krishnan, MS-II; Raelynn Vigue, PGY-1; Carson E. Herman; Jaran White; Erica Peethumnongsin, MD, PhD; Catherine A. Staton, MD MSc; Alexander Gordee, MA; Maragatha Kuchibhatla, PhD; Stephanie Eucker, MD, PhD

Oral Presenter: Rebecca G. Theophanous, MD, MHSc

Objectives: Our primary objective was to identify the facilitators and barriers to optimize our program implementation using the Consolidated Framework for Implementation Research (CFIR) and test for impact at a single local emergency department (ED).

Background: Emergency ultrasound program leaders nationwide recognize that point-of-care ultrasound (POCUS) knowledge retention and utilization are difficult to achieve. Prior studies have identified a lack of provider training with a gap in POCUS knowledge and skills, lack of credentialed ultrasound users, and lack of quality assurance image review as significant barriers to POCUS use. A standardized approach to identifying and addressing barriers to sustainable POCUS implementation is needed to increase POCUS use, reduce radiology ultrasound, and potentially improve ED flow.

Methods: Our mixed-methods study implemented a co-designed, multifaceted intervention at the Durham Veterans Affairs ED from November 2021–October 2022 (12 months) to enhance POCUS usability and sustainability, including education, equipment knowledge, quality review process, and image archiving in the health record. Furthermore, 20/25 (80%) full-time ED providers participated in small- or large-group hands-on educational POCUS training sessions between February–May 2022. We conducted 14 semi-structured interviews to identify emergent themes and codes on ED POCUS use and performed team-based coding using inductive and then deductive analysis using NVivo. For our impact evaluation, we assessed POCUS program acceptability, effectiveness, and feasibility via provider pre/post-course questionnaires, interviews, and health record data (ED POCUS, radiology ultrasound orders, and ED length of stay (LOS)).

Results: Five POCUS themes emerged: convenience and