

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

EVIDENCE AGAINST PRODUCTION OF MAGNETIC MONOPOLES AT FERMILAB

### Permalink

<https://escholarship.org/uc/item/3x80s2w2>

### Authors

Eberhard, P.H.

Ross, R.R.

Taylor, J.D.

et al.

### Publication Date

1975-02-01

00004301797

Submitted to Physical Review D

RECEIVED  
LAWRENCE  
RADIATION LABORATORY

LBL-3680  
Preprint c.1

MAR 25 1975

LIBRARY AND  
DOCUMENTS SECTION

EVIDENCE AGAINST PRODUCTION OF  
MAGNETIC MONOPOLES AT FERMILAB


P. H. Eberhard, R. R. Ross, J. D. Taylor,  
L. W. Alvarez and H. Oberlack

February 1975

Prepared for the U. S. Atomic Energy Commission  
under Contract W-7405-ENG-48

**For Reference**

Not to be taken from this room



LBL-3680  
c.1

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

## EVIDENCE AGAINST PRODUCTION OF MAGNETIC MONOPOLES AT FERMILAB\*

P.H. Eberhard, R.R. Ross, J.D. Taylor, and L.W. Alvarez

Lawrence Berkeley Laboratory, University of California  
Berkeley, California 94720

and

H. Oberlack

Max-Planck-Institut für Physik and Astrophysik  
Munich, Germany

## ABSTRACT

A search for magnetic monopoles that requires very few assumptions about their properties has been performed in material exposed to protons accelerated at Fermilab. No monopoles were found. If monopoles exist with masses less than 12 GeV, the probability of pair production in a proton-nucleon collision is of the order of  $10^{-18}$  or less with 95% confidence.

## INTRODUCTION

Since 1931,<sup>1</sup> the existence of magnetic monopoles has been repeatedly invoked in theories<sup>2</sup> in connection with the observed phenomenon of electric charge quantization.<sup>3</sup> However, increasingly exhaustive experiments designed to find and isolate magnetic charges have all had negative results.<sup>4-10</sup> These experiments would have detected monopoles with various magnetic charges, masses, production cross sections and specific binding properties to matter.<sup>11</sup> In this paper, a very general search for monopoles that can be produced at present accelerators is reported. It uses a modified version of the detector<sup>12,13</sup> used in a previous cosmic ray search.<sup>6,8</sup> It covers a vast domain of charges and masses, requires no extraction of monopoles from material, and, to be valid, needs very few assumptions about the properties of monopoles.

## IRRADIATION AND SAMPLE PREPARATION

In an experiment at the Fermi National Accelerator Laboratory, aluminum targets were irradiated by about  $4 \times 10^{18}$  protons accelerated to several hundreds of GeV in the hope of producing and trapping monopole pairs. The target specifications are given in Table I. From the accelerator records and the geometry of the targets, the number of proton interactions is computed at each energy. Within 30%, it agrees with the estimation based on the  $^{22}\text{Na}$  and  $^7\text{Be}$  radioactivity in the first centimeter of targets 1, 2, and 3. The total for all targets is shown in Table II.

In order to search for monopoles that could have been produced in pairs in some of these interactions and trapped in the targets, the target material was first ground into thin chips to separate the north and south poles of a pair, using a milling machine advancing 10  $\mu\text{m}$  between successive cuts.<sup>14</sup> Then, the chips were placed in a hollow rotating sphere to be randomized. They were divided into 30 samples and the magnetic charge of each sample was measured in an electromagnetic detector.<sup>12,13</sup>

#### THE MEASUREMENT OF THE MAGNETIC CHARGE

The detector is shown schematically in Fig. 1. The sample is carried several times around a path that traverses a coil (sensing coil). This coil is part of a superconducting circuit containing two other coils (field coils) each one wound around a sensitive magnetometer (SQUID).<sup>15</sup> If a sample has a non-zero magnetic charge, it will induce a change of current in the superconducting circuit and a change  $\Delta\phi_1$  and  $\Delta\phi_2$  in the flux measured by SQUIDS 1 and 2. For each SQUID:

$$\frac{\Delta\phi}{\phi_0} = \frac{v_s N_p}{f} \quad (1)$$

where  $v_s$  is the ratio of the sample magnetic charge  $g_s$  to the Dirac unit

$$g_0 = \frac{e}{2\alpha} = \frac{137}{2} e \text{ (in Gaussian units)} \quad (2)$$

$\phi_0$  is the flux quantum of superconductivity ( $2.07 \times 10^{-7}$  Gauss  $\text{cm}^2$ ),  $N_p$  the number of passes through the sensing coil and  $f$  a constant depending on the various inductances of the circuit.<sup>13</sup> For SQUID 1 (SQUID 2),

$$f = 34(290).$$

The magnetic charge measurement was performed by taking magnetometer readings after 1, 3, 9, 27, and 81 passes. This procedure provided an accuracy of 0.03 on the value of  $v_s$  with the restriction described in Appendix 1 and in Ref. 13.

A magnetic charge is found zero if successive measurements of the current stored in the circuit are found to be the same within errors. Equipment instabilities would result in different current readings, i.e., in spurious non-zero magnetic charge measurement. In our procedure they would trigger a thorough check of the equipment and a remeasurement of the sample. Therefore, this method of search for monopoles is quite safe against equipment failures.

Furthermore, in the Maxwell equation,

$$\text{curl } \vec{E} = -\frac{1}{c} \frac{\partial \vec{B}}{\partial t} - \frac{4\pi}{c} \vec{J}_m \quad (4)$$

the magnetic current  $\vec{J}_m$  has the same effect as a time derivative of the magnetic induction  $\vec{B}$ . Therefore, we were able to test the adequacy of our apparatus to detect magnetic charge using induction in the sensing coil, exactly as if magnetic charges were available for that test.

## RESULTS

The magnetic charges  $v_s$  of all the samples were measured consistent with zero and incompatible with any value larger than 0.1 (except for the restrictions of Appendix 1). From this result, a maximum value  $R_{\max}$  for the ratio of the number of monopole pairs to the number

of interactions has been computed at a 95% confidence level.  $R_{\max}$  is shown in Table II for the three different incident proton energies. These figures are valid when the monopole charge  $\nu$  (in units of  $g_0$ ) is included between 1 and 7, where the only sizeable correction comes from the probability that chips of opposite charge end up in the same sample in spite of the randomization. For  $1 \leq \nu \leq 7$ , the expected energy loss of the monopole in aluminum<sup>16</sup>

$$\frac{dE}{dx} = \nu^2 \cdot 21 \text{ GeV/cm} \quad (5)$$

ensures that most monopoles stop in the targets and multiple scattering is large enough to separate the north and south poles of a pair by more than the chip size.

The upper limits  $\sigma_{\max}$  for the cross sections for monopole pair production in proton nucleon collision have been computed at a 95% confidence level and plotted on Fig. 2. The interactions in aluminum were assumed to correspond to a total proton nucleon cross section of 35 mb. Figure 2 shows  $\sigma_{\max}$  for values of  $\nu$  between 0.01 and 100, i.e., even for fractional values forbidden by the Dirac theory but for which our search is still meaningful. For  $\nu > 7$  or  $\nu < 1$ , various corrections are needed as described in Appendix 2. In order to determine upper limits independent of the production process, the most unfavorable case for the monopole detection was considered where monopoles of a pair are produced with  $0^\circ$  opening angle and with the same energy. The upper limits indicated by the solid curve on Fig. 2 correspond to additional plausible assumptions concerning monopole properties<sup>17</sup>; the dashed curve corresponds to assumptions more favorable and the dotted curve extremely



unfavorable for detecting monopoles. Assumptions for all three are described in Appendix 2.

In any case, monopoles with masses larger than 12 GeV (that could not be produced in our proton nucleon interactions), tachyons<sup>18</sup> and zero mass<sup>19</sup> monopoles (that would not stop in the material) would escape this search.

#### OTHER MATERIAL SEARCHED

The detector was also used to search for monopoles in a steel cylinder exposed to 18 GeV electrons at SLAC and in a  $100 \text{ cm}^2 \times 2 \text{ mm}$  thick stainless steel piece of the ISR vacuum chamber that was located near an interaction point. No magnetic charge was found. The number of interactions seen by this material (including a correction for the solid angle in the ISR case) is shown in Table II. The value of  $R_{\text{max}}$ , the maximum ratio of the number of monopole pairs produced to this number of interactions, which is compatible with our results, is also shown.<sup>20</sup>

The SLAC target would have been suited to detect monopoles if their mass was lower than 2.5 GeV but their production cross section by electrons higher than by protons. The ISR material would be more suited if monopoles had a large mass, charge, and opening angle in the center-of-mass system. Unless one of these circumstances is true, the chance to detect monopoles here would be smaller than in cosmic ray experiments.<sup>8</sup>

## CONCLUSIONS

No monopoles were detected in material exposed to  $4 \times 10^{18}$  protons. The ratio of monopole pairs produced to the number of interactions is of the order of  $10^{-18}$  or less for a large range of charge and for different assumptions about monopole properties. If monopoles have masses less than 12 GeV and are subject to strong interactions, pair production is affected by a very strong suppressing mechanism.<sup>21</sup> Indeed, hadrons, produced by strong interactions, are produced with cross sections that range from  $10^{-27}$  to  $10^{-30}$  cm<sup>2</sup> while weak interaction cross sections are of the order of  $10^{-36}$  cm<sup>2</sup> in this energy region. Our limits for monopole production cross sections by proton-nucleon interactions (Figs. 2a and b) are far below the figures for weak interactions. If monopoles exist, their masses are probably higher than 12 GeV.

Similar conclusions could be drawn from previous NAL experiments<sup>9,10</sup> and their upper limits are only one or two orders of magnitude above ours. However, the validity of those earlier experiments is dependent on the assumption that a monopole can be extracted from material with or without a nucleus attached to it. Indeed, the extraction procedure cannot be tested without monopoles, while Eq. (4) shows that the effect of magnetic current used in our detector can be simulated exactly by magnetic induction. Some cosmic ray experiments<sup>4,5</sup> also give cross section limits only a few orders of magnitude above ours, but for them, assumptions about migration in the atmosphere and in the ocean water are needed in addition to the possibility of extraction. With respect to the cosmic ray experiment using the lunar material,<sup>6,8</sup> the present experiment

has the advantage of relying on more controlled conditions for the production and trapping of monopole pairs and it results in limits about 3 orders of magnitude lower.

#### ACKNOWLEDGMENTS

We are indebted to G. Gibson for his help with the measurements, to Drs. D. Eartley, R. Stefanski, D. Theriot, and P. Neeson at Fermilab, to Drs. H. Blewett at the ISR and J. Murray at SLAC for their help in getting the irradiation, to L. Cadra, P. Bringham, A. Smith and the staff of Safety Services at LBL for handling and preparing the radioactive material, and to W. Kelly, M. Attencio and A. Roberts for help in constructing the targets. The experiment was possible only because the monopole detector was loaned to us by NASA.

Footnote and References

\* Work performed under the auspices of the U.S. Energy Research and Development Administration.

1. P.A.M. Dirac, Proc. Roy. Soc. (London) A133, 60 (1931).
2. Some recent examples are: J. Schwinger, Phys. Rev. 173, 1536 (1968); Science 165, 757 (1969); G. 't Hooft, Nuclear Physics 379, 276 (1974); N.H. Christ, Phys. Rev. Lett. 34, 355 (1975).
3. J.G. King, Phys. Rev. Letters 5, 562 (1960); L.J. Fraser, E.R. Carlson, and V.W. Hughes, Bull. Am. Phys. Soc. 13, 636 (1968).
4. R. L. Fleischer, H.R. Hart, Jr., I.S. Jacobs, P.B. Price, W.M. Schwarz, and R.T. Woods, J. Appl. Phys. 41, 958 (1970).
5. H.H. Kolm, F. Villa, and A. Odian, Phys. Rev. D 4, 1285 (1971).
6. P.H. Eberhard, R.R. Ross, L.W. Alvarez, and R.D. Watt, Phys. Rev. D 4, 3260 (1971).
7. I.I. Gurevich, S. Kh. Khakimov, V.P. Martemianov, A.P. Mishakova, L.A. Makar'ina, V.V. Orgurtzov, V.G. Tarasenkov, L.A. Chernishova, L.M. Barkov, M.S. Zolotorev, V.S. Ohapkin, and N.M. Tarakanov, Phys. Lett. 38B, 549 (1972).
8. R.R. Ross, P.H. Eberhard, L.W. Alvarez, and R.D. Watt, Phys. Rev. D 8, 698, (1973).
9. R.A. Carrigan, Jr., F.A. Nezrick, and B. P. Strauss, Phys. Rev. D 8, 3717 (1973).
10. R.A. Carrigan, Jr., F.A. Nezrick, and B.P. Strauss, Phys. Rev. D 10, 3867 (1974).

11. For a summary of the properties assumed in experiments performed up to 1971, see Table III of Ref. 6.
12. L. Alvarez, M. Antuna, Jr., R. Byrns, P. Eberhard, R. Gilmer, E. Hoyer, R. Ross, H. Stellrecht, J. Taylor, and R. Watt, Rev. Sci. Instr. 42, 326 (1971).
13. P.H. Eberhard, R.R. Ross, and J.D. Taylor, Lawrence Berkeley Laboratory Report No. LBL-1732 unpublished and TBP Rev. Sci. Instr.
14. The cut was 100  $\mu\text{m}$  deep and 12 mm wide.
15. R.C. Jaklevic, J. Lambe, J.E. Mercereau, and A.H. Silver, Phys. Rev. 140A, 1928 (1965); J.W. Zimmerman and A.H. Silver. Phys. Rev. 141, 367 (1966); R.L. Forgacs and A. Warnick, IEEE International Convention Record 14, 90 (1966).
16. E. Bauer, Prod. Cambridge Nul. Sec. 47, 777 (1951); H.J.D. Cole, *ibid* 47, 196 (1951).
17. For instance, binding to aluminum nuclei; D. Sivers, Phys. Rev. D 2, 2048 (1970).
18. L. Parker, Phys. Rev. 188, 2287 (1969); D. Weingarten, Ann. of Phys. 76, 510 (1973); R. Mignani and E. Recami, Lett. Nuovo Cimento Ser. 2 9, 367 (1974).
19. A. Salam, Phys. Letters 22, 683 (1966).
20. As for the Fermilab targets,  $R_{\text{max}}$  is corrected for the probability that the north pole and the south pole of a pair would end up in the same sample. The SLAC target was ground in 10  $\mu\text{m}$  chips, randomized and run in 16 different samples. The ISR target was cut in 16 equal size pieces and run in two samples, each one composed of non-adjacent pieces.

21. The possibility of a very strong suppressing mechanism has been suggested by M.A. Ruderman and D. Zwanziger, Phys. Rev. Letters 22, 146 (1969). However, a reliable computation of the corresponding suppressing factor has still to be made.
22. W.C. Carithers, R. Stefanski, R.K. Adair, Phys. Rev. 149, 1070 (1966).
23. The strength of the bond was estimated by considering the magnetically charged aluminum nucleus to be a sphere of  $1.3 \text{ \AA}$  radius that was to be driven through the aluminum of our targets, which has a Brinell hardness no. of  $102 \text{ kG/mm}^2$ .
24. E. Goto, J. Phys. Soc. Japan 13, 1413 (1958).

## APPENDIX 1

Restrictions on the validity of our charge measurements for a few theoretically unexpected values of  $v_s$  result from the periodic response of SQUIDS to magnetic flux changes.<sup>13</sup> After passing the sample  $N_p$  times, each SQUID provides a measurement of  $\Delta\phi$ , modulo  $\phi_0$ , therefore of  $v_s$  modulo  $f/N_p$ . Therefore, a measurement may fail to detect charges such that  $v_s N_p/f_1$  and  $v_s N_p/f_2$  are both equal to an integer within error. To avoid most of these failures, the search in the samples was performed with  $N_p = 81$  with intermediate stops and magnetometer readings after  $N_p = 1, 3, 9$  and  $27$ . Any magnetic charge would have been detected except for the small ones ( $v_s < 0.1$ ) and some very special large ones, (all  $\geq 580$ ) equal within error, to a multiple of  $f_1$  and  $f_2$  at the same time.

## APPENDIX 2

For  $v < 1$  and  $v > 7$ , the values of  $R_{\max}$  given in Table II for the Fermilab targets need further corrections. Those corrections are highly charge dependent and are deduced on the basis of assumptions described below.

For values of  $v$  less than 1 (incompatible with the Dirac theory) the ionization is not sufficient to stop all the monopoles produced. A correction (referred to as correction No. 1 later on) is computed assuming all the monopoles are produced with a typical velocity equal to the velocity of the proton-nucleon center-of-mass system and that they lose 1/2 of their energy every time they collide with an aluminum nucleus, as do protons at high energy when they collide with nuclei.<sup>22</sup>

North and south poles of a pair with large magnetic charges may stop close enough so that the attractive force between them drives them together toward annihilation.<sup>23</sup> Separation due to multiple Coulomb scattering is sufficient to avoid this effect for  $\nu < 20$ . For  $20 < \nu < 60$ , large angle Coulomb scattering, and for  $\nu > 60$  nuclear scattering with half energy loss, are used to estimate a correction (No. 2). In all cases, we considered that monopoles of a pair are produced at the same energy and with  $0^\circ$  opening angle.

Another correction (No.3) takes into account the probability that the two monopoles of a pair end up in the same chip. It is estimated on the basis of multiple Coulomb scattering and affects the upper limits for  $\nu > 7$ .

If monopoles had charges  $\nu < 0.1$ , but there were many of them, the statistical fluctuations would generate some measurable charges for the samples. Therefore, our experiment allows computation of an upper limit for the density of such monopoles, with a reduced sensitivity (correction No. 4.)

These assumptions, which are quite pessimistic about the sensitivity of our experiment, correspond to the solid curve of Fig. 2. The dashed curve corresponds, for  $\nu < 0.5$ , to monopoles coming out of the aluminum nucleus in which they are produced with a very small energy (i.e., without correction No. 1), and, for  $\nu > 20$ , corresponds to magnetically charged aluminum nuclei very strongly bound to the crystal lattice (i.e., without correction No. 2). Therefore, the effect of correction No. 1 (No. 2) is shown by the difference between the solid and the dashed



curve of Fig. 2 for  $\nu < 1$  ( $\nu > 7$ ). The effect of corrections Nos. 3 and 4 are shown by the difference between the dashed curve value and the solid curve value for  $\nu = 1$ .

In order to cover an even more pessimistic but unlikely case for the sensitivity of our experiment, wherein the magnetically charged aluminum nuclei do not bind to the crystal but would be free to move inside the material, a magnetized iron case producing a field of about 1.5 Gauss was built around target No. 5. For a sufficient initial separation, the poles of a pair would drift in opposite directions along field lines to the iron, where they would be trapped.<sup>24</sup> The iron case was divided into separate samples and processed in the detector. The dotted line of Fig. 2 corresponds to that case and to the other pessimistic assumption where monopoles have no nuclear interaction (like muons), and do not stop in the target for  $\nu < 0.5$ .

Table I. Characteristics of the aluminum targets exposed to the Fermilab beams.

Target No.	Length (cm)	Location of irradiation <sup>a</sup>	No. of protons in the beam	No. of interaction lengths before the target
1	30	$\nu$	$1.3 \times 10^{18}$	0
2	30	$\nu$	$1.3 \times 10^{18}$	0
3	45	$\nu$	$1.0 \times 10^{18}$	0
4	16.5	$\nu$	$0.2 \times 10^{18}$	0.3
5	41	P	$0.4 \times 10^{18}$	0.5

<sup>a</sup> $\nu$  stands for neutrino lab.

P stands for proton lab, eastern section.

Table II. Energy distribution of the exposures of the material analyzed.  $R_{\max}$  is the maximum ratio of monopole pairs to primary interactions (95% confidence).

Origin	Beam energy (GeV)	Material	No. of primary interactions	$R_{\max}$	Maximum monopole mass (GeV)	Range of $\nu^a$
FERMILAB	200	Aluminum	$2.0 \times 10^{17}$	$1.6 \times 10^{-17}$	8.8	1 to 7
FERMILAB	300	Aluminum	$2.5 \times 10^{18}$	$1.3 \times 10^{-18}$	10.9	-
FERMILAB	400	Aluminum	$6.6 \times 10^{16}$	$5.0 \times 10^{-17}$	12.8	-
SLAC	18	Iron	$1.4 \times 10^{19}$	$2.3 \times 10^{-19}$	2.5	1 to 3
ISR	11.5	Stainless steel	$3.5 \times 10^9$	$1.1 \times 10^{-9}$	10.5	$\nu \geq 1$
ISR	15	Stainless steel	$4.9 \times 10^9$	$7.8 \times 10^{-10}$	14	-
ISR	22.5	Stainless steel	$4.2 \times 10^{10}$	$9.0 \times 10^{-11}$	21.5	$\nu \geq 2$
ISR	26.5	Stainless steel	$4.6 \times 10^{10}$	$8.3 \times 10^{-11}$	25.5	-
ISR	31.4	Stainless steel	$4.6 \times 10^8$	$8.3 \times 10^{-9}$	30.4	-

<sup>a</sup>Range of  $\nu$  for which  $R_{\max}$  in the table does not need any additional correction.

Figure Captions


FIG. 1. Schematic view of the detector. The sample is moved along the dashed curve labeled sample path. The superconducting circuit is shown with the sensing coil and a field coil connected in series. The magnetometer and an auxiliary coil are also shown inside the cryostat.

FIG. 2. Upper limit (95% Confidence level) on monopole pair-production cross section in proton-nucleon collisions as a function of magnetic charge.

a) 300 GeV/c protons on aluminum


b) 400 GeV/c protons on aluminum.

The solid curve corresponds to corrections 1 to 4 described in Appendix 2, the dashed curve to corrections 1 and 2 equal to 1. The dotted curve has been computed, using target 5 only, with a most pessimistic view of the sensitivity of the experiment.



XBL7410-4456

Fig. 1.



XBL 752 - 2396

Fig. 2a.




Fig. 2b.

**LEGAL NOTICE**

*This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.*



TECHNICAL INFORMATION DIVISION  
LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720