

**UCLA**

**UCLA Previously Published Works**

**Title**

Family Presence for Critically Ill Patients During a Pandemic

**Permalink**

<https://escholarship.org/uc/item/4t20309v>

**Journal**

CHEST Journal, 160(2)

**ISSN**

0012-3692

**Authors**

Hart, Joanna L  
Taylor, Stephanie Parks

**Publication Date**

2021-08-01

**DOI**

10.1016/j.chest.2021.05.003

Peer reviewed

# Family Presence for Critically Ill Patients During a Pandemic



Joanna L. Hart, MD; and Stephanie Parks Taylor, MD

Family engagement is a key component of high-quality critical care, with known benefits for patients, care teams, and family members themselves. The COVID-19 pandemic led to rapid enactment of prohibitions or restrictions on visitation that now persist, particularly for patients with COVID-19. Reevaluation of these policies in response to advances in knowledge and resources since the early pandemic is critical because COVID-19 will continue to be a public health threat for months to years, and future pandemics are likely. This article reviews rationales and evidence for restricting or permitting family members' physical presence and provides broad guidance for health care systems to develop and implement policies that maximize benefit and minimize risk of family visitation during COVID-19 and future similar public health crises.

CHEST 2021; 160(2):549-557

**KEY WORDS:** COVID-19; critical care; ethics; family; policy

Critical illness has lasting effects on patients and their families long after the illness episode.<sup>1-3</sup> Family members of critically ill patients themselves frequently experience symptoms of depression and traumatic stress.<sup>1</sup> Family-centered care improves quality of critical care and the experience for patients and families.<sup>2,4</sup> Over the past two decades, ICUs have made significant progress toward family-centered care, including building the evidence base and implementing effective approaches.<sup>2</sup> Central to this progress has been the designing of structures in which family members are an integrated part of the critical care team rather than visitors; such designs include open visitation, inclusion in interdisciplinary team rounds, participation in direct

caregiving, frequent communication, shared decision-making, and opportunities to be present during resuscitative efforts.<sup>2</sup> In-person family presence is a key facilitator of these family-centered care practices.

However, the emergence of the COVID-19 pandemic in 2020 resulted in deimplementation of family-centered care because of prohibitions or restrictions on the presence of patient family members.<sup>4-7</sup> Avoiding gathering and maintaining social distancing have been central strategies in reducing the transmission of SARS-CoV-2, the virus causing COVID-19. The application of these principles in hospitals led to restrictions on the presence of family members.<sup>8,9</sup> Although such restrictions were

**ABBREVIATIONS:** PPE = personal protective equipment

**AFFILIATIONS:** From the Division of Pulmonary, Allergy, and Critical Care, Department of Medicine (J. L. Hart), the Department of Medical Ethics and Health Policy (J. L. Hart), and the Palliative and Advanced Illness Research Center (J. L. Hart), University of Pennsylvania Perelman School of Medicine, Philadelphia, PA; and the Department of Internal Medicine (S. P. Taylor), Atrium Health's Carolinas Medical Center, Charlotte, NC.

**FUNDING/SUPPORT:** NIH K23HL132065 provided salary support for J. L. H.

**CORRESPONDENCE TO:** Joanna Hart, MD; email: [joanna.hart@penn-medicine.upenn.edu](mailto:joanna.hart@penn-medicine.upenn.edu)

Copyright © 2021 American College of Chest Physicians. Published by Elsevier Inc. All rights reserved.

**DOI:** <https://doi.org/10.1016/j.chest.2021.05.003>

reasonable in the early periods of limited scientific knowledge about viral transmission, supply shortages, and rapid efforts to restructure hospitals to expand capacity,<sup>10</sup> reevaluation of these policies is now critical given our increased information and resources.<sup>11-13</sup> ICU visitation restrictions such as those enacted because of COVID-19 lead to incomplete grief (eg, inability to fully experience the grieving process)<sup>11,14-18</sup>; emotional distress experienced by patients, families, and physicians<sup>11,14-23</sup>; barriers to high-quality communication and decision-making<sup>11,14-22,24</sup>; perpetuation of existing inequities<sup>11,20</sup>; and poor clinical outcomes.<sup>11,14,17-20,24-26</sup> Given the serious and long-lasting harm these restrictions cause for patients, family members, staff, and communities, we believe broadly prohibiting visitation of family members of patients with critical illness, including those with COVID-19, is no longer justified.

Although improved COVID vaccine availability and decreased COVID-related hospital strain have led hospitals to relax some restrictive visitation policies, prohibitions and restrictions on family visits to patients with and without COVID-19 persist.<sup>27,28</sup> Because COVID-19 will continue to be a public health threat for months to years and similar future pandemics are likely,<sup>29</sup> hospitals urgently need to develop visitation policies that maximize family-centered care while mitigating the risks of in-person family presence to patients, family members, and hospital staff. This article provides broad guidance for hospitals to develop and implement policies that maximize benefit and minimize risk of family visitation during COVID-19 and future similar public health crises.

## Common Rationales for Limiting Family Presence

During the pandemic, hospitals broadly prohibited visitation because of concerns that family members of admitted patients may transmit virus through person-to-person contact within the hospital.<sup>4,30,31</sup> Persistence of visitation restrictions for patients with COVID-19 reflects the desire to protect family members from contracting and transmitting COVID-19 after exposure to the infected patient.<sup>32</sup> However, there is no existing evidence that family members of admitted patients are a significant contributor to nosocomial respiratory viral spread, or that prohibiting visitation to patients with COVID-19 protects against viral transmission to patients, staff, or visitors.<sup>33,34</sup> Existing Centers for Disease Control and Prevention guidelines

detail infection control processes that effectively prevent viral transmission in health care settings, particularly the proper use of personal protective equipment (PPE).<sup>32</sup> The increased availability of effective COVID vaccines, including for health care workers and individuals at high personal risk of severe COVID-19, have further reduced the direct risks to staff and patients.<sup>35</sup> Providing family members with the opportunity to discuss and judge the benefits and potential harms of visitation, using these risk reduction methods (eg, PPE use, prior vaccination, and distancing when out of the patient's room), promotes their individual autonomy.<sup>34</sup> Protecting families' and patients' self-determination by allowing them to decide whether in-person presence is preferred, based on their personal risk of severe COVID-19 and individual values, may mitigate against the loss of control they commonly experience during an ICU stay.<sup>36</sup> Health care systems and physicians may view family presence as a potential threat to resources, specifically PPE and hospital staff time. Initially, the decision to prohibit family presence reflected the primacy of dedicating space, equipment, and personnel resources to provide life-saving care to the surge of patients with COVID-19.<sup>10,30</sup> However, advancement in our knowledge and stabilization of resources since those early months have led to an enhanced understanding of the anticipated effect of family visitation on health care system resources.

PPE was in short supply early in the pandemic, and health care systems experienced challenges in acquiring and maintaining sufficient PPE to protect health care workers.<sup>37,38</sup> PPE is now more readily available and proven effective in reducing viral transmission.<sup>32</sup> Health care systems should incorporate the PPE needs of family members into supply chain calculations without concern for compromising the availability of adequate PPE for health care workers.

Hospital staff are a valuable health care resource, particularly during the COVID-19 crisis. Implementing processes to allow families to safely visit patients with COVID-19 will likely require additional effort by health-care personnel, specific to the need of individual sites. The additional needs might include dedicated personnel for screening and logging family members entering the hospital and assisting them with safely donning and doffing PPE. During times of extreme resource strain when no dedicated personnel are available, bedside physicians can facilitate PPE use to family members, although this should be a short-term solution if possible.

Overall, the benefits of permissive family visitation are expected to more than offset the additional personnel effort required to facilitate safe visitation.<sup>13</sup> Physicians are now more comfortable in caring for patients with COVID-19, and the need to simplify bedside workflow to enable adaptation to a dramatically altered environment is no longer pressing. In the typical ICU environment, there is no evidence that family presence contributes to strain.<sup>39,40</sup>

### Benefits of Unrestricted Family Presence During COVID-19

First, the physical presence of family members enhances integration of family members into the ICU workflow. Communication strategies, such as videoconferencing processes and routinized outreach to families, have partially filled gaps left by visitation prohibitions.<sup>4,19,22</sup> Yet, these strategies also create demands on health care workers to facilitate patient-family connections, explain technological tools, and establish longitudinal relationships with distant family members.<sup>41</sup> Family-centered care guidelines support family presence during ICU rounds, and the vast majority of families prefer to be present.<sup>2,42</sup> In observational work, family presence was associated with higher satisfaction with care, collaboration with the medical team, shared decision-making, and facilitated information gathering for family members, with minimal additional rounding time.<sup>43,44</sup> COVID-era visitation restrictions lengthened ICU stays and delayed decisions to limit treatments before death.<sup>24</sup> Therefore, supporting family presence for patients with COVID-19 is expected to improve the efficiency and quality of relationship-building between families and the medical team and preserve ICUs and their staff as critical resources.

Second, family presence may reduce delirium, which is common among patients with COVID-19.<sup>25,26,45,46</sup> Delirium is associated with longer ICU and hospital lengths of stay, reintubations, ICU readmissions, and higher workloads for nursing staff.<sup>46,47</sup> An observational study of more than 4,500 critically ill patients with COVID-19 from 69 ICUs revealed that family visitation was associated with a lower risk of delirium among critically ill patients with COVID-19, one of only two modifiable risk factors the study identified.<sup>26</sup> Therefore, supporting the presence of family members will lead to multiple improvements in patient- and ICU-level outcomes related to reduction in delirium.

Finally, family presence is critical to optimizing end-of-life and grief experiences for patients, families, and physicians. For families, deaths in the ICU are associated with complicated grief, posttraumatic stress disorder, and prolonged grief.<sup>17</sup> This is undoubtedly exacerbated in the setting of family separation around the time of death, due to incomplete grief resulting from health care system limits on the opportunities of families to carry out mourning rituals, or doing so in an abbreviated or delayed manner.<sup>4,15,17</sup> Prohibiting family presence is likely to further strain the ICU workforce in the short and long term.<sup>27</sup> Physicians may experience secondary trauma and moral injury due to participating in separation of families during critical illness and death, which has been widespread during the COVID-19 pandemic.<sup>5,27,48-51</sup> Added to the already high baseline levels of distress among critical care physicians, this may contribute to long-term attrition of the ICU workforce due to burnout, depression, anxiety, and posttraumatic distress.<sup>52,53</sup> Supporting family cohesion during critical illness may, in turn, protect ICU physicians from psychological distress associated with their role in the care of patients with COVID-19.

### Policy Recommendations

To guide health care systems in creating safe, equitable, and family-centered visitation policies, we provide a blueprint for supporting family presence for patients with COVID-19 in adult ICUs that can be adapted to the individual health care system's culture and structure (Table 1). Although these recommendations are directly relevant to pediatric and non-ICU settings as well, there may be additional considerations and logistics relevant for families of hospitalized children to respond to the children's developmental needs.

Development of equitable and acceptable visitation policies requires a diverse, interdisciplinary approach inclusive of key stakeholders,<sup>54,55</sup> such as patient and family advocates or representatives, community members, physicians from representative service lines, ethics staff, safety and quality experts, infection control experts, and security staff. Health care systems should leverage existing or novel community partnerships and patient advisory committees to participate in policy creation. After drafting and implementing an initial proposal, revisiting the procedures and policies can mitigate concerns through ongoing evolution and adaptation to meet or balance the needs of all stakeholders.

**TABLE 1 ] Recommendations for Visitation Policy Design and Implementation**

Development of policy
Convene key stakeholders to draft policy
Avoid policies that are highly dependent on levels of community viral transmission
Coordinate approach across units, hospitals, and health care systems
Establish and communicate plans to rapidly reassess and adapt policy
General visitation rules
Permit at least two family members per patient, including those with COVID-19
Avoid exceptions based on clinical condition or prognostication (eg, end-of-life periods)
Avoid establishing visiting hours but, if enacted, maximize hours to allow for overlap with rounds and access to family members with caregiving or work responsibilities
Entry and screening procedures
Encourage family members to make individualized decisions to visit based on personal risk of severe COVID-19, vaccination status, and values
Do not restrict visitors on the basis of relationship to the patient to promote opportunities for low-risk family members to visit rather than individuals at higher personal risk of COVID-19
Identify and mark entrances for visitors to undergo standardized screening for COVID-19
Establish a centralized visitor logging and pass process coordinated with the bedside teams
Educate visitors on safety protocols, such as universal masking and distancing, and use the built environment to encourage adherence
Personal protective equipment (PPE) use
Establish rapid N95 fit assessment protocols for visitors with local infection control
Mandate that visitors adhere to federal and local guidance for PPE use
Identify and train staff to observe visitors doffing and donning PPE
Nonphysical family presence
Support robust alternatives to in-person visitation, such as routinized communication standards and videoconferencing
Provide technology support to patients and families, particularly when restricting visitation
Communication of the policy to patients, families, and staff
Provide a clear rationale for the established policies, including the decision-making process leading to the policy
Written communication should be at a fifth to eighth grade reading level to promote public understanding
Provide information in Spanish, as well as other locally prevalent non-English languages
Navigating exceptions
Centralize visitation adjudication when possible to preserve the role of the clinical team and promote consistency and equity
Use a standardized approach to evaluate and grant exceptions that promote family-centered care and reduce moral distress while avoiding bias and harm associated with viral transmission
Other policy considerations
Visitors should not eat in the rooms of patients with COVID-19, due to inability to maintain PPE; other accommodations should be arranged
Health care systems choosing to restrict visitation to patients with COVID-19 should establish clear procedures for removing patients from isolation once no longer infectious
Clergy and spiritual leaders should not be considered visitors; visitation policies do not apply to these individuals
Hospitalized patients with disabilities may require reasonable accommodations and protections, including the presence of a caregiver. Caregivers of such patients should not be considered visitors; visitation policies do not apply to these individuals

Policies that are stable over time are likely to cause less confusion and eliminate additional strain on staff members who must learn and implement visitation procedures. Models for stable policies include those that do not need to be altered during viral surges in the

community (ie, that permit visitation throughout periods of strain and to patients with and without COVID-19) or those that include planned adaptations to changes in viral spread or health care system strain (ie, with clearly communicated thresholds for procedural changes).

Policies affecting family members of admitted patients should promote a standardized experience throughout the hospitalization, including transitions between ICUs and ward locations. When possible, this should include all health care systems in a referral area, similar to processes proposed for regional allocation of scarce resources.<sup>56</sup> Regional coordination and collaboration across health care systems, or across entities within a single system, can improve continuity for patients and family members.

### *General Visitation Rules*

A standard policy that permits at least two visitors for each patient with COVID-19 allows family members to provide mutual support and reduces the demands on a single individual. Flexibility and clearly delineated common exceptions should be included in the visitation policy to promote the respect and well-being of patients, family members, and staff. Adhering too strictly or too loosely to the standard policy is likely to result in distress among these groups given the complexity of family relationships and situations.

ICUs have been challenged by what has become a common exception allowing for more permissive visitation for patients at the end of life.<sup>57</sup> Although this is intended to promote family cohesion and grief during the dying process, the “end-of-life period” for a critically ill patient is difficult to define.<sup>58</sup> Indeed, family members of critically ill patients with COVID-19 experience preloss grief.<sup>59</sup> Because ICUs serve patients with life-threatening conditions requiring advanced monitoring and life support, rapid clinical decline is common. Even among ICU physicians, there is little agreement on when a patient is “acutely dying.”<sup>58</sup> Eliminating the end-of-life exception protects ICU staff from the stress of prognostication errors or disagreements. Promoting family presence throughout the critical illness also prevents unintended coercion leading patients or families to choose treatment limitations for the primary purpose of reuniting and avoids missed opportunities for meaningful final connections between patients and family members that are important for the grieving process.<sup>17,18</sup> Further, linking policy exceptions to discretionary judgments such as prognostication may predispose to inequitable application due to implicit or explicit biases.<sup>60</sup> We recommend making the standard policy for family visitation, particularly in the ICU, as permissible as is feasible rather than relying on exceptions based on patient clinical conditions.

There is no reason to limit family presence on the basis of time of day in most circumstances. Providing flexible or “open” visitation reduces distress experienced by family members.<sup>39</sup> However, if drafting a policy with visiting hours, feedback from those affected is critical. Consider overlapping with common ICU rounding times to facilitate family member presence on rounds, which can improve the efficiency of communication and integration of the family member into the critical care team.<sup>42</sup> Provide family-centered times that avoid unnecessary gaps during the day and that can accommodate visits by family members traveling from long distances and those with variable work schedules or other caregiving responsibilities to avoid inequity in access.

### *Entry and Screening Procedures*

Establish and communicate a clear structure for visitors to obtain entry. For families of patients with COVID-19, discussion with the family members of their personal risks and benefits of visitation should ideally occur before their arrival at the hospital and be based on their own health conditions, vaccination status, and values. Site-specific legal consultation can inform whether obtaining a liability release and waiver is appropriate. Visitation policies should not restrict on the basis of relationship to patient (eg, immediate family members only), to allow families and patients to identify the most appropriate individual to provide in-person support based on many factors, including personal risk related to COVID-19.

Entry procedures may include a dedicated entrance for visitors, a visitor logging process coordinated with the bedside teams, and standard screening procedures to identify visitor COVID-19 symptoms or high-risk exposures. Provide basic PPE education as they enter, using signs and other environmental cues throughout the hospital to reinforce the need for universal masking, hand washing/sanitizing, and distancing.<sup>61</sup> Dedicating staff to manage visitation logs, visitor screening and approvals, and enforcement of policies using deescalation techniques when necessary will reduce the potential for perceived visitation-related strain on ICU physicians.

### *PPE Use by Visitors*

All family members present in rooms with patients must adhere to recommended PPE guidelines.<sup>32</sup> For patients with COVID-19, this includes N95 masking, eye



protection, and gowning and gloving. Identifying ICU staff to teach and supervise donning and doffing by family members of patients with COVID-19 is essential. Discussions with the health care system's local infection control experts should establish strategies to rapidly assess the fit of N95 models available to visitors. Visitors who are unable or unwilling to maintain PPE in place should be offered alternatives to physical presence, such as videoconferencing.

### *Nonphysical Family Presence*

Providing robust alternatives to in-person family presence is important to ensure high-quality decision-making about the benefits and harms of in-person presence by family members and patients. ICUs should deliver family-centered care regardless of the personal situations and decisions of family members.<sup>4</sup> This is particularly relevant for critically ill patients with COVID-19 because family members may also have COVID-19, be under quarantine, or be at high personal risk of developing severe COVID-19 if exposed to their ill family member. ICUs should capitalize on gains made in the advancement of technological communication strategies during COVID-19 and aim to optimize family engagement through digital platforms, routinized family-physician communication, and extended bereavement support.<sup>4,62</sup> For example, prior work primarily in neonatal ICUs demonstrates that continuous bedside videoconferencing may improve the experiences and outcomes of family members and has no negative effects on physicians.<sup>62-65</sup> Further implementation of such innovative approaches to support the nonphysical presence of family members is relevant during and after visitation restrictions to expand access to family members who feel it is too high risk or who have been exposed to SARS-CoV-2. Because many families of critically ill patients may experience barriers to in-person presence because of work or caregiving responsibilities, geographic distance from the hospital, and transportation or health limitations, maintaining these infrastructures and processes of care will have lasting positive impacts even when the burden of COVID-19 subsides.

### *Communication of the Policy*

The visitation policy itself should make the health care system decision-making process transparent (eg, who was involved in policy decisions), include a clear rationale for the policy, and express concern for the

maintenance of family-centered care. These elements are important for staff as well as families and patients.

Written communication of the policy should be publicly available, including to all patients, and target a fifth to eighth grade reading level.<sup>66,67</sup> On the basis of the individual health care system's patient population, non-English language versions of the policy should be readily accessible. We recommend that all health care systems provide Spanish language materials because of the disproportionate effects of COVID-19 on individuals identifying as Latinx or Hispanic.<sup>68,69</sup>

On admission, a designated staff member or physician should explain the visitation policy to patients and family members, as applicable. Staff should recommend that family members evaluate their personal potential for harm and benefit from visiting a patient with COVID-19, as well as their willingness to take the necessary steps to mitigate risks. Alternatives to in-person visitation should be clearly explained and supported, including establishing plans for routine family-physician communication and family-patient engagement.<sup>41</sup> If visitation prohibitions are in place, the health care system should provide families and patients with technological resources and support to overcome any barriers to such routinized communication.<sup>41,70</sup>

### *Navigating Exceptions*

Although we encourage health care systems to establish permissive visitation policies, it is still necessary to avoid gatherings of people while community transmission rates of SARS-CoV-2 are high. Therefore, visitation policies are likely to have some restrictions in place, such as the number of visitors at a time or throughout the hospitalization. Navigating individual exceptions to the policies established by the health care system requires particular attention to reduce inequity, prevent moral distress among physicians, and protect relationships. Clear processes for adjudicating and granting exceptions will help avoid conflict within clinical teams or across units.

We recommend that health care systems separate clinical teams from visitation adjudication when possible, akin to proposed allocation of scarce resource plans.<sup>56</sup> Such centralized decision-making can preserve the clinical relationship, avoid exacerbating moral distress experienced by physicians, and promote equity.<sup>56</sup> However, input from clinical leaders and bedside physicians, patient advocates, and ethicists is

important to maintain acceptable and appropriate procedures. When considering a case-by-case exception to a standard policy, attention should be given to (1) what specific situational attributes make the exception necessary; (2) the potential impact on other patients, visitors, and staff; (3) whether the exception could be reasonably applied to all patients in similar situations; and (4) how individual biases influence the desire to grant an exception (A. Narva, JD, MSN, email communication, March 2021). Because biases may lead directly to perpetuating or reinforcing health inequities for disenfranchised or frequently underresourced groups, decision makers should carefully interrogate this before granting (or not granting) exceptions.

For example, consider a patient with two adult children and a spouse who requests an exception to a policy that only two family members may visit during the hospitalization. Similarly, consider an 11-year-old child who asks to visit a critically ill parent, yet a policy does not permit children under 12 years of age to visit. In both cases, the health care system could reasonably support all patients and family members being granted exceptions under such circumstances, and these exceptions would respect relationships, promote patient and family well-being, and cause little additional threat to infection control.

### Other Policy Considerations

Family members of patients with COVID-19 should not eat in the patient room, as this requires removal of PPE. Consider discussions with food services and facilities staff to facilitate options for family members, such as outdoor or designated seating areas and grab-and-go or delivered meal options to avoid cafeteria exposures or congregation.

Patient- and family-centered ICU care includes meeting the spiritual needs of patients.<sup>2</sup> In the United States, clergy should not be considered visitors under federal civil rights protections.<sup>71</sup> Their access to the patient should not be restricted and they should not count toward any visitor limits placed by the health care system.

Patients with disabilities may require accommodations during hospitalization to ensure they receive a high quality of care. In countries such as the United States and the United Kingdom, hospitals are legally obligated to provide such accommodations.<sup>72,73</sup> This includes access to essential supporters or communicators when necessary to provide care during a hospitalization.<sup>73</sup> Health care systems should not consider these supportive, caregiving individuals as visitors.

Health care systems that choose to restrict visitation to patients with COVID-19 should develop clear policies and procedures for removing such patients from isolation.<sup>74</sup> Family members may then visit under the health care system's standard policies once the patients are deemed noninfectious.

Given existing variability in local public health regulations, health care systems may choose to consult with legal counsel to ensure compliance. However, the existing evidence and the family-centered care approach frames family members as essential to the clinical care of critically ill patients. As such, similar to health care workers, they may be exempt from some regulations limiting "congregating" to facilitate caregiving.

### Conclusion

The prolonged COVID-19 pandemic has the potential to set family-centered critical care back decades through ongoing visitation restrictions, with a particularly devastating impact on patients with COVID-19 and their families. Health care systems can now leverage knowledge and evidence gained over the first year of the pandemic to promote the safe return of families to the bedside, which will promote the well-being of patients, families, and physicians.

### Acknowledgments

**Financial/nonfinancial disclosures:** None declared.

**Role of sponsors:** The sponsor had no role in the design of the study or the preparation of the manuscript. The content is solely the responsibility of the authors and does not necessarily represent the views of their institutions or the National Institutes of Health.

### References

1. Davidson JE, Jones C, Biennu OJ. Family response to critical illness: postintensive care syndrome—family. *Crit Care Med.* 2012;40(2):618-624.
2. Davidson JE, Aslakson RA, Long AC, et al. Guidelines for family-centered care in the neonatal, pediatric, and adult ICU. *Crit Care Med.* 2017;45(1):103-128.
3. Harvey MA. The truth about consequences—post-intensive care syndrome in intensive care unit survivors and their families. *Crit Care Med.* 2012;40(8):2506-2507.
4. Hart JL, Turnbull AE, Oppenheim IM, Courtright KR. Family-centered care during the COVID-19 era. *J Pain Symptom Manage.* 2020;60(2):e93-e97.
5. Andrist E, Clarke RG, Harding M. Paved with good intentions: hospital visitation restrictions in the age of coronavirus disease 2019. *Pediatr Crit Care Med.* 2020;21(10):e924-e926.
6. Siddiqi H. To suffer alone: hospital visitation policies during COVID-19. *J Hosp Med.* 2020;15(11):694-695.
7. Darcy Mahoney A, White RD, Velasquez A, Barrett TS, Clark RH, Ahmad KA. Impact of restrictions on parental presence in neonatal intensive care units related to coronavirus disease 2019. *J Perinatol.* 2020;40(suppl 1):36-46.



8. Aziz S, Arabi YM, Alhazzani W, et al. Managing ICU surge during the COVID-19 crisis: rapid guidelines. *Intensive Care Med.* 2020;46(7):1303-1325.
9. Mehrotra P, Shane AL, Milstone AM. Family-centered care and high-consequence pathogens: thinking outside the room. *JAMA Pediatr.* 2015;169(11):985-986.
10. Kerlin MP, Costa DK, Davis BS, Admon AJ, Vranas KC, Kahn JM. Actions taken by US hospitals to prepare for increased demand for intensive care during the first wave of COVID-19: a national survey. *Chest.* 2021;160(2):519-528.
11. Voo TC, Senguttuvan M, Tam CC. Family presence for patients and separated relatives during COVID-19: physical, virtual, and surrogate. *J Bioeth Inq.* 2020;17(4):767-772.
12. Curley MAQ, Meyer EC. Upholding family-centered care in the face of high-consequence pathogens—thinking inside the room. *JAMA Pediatr.* 2016;170(3):298-299.
13. Navathe A, Liao J. Barring visitors from hospitals can worsen patient care during COVID. *The Philadelphia Inquirer.* July 27, 2020. <https://www.inquirer.com/opinion/commentary/hospital-visits-coronavirus-philadelphia-pennsylvania-regulations-20200727.html>. Accessed May 13, 2021.
14. Yardley S, Rolph M. Death and dying during the pandemic. *BMJ.* 2020;369:m1472.
15. Moore B. Dying during Covid-19. *Hastings Cent Rep.* 2020;50(3):13-15.
16. Moore KJ, Sampson EL, Kupeli N, Davies N. Supporting families in end-of-life care and bereavement in the COVID-19 era. *Int Psychogeriatr.* 2020;32(10):1245-1248.
17. Morris SE, Moment A, deLima Thomas J. Caring for bereaved family members during the COVID-19 pandemic: before and after the death of a patient. *J Pain Symptom Manage.* 2020;60(2):e70-e74.
18. Montauk TR, Kuhl EA. COVID-related family separation and trauma in the intensive care unit. *Psychol Trauma.* 2020;12(S1):S96-S97.
19. Sutherland AE, Stickland J, Wee B. Can video consultations replace face-to-face interviews? Palliative medicine and the Covid-19 pandemic: rapid review. *BMJ Support Palliat Care.* 2020;10(3):271-275.
20. Murray PD, Swanson JR. Visitation restrictions: is it right and how do we support families in the NICU during COVID-19? *J Perinatol.* 2020;40(10):1576-1581.
21. Koller DF, Nicholas DB, Goldie RS, Gearing R, Selkirk EK. When family-centered care is challenged by infectious disease: pediatric health care delivery during the SARS outbreaks. *Qual Health Res.* 2006;16(1):47-60.
22. Kuntz JG, Kavalieratos D, Esper GJ, et al. Feasibility and acceptability of inpatient palliative care e-family meetings during COVID-19 pandemic. *J Pain Symptom Manage.* 2020;60(3):e28-e32.
23. Greenberg JA, Basapur S, Quinn TV, Bulger JL, Glover CM, Shah RC. Psychological symptoms among surrogates of critically ill patients during and before the COVID-19 pandemic. *Chest.* 2021;159(6):2318-2320.
24. Azad TD, Al-Kawaz MN, Turnbull AE, Rivera-Lara L. Coronavirus disease 2019 policy restricting family presence may have delayed end-of-life decisions for critically ill patients [published online ahead of print April 8, 2021]. *Crit Care Med.* <https://doi.org/10.1097/CCM.0000000000005044>.
25. Kandori K, Okada Y, Ishii W, Narumiya H, Maebayashi Y, Iizuka R. Association between visitation restriction during the COVID-19 pandemic and delirium incidence among emergency admission patients: a single-center retrospective observational cohort study in Japan. *J Intensive Care.* 2020;8:90.
26. Pun BT, Badenes R, Heras La Calle G, et al. COVID-19 Intensive Care Study Group. Prevalence and risk factors for delirium in critically ill patients with COVID-19 (COVID-D): a multicentre cohort study. *Lancet Respir Med.* 2021;9(3):239-250.
27. Gray N. A year of “I’m sorry”—for your loss and the pandemic rules that cloud hospital care. *Los Angeles Times.* March 22, 2021. <https://www.latimes.com/opinion/story/2021-03-22/pandemic-hospital-visitation-rules>. Accessed May 13, 2021.
28. Pennsylvania Department of Health. Guidance on hospitals’ responses to COVID-19. Updated February 22, 2021. <https://www.health.pa.gov/topics/disease/coronavirus/Pages/Guidance/Hospital-Guidance.aspx>. Accessed May 12, 2021.
29. Narain JP, Dawa N, Bhatia R. Health system response to COVID-19 and future pandemics. *J Health Manage.* 2020;22(2):138-145.
30. Valley TS, Schutz A, Nagle MT, et al. Changes to visitation policies and communication practices in Michigan ICUs during the COVID-19 pandemic. *Am J Respir Crit Care Med.* 2020;202(6):883-885.
31. Centers for Medicare and Medicaid Services. Hospital visitation—phase II visitation for patients who are Covid-19 negative. June 26, 2020. <https://www.cms.gov/files/document/covid-hospital-visitation-phase-ii-visitation-covid-negative-patients.pdf>. Accessed May 12, 2021.
32. Centers for Disease Control and Prevention. Interim infection prevention and control recommendations for healthcare personnel during the coronavirus disease 2019 (COVID-19) pandemic. Updated February 10, 2021. <https://www.cdc.gov/coronavirus/2019-ncov/hcp/infection-control-recommendations.html>. Accessed May 12, 2021.
33. Zhou Q, Gao Y, Wang X, et al. COVID-19 Evidence and Recommendations Working Group. Nosocomial infections among patients with COVID-19, SARS and MERS: a rapid review and meta-analysis. *Ann Transl Med.* 2020;8(10):629.
34. Bergman C, Stall NM, Haimowitz D, et al. Recommendations for welcoming back nursing home visitors during the COVID-19 pandemic: results of a Delphi panel. *J Am Med Dir Assoc.* 2020;21(12):1759-1766.
35. Centers for Medicare and Medicaid Services. Nursing home visitation—COVID-19 (revised). March 10, 2021. <https://www.cms.gov/medicareprovider-enrollment-and-certificationsurveycertificationgeninfo/policy-and-memos-states-and-nursing-home-visitation-covid-19-revised>. Accessed May 12, 2021.
36. Wong P, Liamputtong P, Koch S, Rawson H. Barriers to regaining control within a constructivist grounded theory of family resilience in ICU: living with uncertainty. *J Clin Nurs.* 2017;26(23-24):4390-4403.
37. Jessop ZM, Dobbs TD, Ali SR, et al. Personal protective equipment for surgeons during COVID-19 pandemic: systematic review of availability, usage and rationing. *Br J Surg.* 2020;107(10):1262-1280.
38. Hoernke K, Djellouli N, Andrews L, et al. Frontline healthcare workers’ experiences with personal protective equipment during the COVID-19 pandemic in the UK: a rapid qualitative appraisal. *BMJ Open.* 2021;11(1):e046199.
39. Rosa RG, Falavigna M, da Silva DB, et al; ICU Visits Study Group Investigators and the Brazilian Research in Intensive Care Network (BRICNet). Effect of flexible family visitation on delirium among patients in the intensive care unit: the ICU Visits Randomized Clinical Trial. *JAMA.* 2019;322(3):216-228.
40. Nassar Junior AP, Besen BAMP, Robinson CC, Falavigna M, Teixeira C, Rosa RG. Flexible versus restrictive visiting policies in ICUs: a systematic review and meta-analysis. *Crit Care Med.* 2018;46(7):1175-1180.
41. Taylor SP, Short RT III, Asher AM, Taylor B, Beidas RS. A rapid pre-implementation evaluation to inform a family engagement navigator program during COVID-19. *Implement Sci Commun.* 2020;1(1):110.
42. Davidson JE. Family presence on rounds in neonatal, pediatric, and adult intensive care units. *Ann Am Thorac Soc.* 2013;10(2):152-156.
43. Au SS, Roze des Ordon AL, Parsons Leigh J, et al. A multicenter observational study of family participation in ICU rounds. *Crit Care Med.* 2018;46(8):1255-1262.
44. Gupta PR, Perkins RS, Hascall RL, Shelak CF, Demirel S, Buchholz MT. The effect of family presence on rounding duration in the PICU. *Hosp Pediatr.* 2017;7(2):103-107.
45. Kotfis K, Williams Roberson S, Wilson JE, Dabrowski W, Pun BT, Ely EW. COVID-19: ICU delirium management during SARS-CoV-2 pandemic. *Crit Care.* 2020;24(1):176.
46. LaHue SC, James TC, Newman JC, Esmaili AM, Ormseth CH, Ely EW. Collaborative delirium prevention in the age of COVID-19. *J Am Geriatr Soc.* 2020;68(5):947-949.

47. Wassenaar A, Rood P, Schoonhoven L, et al. The impact of nUrsiNg DEliRium Preventive INterventions in the Intensive Care Unit (UNDERPIN-ICU): a study protocol for a multi-centre, stepped wedge randomized controlled trial. *Int J Nurs Stud.* 2017;68:1-8.
48. Trumello C, Bramanti SM, Ballarotto G, et al. Psychological adjustment of healthcare workers in Italy during the COVID-19 pandemic: differences in stress, anxiety, depression, burnout, secondary trauma, and compassion satisfaction between frontline and non-frontline professionals. *Int J Environ Res Public Health.* 2020;17(22):8358.
49. Ruiz-Fernández MD, Ramos-Pichardo JD, Ibáñez-Masero O, Cabrera-Troya J, Carmona-Rega MI, Ortega-Galán AM. Compassion fatigue, burnout, compassion satisfaction and perceived stress in healthcare professionals during the COVID-19 health crisis in Spain. *J Clin Nurs.* 2020;29(21-22):4321-4330.
50. Vagni M, Maiorano T, Giostra V, Pajardi D. Coping with COVID-19: emergency stress, secondary trauma and self-efficacy in healthcare and emergency workers in Italy. *Front Psychol.* 2020;11:566912.
51. Werner EA, Aloisio CE, Butler AD, et al. Addressing mental health in patients and providers during the COVID-19 pandemic. *Semin Perinatol.* 2020;44(7):151279.
52. Chuang CH, Tseng PC, Lin CY, Lin KH, Chen YY. Burnout in the intensive care unit professionals: a systematic review. *Medicine.* 2016;95(50):e5629.
53. Embriaco N, Azoulay E, Barrau K, et al. High level of burnout in intensivists: prevalence and associated factors. *Am J Respir Crit Care Med.* 2007;175(7):686-692.
54. Thornton KC, Schwarz JJ, Gross AK, et al; Project Emerge Collaborators. Preventing harm in the ICU: building a culture of safety and engaging patients and families. *Crit Care Med.* 2017;45(9):1531-1537.
55. Petkovic J, Riddle A, Akl EA, et al. Protocol for the development of guidance for stakeholder engagement in health and healthcare guideline development and implementation. *Syst Rev.* 2020;9(1):21.
56. Maves RC, Downar J, Dichter JR, et al; ACCP Task Force for Mass Critical Care. Triage of scarce critical care resources in COVID-19: an implementation guide for regional allocation: an expert panel report of the Task Force for Mass Critical Care and the American College of Chest Physicians. *Chest.* 2020;158(1):212-225.
57. Robert R, Kentish-Barnes N, Boyer A, Laurent A, Azoulay E, Reignier J. Ethical dilemmas due to the Covid-19 pandemic. *Ann Intensive Care.* 2020;10(1):84.
58. Iwashyna TJ. Recognizing a patient is acutely dying. *Ann Am Thorac Soc.* 2020;17(10):1195-1198.
59. Singer J, Spiegel JA, Papa A. Preloss grief in family members of COVID-19 patients: recommendations for clinicians and researchers. *Psychol Trauma.* 2020;12(S1):S90-S93.
60. Starr LT, O'Connor NR, Meghani SH. Improved serious illness communication may help mitigate racial disparities in care among black Americans with COVID-19. *J Gen Intern Med.* 2021;36(4):1071-1076.
61. Gravina N, Nastasi JA, Sleiman AA, Matey N, Simmons DE. Behavioral strategies for reducing disease transmission in the workplace. *J Appl Behav Anal.* 2020;53(4):1935-1954.
62. Guttman K, Patterson C, Haines T, et al. Parent stress in relation to use of bedside telehealth, an initiative to improve family-centeredness of care in the neonatal intensive care unit. *J Patient Exp.* 2020;7(6):1378-1383.
63. Weber JC, Sohn K, Sauers-Ford HS, et al. Impact of a parent video viewing program in the neonatal intensive care unit. *Telemed J E Health.* 2021;27(6):679-685.
64. Kubicka Z, Zahr E, Clark P, Williams D, Berbert L, Arzuaga B. Use of an internet camera system in the neonatal intensive care unit: parental and nursing perspectives and its effects on stress. *J Perinatol.* 2021:1-9.
65. Epstein EG, Arechiga J, Dancy M, Simon J, Wilson D, Alhusen JL. Integrative review of technology to support communication with parents of infants in the NICU. *J Obstet Gynecol Neonatal Nurs.* 2017;46(3):357-366.
66. Kutner M, Greenberg E, Jin Y, Paulsen C; Institute of Educational Sciences, National Center for Educational Statistics. The health literacy of America's adults: results from the 2003 National Assessment of Adult Literacy. September 2006. <https://nces.ed.gov/pubs2006/2006483.pdf>. Accessed May 13, 2021.
67. Szmuda T, Özdemir C, Ali S, Singh A, Syed MT, Sloniewski P. Readability of online patient education material for the novel coronavirus disease (COVID-19): a cross-sectional health literacy study. *Public Health.* 2020;185:21-25.
68. Williams DR, Cooper LA. COVID-19 and health equity—a new kind of “herd immunity.” *JAMA.* 2020;323(24):2478-2480.
69. Macias Gil R, Marcelin JR, Zuniga-Blanco B, Marquez C, Mathew T, Piggott DA. COVID-19 pandemic: disparate health impact on the Hispanic/Latinx population in the United States. *J Infect Dis.* 2020;222(10):1592-1595.
70. Fang ML, Canham SL, Battersby L, Sixsmith J, Wada M, Sixsmith A. Exploring privilege in the digital divide: implications for theory, policy, and practice. *Gerontologist.* 2019;59(1):e1-e15.
71. US Department of Health and Human Services. OCR resolves religious discrimination complaints after Maryland and Virginia hospitals ensure patients can receive safe religious visitations during COVID-19. October 20, 2020. <https://www.hhs.gov/about/news/2020/10/20/ocr-resolves-religious-discrimination-complaints-after-maryland-and-virginia-hospitals-ensure.html#:~:text=media%40hhs.gov,OCR%20Resolves%20Religious%20Discrimination%20Complaints%20after%20Maryland%20and%20Virginia%20Hospitals,Religious%20Visitations%20During%20COVID%2D19&text=Shortly%20after%2C%20the%20family%20of,to%20provide%20the%20religious%20sacraments>. Accessed May 12, 2021.
72. Read S, Heslop P, Turner S, et al. Disabled people's experiences of accessing reasonable adjustments in hospitals: a qualitative study. *BMC Health Serv Res.* 2018;18(1):931.
73. US Department of Health and Human Services. OCR resolves complaints after state of Connecticut and private hospital safeguard the rights of persons with disabilities to have reasonable access to support persons in hospital settings during COVID-19. June 9, 2020. <https://www.hhs.gov/about/news/2020/06/09/ocr-resolves-complaints-after-state-connecticut-private-hospital-safeguard-rights-persons.html>. Accessed May 12, 2021.
74. Centers for Disease Control and Prevention. Discontinuation of transmission-based precautions and disposition of patients with SARS-CoV-2 infection in healthcare settings. Updated February 16, 2021. <https://www.cdc.gov/coronavirus/2019-ncov/hcp/disposition-hospitalized-patients.html>. Accessed May 12, 2021.