

# UCSF

## UC San Francisco Previously Published Works

### Title

Contemporary screen time modalities among children 9–10 years old and binge-eating disorder at one-year follow-up: A prospective cohort study

### Permalink

<https://escholarship.org/uc/item/5436f94n>

### Journal

International Journal of Eating Disorders, 54(5)

### ISSN

0276-3478

### Authors

Nagata, Jason M  
Iyer, Puja  
Chu, Jonathan  
[et al.](#)

### Publication Date

2021-05-01

### DOI

10.1002/eat.23489

Peer reviewed



Published in final edited form as:

*Int J Eat Disord.* 2021 May ; 54(5): 887–892. doi:10.1002/eat.23489.

## Contemporary Screen Time Modalities among Children 9–10 years old and Binge-Eating Disorder at One-Year Follow-Up: A Prospective Cohort Study

Jason M. Nagata<sup>1</sup>, Puja Iyer<sup>1</sup>, Jonathan Chu<sup>1</sup>, Fiona C. Baker<sup>2,3</sup>, Kelley Pettee Gabriel<sup>4</sup>, Andrea K. Garber<sup>1</sup>, Stuart B. Murray<sup>5</sup>, Kirsten Bibbins-Domingo<sup>6</sup>, Kyle T. Ganson<sup>7</sup>

<sup>1</sup>Division of Adolescent and Young Adult Medicine, Department of Pediatrics, University of California, San Francisco, San Francisco, California, USA

<sup>2</sup>Biosciences Division, Center for Health Sciences, SRI International, Menlo Park, California, USA

<sup>3</sup>Department of Physiology, School of Physiology, University of the Witwatersrand, Johannesburg, South Africa

<sup>4</sup>Department of Epidemiology, University of Alabama at Birmingham, Birmingham, Alabama, USA

<sup>5</sup>Department of Psychiatry and Behavioral Sciences, University of Southern California, Los Angeles, California, USA

<sup>6</sup>Department of Epidemiology and Biostatistics, University of California, San Francisco, San Francisco, California, USA

<sup>7</sup>Factor-Inwentash Faculty of Social Work, University of Toronto, Toronto, Ontario, Canada

### Abstract

**Objective:** To determine the prospective associations between contemporary screen time modalities in a nationally representative cohort of 9–11-year-old children and binge-eating disorder at one-year follow-up.

**Method:** We analyzed prospective cohort data from the Adolescent Brain Cognitive Development (ABCD) Study (N=11,025). Logistic regression analyses were conducted to estimate associations between baseline child-reported screen time (exposure) and parent-reported binge-eating disorder based on the Kiddie Schedule for Affective Disorders and Schizophrenia (KSADS-5, outcome) at one-year follow-up, adjusting for race/ethnicity, sex, household income, parent education, BMI percentile, site, and baseline binge-eating disorder.

**Results:** Each additional hour of total screen time per day was prospectively associated with 1.11 higher odds of binge-eating disorder at 1-year follow-up (95% CI 1.05–1.18) after adjusting for covariates. In particular, each additional hour of social networking (aOR 1.62, 95% CI 1.18–2.22), texting (aOR 1.40, 95% CI 1.08–1.82), and watching/streaming television shows/movies (aOR 1.39, 95% CI 1.14–1.69) was significantly associated with binge-eating disorder.

**Corresponding Author:** Jason M. Nagata, 550 16<sup>th</sup> Street, 4<sup>th</sup> Floor, Box 0110, San Francisco, California 94158, Telephone: +1 (415) 476-3610, jasonmnagata@gmail.com.

**Conflicts of interest statement:** The authors have no conflict to declare.

**Discussion:** Clinicians should assess screen time usage and binge eating in children and adolescents and advise parents about the potential risks associated with excessive screen time.

### Keywords

Screen time; television; social media; smart phone; binge eating; binge-eating disorder; eating disorder; disordered eating; pediatrics; adolescents

## Introduction

The advancement and accessibility of technology has led to the rapid increase in children and adolescents using screens to interact with the world (Hill et al., 2016). As time spent in front of screens has risen, so have concerns regarding the effects of excessive screen time on young people's wellbeing (Hill et al., 2016; Viner et al., 2019). Recent research has linked excessive screen time with adverse effects on children's health, including depression, anxiety, inattention, poor sleep, and physical inactivity (Lissak, 2018; Viner et al., 2019), although it is increasingly apparent that effects of screen time are nuanced, depending on various factors, such as level of engagement and interaction (Orben & Przybylski, 2019; Przybylski et al., 2020).

One particular area of interest is the relationship between excessive screen time and binge eating. Prior studies have demonstrated links between screen time and snacking (Fiechtner et al., 2018; Kim et al., 2020); however, fewer studies have specifically addressed the relationship between screen time and binge eating. Screen time could be linked to binge eating through overeating in the absence of hunger during passive (as opposed to interactive) screen time (Fiechtner et al., 2018; Kim et al., 2020), binge-watching behaviors (Flayelle et al., 2019; Vizcaino et al., 2020), and negative body image (Dakanalis et al., 2015). Furthermore, while research in this area has progressed from the initial focus on television-watching, screen interactions have continued to diversify rapidly since the advent of video games, texting, and social media, and require further continual investigation to differentiate potential effects.

Moreover, the majority of current literature focuses on older adolescents or adults (Burmeister & Carels, 2014; Smith et al., 2013). However, children's screen time usage increases at the greatest rate in early adolescence (Smink et al., 2012; Twenge & Campbell, 2018). Some studies have examined the cross-sectional association between screen time and binge eating in children and adults (Burmeister & Carels, 2014; Fiechtner et al., 2018; Vizcaino et al., 2020), but few have used longitudinal study designs or focused on younger adolescents. One study analyzing clinical samples of children and adolescents for weight loss treatment found a subgroup (n=15) where eating alone, in some cases while watching television, was associated with binge eating (Tanofsky-Kraff et al., 2007). Another population-based study of adolescents in Minnesota did not find an association between television viewing and binge eating (Neumark-Sztainer et al., 2007). However, there is a paucity of data using large, diverse, longitudinal samples examining the association between specific types of screen time and binge-eating disorder in early adolescents using

the Diagnostic and Statistical Manual, 5<sup>th</sup> Edition (DSM-5) criteria (American Psychiatric Association, 2013).

The objective of this study was to determine the prospective associations between screen time in a population-based, demographically diverse cohort of 9–10-year-old children in the U.S and binge-eating disorder at one-year follow-up. In addition, we sought to identify the specific types of screen time (television, videos, video games, texting, video chat, and social networking) that are associated with binge eating. We hypothesized that excessive screen time, particularly passive forms (television, videos) and those that may exacerbate a negative body image (social networking), would be prospectively associated with binge-eating disorder at one-year follow-up.

## 2. Methods

### 2.1 Study Population

We analyzed prospective cohort data from the Adolescent Brain Cognitive Development (ABCD) Study, a longitudinal study of brain development and health across adolescence consisting of 11,875 children recruited from 21 sites around the U.S. (See (Barch et al., 2018) for descriptions of study sample, recruitment, procedures, and measures). Data analyzed here are from the ABCD 3.0 release for the baseline (2016–2018, ages 9–10 years) and one-year follow-up (2017–2019, ages 10–11 years) assessments. Participants with missing data for baseline screen time (N=63) or binge-eating disorder at one-year follow-up (N=793) were excluded, leaving a total of 11,025 participants in the cohort. For participants with missing covariate data (N=1,016), Gaussian normal regression imputation was used to impute missing covariate data. Centralized institutional review board (IRB) approval was obtained from the University of California, San Diego. Study sites obtained approval from their local IRBs. Caregivers provided written informed consent and each child provided written assent.

### 2.2 Measures

**Exposures: Screen Time**—Screen time was determined using the self-reported ABCD Youth Screen Time Survey. Participants answered questions about typical hours per day spent on six different screen modalities (viewing/streaming television shows or movies, watching/streaming videos [e.g. YouTube], playing videogames, texting, video chatting [e.g. Skype, Facetime], and social networking [e.g. Facebook, Instagram, Twitter]) separately for weekdays and weekend days, based on a previously validated measure (Bagot et al., 2018; Gray et al., 2019; Paulus et al., 2019; Sharif et al., 2010). We calculated a weighted average calculation of the participants' typical weekday and weekend screen time use  $((\text{weekday average} \times 5) + (\text{weekend average} \times 2))/7$  to report a single typical hours per day measure (Guerrero et al., 2019). After obtaining this screen time total for each type of media utilized by the participants, we reported the weighted average as a continuous variable.

**Outcome: Binge-Eating Disorder**—The ABCD Study utilized the Kiddie Schedule for Affective Disorders and Schizophrenia (KSADS-5), a computerized tool for categorizing child and adolescent mental health concerns based on the DSM-5 (American Psychiatric

Association, 2013), for assessment of current binge-eating disorder at baseline and one-year follow-up (Townsend et al., 2020). Parents/caregivers completed the binge-eating disorder modules of the KSADS-5 (assessing frequency, duration, characteristics, and associated distress of their child's binge eating) on behalf of their child given evidence that parents are particularly important reporters for these behaviors in this age range (Barch et al., 2018) and young children may have less insight regarding their eating behaviors (Braet et al., 2007). Using the KSADS-5 computerized scoring system, responses to the interview questions from parents were extrapolated into the diagnosis of current binge-eating disorder from reported symptoms corresponding to the DSM-5 (American Psychiatric Association, 2013).

**Covariates:** Sociodemographic covariates were selected based on previous literature and theory as being potential confounders for the association between screen time and binge-eating disorder (Fiechtner et al., 2018; Neumark-Sztainer et al., 2007; Tanofsky-Kraff et al., 2007). Age (years), sex assigned at birth (female, male (including three intersex-male participants)), race/ethnicity (White, Latino/Hispanic, Black, Asian, Native American, other), past year household income (dollars, six categories), and highest parent education (high school or less vs. college or more) were based on parents' self-report. Body mass index (BMI) was based on measured height and weight ( $BMI = \text{weight}/\text{height}^2$ ) and converted into sex and age-specific percentiles in accordance with CDC growth curves (Centers for Disease Control and Prevention, 2019).

**Statistical Analysis**—Data analysis was performed in 2020 using Stata 15.1 (StataCorp, College Station, TX). Unadjusted logistic regression analyses estimated the association between baseline screen time (total and each type of screen time independently) and binge-eating disorder at one-year follow-up. Multiple logistic regression analyses were conducted to estimate the association between baseline screen time (total and each type of screen time independently; exposure) and binge-eating disorder at one-year follow-up (outcome), adjusting for sex, race/ethnicity, household income, parent education, site, and baseline binge-eating disorder. Some children within the sample were twins or siblings. Sensitivity analyses were conducted including only one sibling per family and findings did not differ; therefore, we present results from the full sample. We tested for effect modification by sex and present sex-stratified results where sex significantly modifies the association between screen time and binge-eating disorder ( $p < 0.05$ ). Propensity weights were applied to yield nationally representative estimates based on the American Community Survey from the US Census (Heeringa & Berglund, 2020).

## Results

Table 1 describes sociodemographic characteristics of the 11,025 participants included. The sample was approximately matched by sex (48.8% female) and was racially and ethnically diverse (47.6% non-White). On average, at baseline, children reported  $4.0 \pm 3.2$  hours (mean  $\pm$  SD) of screen time per day, with the most time spent watching/streaming television shows/movies ( $1.3 \pm 1.3$  hours), playing video games ( $1.1 \pm 1.1$  hours), and watching/streaming videos ( $1.1 \pm 1.2$  hours). At one-year follow-up, 1.1% of participants met criteria for binge-eating disorder.

Table 2 shows logistic regression analyses examining the prospective associations between baseline screen time and binge-eating disorder at one-year follow-up. In unadjusted models, all forms of screen time were associated with binge-eating disorder at one-year follow-up. In models adjusting for covariates, each additional hour of total screen time per week at baseline was prospectively associated with 1.11 higher odds of binge-eating disorder at one-year follow-up (95% confidence interval [CI] 1.05–1.18). The screen time types that were most strongly associated with binge-eating disorder in fully-adjusted models were social networking, texting, and television/movie viewing. There was no evidence of effect modification by sex for any of the screen time exposures (all  $p > 0.05$ ) except for video games ( $p = 0.013$ ). Each additional hour of video games at baseline was prospectively associated with higher odds of binge-eating disorder in females (AOR 1.54, 95% CI 1.11–2.14,  $p = 0.010$ ) but not males (AOR 0.92, 95% CI 0.73–1.16,  $p = 0.486$ ).

## Discussion

In a population-based, demographically diverse cohort of 9–10-year-old children in the U.S., we found that greater screen time was prospectively associated with binge-eating disorder at one-year follow-up. In particular, more time spent social networking, texting, and watching/streaming television were most strongly associated with incident binge-eating disorder.

Our findings confirm those of mostly cross-sectional studies (Burmeister & Carels, 2014; Fiechtner et al., 2018; Vizcaino et al., 2020) in older adolescents or adults (Burmeister & Carels, 2014; Smith et al., 2013) examining the relationship between screen time and binge eating. We add to the prior literature on screen time and binge eating by: 1) using a nationally representative prospective cohort design, 2) focusing on an important developmental period for screen time and binge eating (children 9–10 years old followed for one year), and 3) assessing DSM-5 binge-eating disorder as an outcome. The estimates of daily screen time (four hours per day, on average) and binge-eating disorder (0.7%–1.1%) in the ABCD Study were consistent with those from other epidemiological studies with overlapping age ranges (Fiechtner et al., 2018; Marzilli et al., 2018).

As technology platforms consistently evolve and diversify, the need to study new specific screen time mediums now accessible to children and their impacts on binge eating is important. We find that greater amounts of time spent on social networking, texting, and watching television shows/movies are associated with binge-eating disorder. This finding is similar to prior studies that identified television and social networking as associated with triggering binge-eating episodes (Burmeister & Carels, 2014; Smith et al., 2013; Tanofsky-Kraff et al., 2007). We add to this literature by showing that more time spent texting, a relatively new form of screen time for children, is a potential risk factor for subsequent binge-eating disorder, as well. Of note, we did not find significant associations between video chat or video games (except in females) and binge-eating disorder. These forms of screen time may be more interactive and, thus, children may be less prone to binge eating during these more interactive pursuits (Kim et al., 2020; Yland et al., 2015).

Several mechanisms may help to explain the prospective association between screen time and binge eating, including overeating in the absence of hunger, binge-watching behaviors,

and negative body image. First, children may be more prone to overeating in the absence of hunger while distracted in front of screens (Fiechtner et al., 2018; Kim et al., 2020). Second, binge-watching behaviors may lead to overconsumption and a loss of control, similar to binge-eating behaviors (Flayelle et al., 2019; Vizcaino et al., 2020). Third, adolescents who hold negative feelings towards their own body image are more likely to binge eat, and researchers posit that media or advertising content reflecting an unattainable body ideal may exacerbate binge eating (Dakanalis et al., 2015).

Limitations of the study should be noted. Although we adjusted for several potential confounders, including baseline levels of binge eating and BMI percentile, there is the possibility of residual confounding. There may be a bidirectional relationship between screen time and binge-eating disorder, which should be explored in future research. While the prospective study design improves on prior cross-sectional evidence, given the observational study design, we cannot definitively establish causality. The screen time measures were based on self-report, which could be subject to reporting bias. Future studies could use automated measurements of device use to assess screen time. It is important to note that the effect sizes of the associations between screen time and binge-eating disorder were relatively small. Screen time use and incidence of binge-eating disorder may rise after ages 9–11; thus, studies following the ABCD cohort into later adolescence will be an important area of future research. Although parent and child reports of binge eating tend to have low concordance (Bartholdy et al., 2017; Tanofsky-Kraff et al., 2005), parents are particularly important reporters for eating disorders in this age range (Barch et al., 2018) since young children may have less insight regarding their eating behaviors (Braet et al., 2007). The binge-eating questions came from a reliable and validated tool (KSADS-5) that was based on DSM-5 diagnostic criteria.

## Conclusion

In a population-based, demographically diverse cohort of 9–10-year-old children in the U.S., we found that greater television and social media screen time was prospectively associated with binge-eating disorder at one-year follow-up. Given the rapid rise in screen time and disordered eating (Nagata et al., 2020; Termorshuizen et al., 2020) during the COVID-19 pandemic, future research should study these associations during the pandemic. Health care providers should assess for associations between excess screen time usage and binge eating and advise about potential risks associated with excessive screen time. Professional organizations, such as the American Academy of Pediatrics, should provide further specific guidance for families regarding screen time usage and strategies to prevent binge-eating disorder related to screen time usage (Chassiakos et al., 2016).

## Acknowledgements:

J.M.N. is a recipient of the American Heart Association Career Development Award (CDA34760281). S.B.M. was supported by the National Institutes of Health (K23 MH115184). K.B.D. is supported by the National Institutes of Health (K24DK103992). The ABCD Study was supported by the National Institutes of Health and additional federal partners under award numbers U01DA041022, U01DA041025, U01DA041028, U01DA041048, U01DA041089, U01DA041093, U01DA041106, U01DA041117, U01DA041120, U01DA041134, U01DA041148, U01DA041156, U01DA041174, U24DA041123, and U24DA041147. A full list of supporters is available at <https://abcdstudy.org/nihcollaborators>. A listing of participating sites and a complete listing of the study investigators

can be found at <https://abcdstudy.org/principal-investigators.html>. ABCD consortium investigators designed and implemented the study and/or provided data but did not necessarily participate in analysis or writing of this report.

## Data availability statement:

Data used in the preparation of this article were obtained from the ABCD Study (<https://abcdstudy.org>), held in the NIMH Data Archive (NDA). This is a multisite, longitudinal study designed to recruit more than 10,000 children aged 9–10 years and follow them over 10 years into early adulthood.

## References

- American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders* (5th ed.). American Psychiatric Publishing.
- Bagot KS, Matthews SA, Mason M, Squeglia LM, Fowler J, Gray K, Herting M, May A, Colrain I, Godino J, Tapert S, Brown S, & Patrick K (2018). Current, future and potential use of mobile and wearable technologies and social media data in the ABCD study to increase understanding of contributors to child health. *Developmental Cognitive Neuroscience*, 32, 121–129. 10.1016/j.dcn.2018.03.008 [PubMed: 29636283]
- Barch DM, Albaugh MD, Avenevoli S, Chang L, Clark DB, Glantz MD, Hudziak JJ, Jernigan TL, Tapert SF, Yurgelun-Todd D, Alia-Klein N, Potter AS, Paulus MP, Prouty D, Zucker RA, & Sher KJ (2018). Demographic, physical and mental health assessments in the adolescent brain and cognitive development study: Rationale and description. In *Developmental Cognitive Neuroscience* (Vol. 32, pp. 55–66). Elsevier Ltd. 10.1016/j.dcn.2017.10.010 [PubMed: 29113758]
- Bartholdy S, Allen K, Hodsoll J, O’Daly OG, Campbell IC, Banaschewski T, Bokde ALW, Bromberg U, Büchel C, Quinlan EB, Conrod PJ, Desrivières S, Flor H, Frouin V, Gallinat J, Garavan H, Heinz A, Ittermann B, Martinot JL, ... Schmidt U (2017). Identifying disordered eating behaviours in adolescents: how do parent and adolescent reports differ by sex and age? *European Child and Adolescent Psychiatry*, 26(6), 691–701. 10.1007/s00787-016-0935-1 [PubMed: 28050706]
- Braet C, Soetens B, Moens E, Mels S, Goossens L, & Van Vlierberghe L (2007). Are two informants better than one? Parent-child agreement on the eating styles of children who are overweight. *European Eating Disorders Review*, 15(6), 410–417. 10.1002/erv.798 [PubMed: 17960860]
- Burmeister JM, & Carels RA (2014). Television use and binge eating in adults seeking weight loss treatment. *Eating Behaviors*, 15(1), 83–86. 10.1016/j.eatbeh.2013.10.001 [PubMed: 24411756]
- Centers for Disease Control and Prevention. (2019). A SAS Program for the 2000 CDC Growth Charts (ages 0 to <20 years). <https://www.cdc.gov/nccdphp/dnpao/growthcharts/resources/sas.htm>
- Chassiakos YR, Radesky J, Christakis D, Moreno MA, Cross C, Hill D, Ameenuddin N, Hutchinson J, Boyd R, Mendelson R, Smith J, & Swanson WS (2016). Children and adolescents and digital media. *Pediatrics*, 138(5). 10.1542/peds.2016-2593
- Dakanalis A, Carrà G, Calogero R, Fida R, Clerici M, Zanetti MA, & Riva G (2015). The developmental effects of media-ideal internalization and self-objectification processes on adolescents’ negative body-feelings, dietary restraint, and binge eating. *European Child and Adolescent Psychiatry*, 24(8), 997–1010. 10.1007/s00787-014-0649-1 [PubMed: 25416025]
- Fiechtner L, Fonte ML, Castro I, Gerber M, Horan C, Sharifi M, Cena H, & Taveras EM (2018). Determinants of Binge Eating Symptoms in Children with Overweight/Obesity. *Childhood Obesity*, 14(8), 510–517. 10.1089/chi.2017.0311 [PubMed: 30153037]
- Flayelle M, Canale N, Vögele C, Karila L, Maurage P, & Billieux J (2019). Assessing binge-watching behaviors: Development and validation of the “Watching TV Series Motives” and “Binge-watching Engagement and Symptoms” questionnaires. *Computers in Human Behavior*, 90, 26–36. 10.1016/j.chb.2018.08.022
- Gray JC, Schvey NA, & Tanofsky-Kraff M (2019). Demographic, psychological, behavioral, and cognitive correlates of BMI in youth: Findings from the Adolescent Brain Cognitive Development (ABCD) study. *Psychological Medicine*, 50(9), 1539–1547. 10.1017/S0033291719001545 [PubMed: 31288867]



- Guerrero MD, Barnes JD, Chaput JP, & Tremblay MS (2019). Screen time and problem behaviors in children: Exploring the mediating role of sleep duration. *International Journal of Behavioral Nutrition and Physical Activity*, 16(1), 105. 10.1186/s12966-019-0862-x [PubMed: 31727084]
- Heeringa S, & Berglund P (2020). A Guide for Population-based Analysis of the Adolescent Brain Cognitive Development (ABCD) Study Baseline Data. *BioRxiv*, 2020.02.10.942011. 10.1101/2020.02.10.942011
- Hill D, Ameenuddin N, Chassiakos YR, Cross C, Radesky J, Hutchinson J, Boyd R, Mendelson R, Moreno MA, Smith J, & Swanson WS (2016). Media and young minds. *Pediatrics*, 138(5). 10.1542/peds.2016-2591
- Kim S, Favotto L, Halladay J, Wang L, Boyle MH, & Georgiades K (2020). Differential associations between passive and active forms of screen time and adolescent mood and anxiety disorders. *Social Psychiatry and Psychiatric Epidemiology*, 55(11). 10.1007/s00127-020-01833-9
- Lissak G (2018). Adverse physiological and psychological effects of screen time on children and adolescents: Literature review and case study. *Environmental Research*, 164, 149–157. 10.1016/j.envres.2018.01.015 [PubMed: 29499467]
- Marzilli E, Cerniglia L, & Cimino S (2018). A narrative review of binge eating disorder in adolescence: prevalence, impact, and psychological treatment strategies. *Adolescent Health, Medicine and Therapeutics*, *Volume 9*, 17–30. 10.2147/ahmt.s148050 [PubMed: 29379325]
- Nagata JM, Abdel Magid HS, & Pettee Gabriel K (2020). Screen Time for Children and Adolescents During the Coronavirus Disease 2019 Pandemic. *Obesity*, 28(9), 1582–1583. 10.1002/oby.22917 [PubMed: 32463530]
- Neumark-Sztainer DR, Wall MM, Haines JI, Story MT, Sherwood NE, & van den Berg PA (2007). Shared Risk and Protective Factors for Overweight and Disordered Eating in Adolescents. *American Journal of Preventive Medicine*, 33(5). 10.1016/j.amepre.2007.07.031
- Orben A, & Przybylski AK (2019). Screens, Teens, and Psychological Well-Being: Evidence From Three Time-Use-Diary Studies. *Psychological Science*, 30(5), 682–696. 10.1177/0956797619830329 [PubMed: 30939250]
- Paulus MP, Squeglia LM, Bagot K, Jacobus J, Kuplicki R, Breslin FJ, Bodurka J, Morris AS, Thompson WK, Bartsch H, & Tapert SF (2019). Screen media activity and brain structure in youth: Evidence for diverse structural correlation networks from the ABCD study. *NeuroImage*, 185, 140–153. 10.1016/j.neuroimage.2018.10.040 [PubMed: 30339913]
- Przybylski AK, Orben A, & Weinstein N (2020). How Much Is Too Much? Examining the Relationship Between Digital Screen Engagement and Psychosocial Functioning in a Confirmatory Cohort Study. *Journal of the American Academy of Child and Adolescent Psychiatry*, 59(9), 1080–1088. 10.1016/j.jaac.2019.06.017 [PubMed: 31400437]
- Sharif I, Wills TA, & Sargent JD (2010). Effect of visual media use on school performance: A prospective study. *Journal of Adolescent Health*, 46(1), 52–61. 10.1016/j.jadohealth.2009.05.012
- Smink FRE, Van Hoeken D, & Hoek HW (2012). Epidemiology of eating disorders: Incidence, prevalence and mortality rates. *Current Psychiatry Reports*, 14(4), 406–414. 10.1007/s11920-012-0282-y [PubMed: 22644309]
- Smith AR, Hames JL, & Joiner TE (2013). Status Update: Maladaptive Facebook usage predicts increases in body dissatisfaction and bulimic symptoms. *Journal of Affective Disorders*, 149(1–3), 235–240. 10.1016/j.jad.2013.01.032 [PubMed: 23453676]
- Tanofsky-Kraff M, Goossens L, Eddy KT, Ringham R, Goldschmidt A, Yanovski SZ, Braet C, Marcus MD, Wilfley DE, Olsen C, & Yanovski JA (2007). A Multisite Investigation of Binge Eating Behaviors in Children and Adolescents. *Journal of Consulting and Clinical Psychology*, 75(6), 901–913. 10.1037/0022-006X.75.6.901 [PubMed: 18085907]
- Tanofsky-Kraff M, Yanovski SZ, & Yanovski JA (2005). Comparison of child interview and parent reports of children's eating disordered behaviors. *Eating Behaviors*, 6(1), 95–99. 10.1016/j.eatbeh.2004.03.001 [PubMed: 15567115]
- Termorshuizen JD, Watson HJ, Thornton LM, Borg S, Flatt RE, MacDermod CM, Harper LE, van Furth EF, Peat CM, & Bulik CM (2020). Early impact of COVID-19 on individuals with self-reported eating disorders: A survey of ~1,000 individuals in the United States and the

Netherlands. *International Journal of Eating Disorders*, 53(11), 1780–1790. 10.1002/eat.23353 [PubMed: 32720399]

Townsend L, Kobak K, Kearney C, Milham M, Andreotti C, Escalera J, Alexander L, Gill MK, Birmaher B, Sylvester R, Rice D, Deep A, & Kaufman J (2020). Development of Three Web-Based Computerized Versions of the Kiddie Schedule for Affective Disorders and Schizophrenia Child Psychiatric Diagnostic Interview: Preliminary Validity Data. *Journal of the American Academy of Child and Adolescent Psychiatry*, 59(2), 309–325. 10.1016/j.jaac.2019.05.009 [PubMed: 31108163]

Twenge JM, & Campbell WK (2018). Associations between screen time and lower psychological well-being among children and adolescents: Evidence from a population-based study. *Preventive Medicine Reports*, 12, 271–283. 10.1016/j.pmedr.2018.10.003 [PubMed: 30406005]

Viner RM, Davie M, & Firth A (2019). The health impacts of screen time: a guide for clinicians and parents. [https://www.rcpch.ac.uk/sites/default/files/2018-12/rcpch\\_screen\\_time\\_guide\\_-\\_final.pdf](https://www.rcpch.ac.uk/sites/default/files/2018-12/rcpch_screen_time_guide_-_final.pdf)

Vizcaino M, Buman M, Desroches T, & Wharton C (2020). From TVs to tablets: The relation between device-specific screen time and health-related behaviors and characteristics. *BMC Public Health*, 20(1). 10.1186/s12889-020-09410-0

Yland J, Guan S, Emanuele E, & Hale L (2015). Interactive vs passive screen time and nighttime sleep duration among school-aged children. *Sleep Health*, 1(3), 191–196. 10.1016/j.sleh.2015.06.007 [PubMed: 27540566]

**Table 1.**

Sociodemographic, screen time, and binge-eating characteristics of 11,025 Adolescent Brain Cognitive Development (ABCD) Study participants

| Sociodemographic characteristics (baseline)            | Total         |
|--|---------------|
|  | Mean (SD) / % |
| Age (years)  | 9.95 (0.63)   |
| Sex assigned at birth (%)                              |               |
| Female   | 48.8%         |
| Male <sup>a</sup>                                      | 51.2%         |
| Race/ethnicity (%)                                     |               |
| White  | 52.4%         |
| Latino / Hispanic                                      | 20.1%         |
| Black  | 17.3%         |
| Asian  | 5.5%          |
| Native American  | 3.2%          |
| Other  | 1.5%          |
| Household income (%)                                   |               |
| Less than \$25,000                                     | 18.1%         |
| \$25,000 through \$49,999                              | 20.7%         |
| \$50,000 through \$74,999                              | 18.0%         |
| \$75,000 through \$99,999                              | 15.7%         |
| \$100,000 through \$199,999                            | 20.1%         |
| \$200,000 and greater                                  | 6.7%          |
| Parent with college education or more (%)              | 79.7%         |
| Body mass index (BMI) percentile                       | 62.13 (30.70) |
| <b>Screen time variables (hours per day, baseline)</b> |               |
| Total screen time                                      | 3.99 (3.16)   |
| Television shows/movies                                | 1.31 (1.31)   |
| Videos (e.g. YouTube)                                  | 1.05 (1.18)   |
| Video games  | 1.06 (1.13)   |
| Texting  | 0.24 (0.56)   |
| Video chat   | 0.21 (0.52)   |
| Social networking                                      | 0.13 (0.45)   |
| <b>Binge-eating disorder, DSM-5</b>                    |               |
| Binge-eating disorder, baseline (%)                    | 0.7%          |
| Binge-eating disorder, one-year follow-up (%)          | 1.1%          |

Propensity weights were applied to yield representative estimates based on the American Community Survey from the US Census. SD = standard deviation

<sup>a</sup>Includes three participants whose sex at birth was intersex-male

Associations between baseline screen time and binge-eating disorder at one-year follow-up in the Adolescent Brain Cognitive Development Study

**Table 2.**

|                                   | Binge-eating disorder, unadjusted |                | Binge-eating disorder, adjusted <sup>a</sup> |              |
|-----------------------------------|-----------------------------------|----------------|--|--------------|
|                                   | OR (95% CI)                       | P              | aOR (95% CI)                                 | P            |
| Total screen time (hours per day) | <b>1.13 (1.08–1.17)</b>           | < <b>0.001</b> | <b>1.11 (1.05 – 1.18)</b>                    | <b>0.001</b> |
| Television shows/movies           | <b>1.48 (1.24–1.77)</b>           | < <b>0.001</b> | <b>1.39 (1.14 – 1.69)</b>                    | <b>0.001</b> |
| Videos (YouTube)                  | <b>1.23 (1.05–1.45)</b>           | <b>0.01</b>    | 1.09 (0.89 – 1.32)                           | 0.407        |
| Video games                       | <b>1.22 (1.03–1.44)</b>           | <b>0.019</b>   | 1.13 (0.90 – 1.41)                           | 0.310        |
| Texting                           | <b>1.48 (1.18–1.87)</b>           | <b>0.001</b>   | <b>1.40 (1.08 – 1.82)</b>                    | <b>0.011</b> |
| Video chat                        | <b>1.38 (1.11–1.72)</b>           | <b>0.004</b>   | 1.32 (0.99 – 1.76)                           | 0.057        |
| Social networking                 | <b>1.66 (1.29–2.12)</b>           | < <b>0.001</b> | <b>1.62 (1.18 – 2.22)</b>                    | <b>0.003</b> |

Bold indicates p<0.05

<sup>a</sup>Covariates: race/ethnicity, sex, household income, parent education, BMI percentile, site, and baseline binge-eating disorder