

UCLA

UCLA Previously Published Works

Title

cis-Acting Determinant Limiting Expression of Sphingomyelinase Gene sph2 in *Leptospira interrogans*, Identified with a gfp Reporter Plasmid.

Permalink

<https://escholarship.org/uc/item/7qv2v583>

Journal

Applied and Environmental Microbiology, 84(23)

ISSN

0099-2240

Authors

Matsunaga, James
Haake, David A

Publication Date

2018-12-01

DOI

10.1128/aem.02068-18

Peer reviewed



cis-Acting Determinant Limiting Expression of Sphingomyelinase Gene *sph2* in *Leptospira interrogans*, Identified with a *gfp* Reporter Plasmid

James Matsunaga,^{a,c} David A. Haake^{b,c,d,e}

^aResearch Service, VA Greater Los Angeles Healthcare System, Los Angeles, California, USA

^bDivision of Infectious Diseases, VA Greater Los Angeles Healthcare System, Los Angeles, California, USA

^cDepartment of Medicine, UCLA, David Geffen School of Medicine, Los Angeles, California, USA

^dDepartment of Microbiology, Immunology, and Molecular Genetics, UCLA, Los Angeles, California, USA

^eDepartment of Urology, UCLA, Los Angeles, California, USA

ABSTRACT Many strains of the spirochete *Leptospira interrogans* serovar Pomona express the osmotically inducible sphingomyelinase gene *sph2* at much higher levels than strains from other serovars. We developed a new green fluorescent protein (GFP) reporter plasmid to examine *sph2* gene expression determinants. The vector enables the fusion of the test promoter to the ribosome-binding site and coding region of *gfp*. We fused the *sph2* promoters from the *L. interrogans* serovar Lai strain 56601 and from the *L. interrogans* serovar Pomona strain LC82-25 to *gfp* to examine the molecular determinants of differential *sph2* expression between the two strains. Similar to what was observed with the native *sph2* genes, the introduction of the plasmids into the Lai 56601 strain resulted in near background levels of *gfp* expression from the Lai *sph2* promoter, while the expression from the Pomona *sph2* promoter was high. The expression of both fusions increased at physiologic levels of osmolarity achieved by adding sodium chloride to the culture medium. We examined the role of a 17-bp upstream element found in all *L. interrogans* strains expressing low basal levels of *sph2* and missing from Pomona strains that express *sph2* at high levels. When the 17-bp sequence present upstream of the Lai *sph2* promoter was deleted or scrambled, the fusion expression increased substantially. Conversely, the insertion of the 17-bp sequence upstream of the Pomona *sph2* promoter diminished fusion expression. In contrast, the removal of an insertion sequence-like element that is found only in the Pomona *sph2* upstream sequence had no effect on the expression from the Pomona *sph2* fusion in the Lai strain. These findings demonstrate the utility of the *gfp* reporter plasmid in analyzing gene expression in *L. interrogans*.

IMPORTANCE Genetic tools are needed to examine gene expression in the pathogen *Leptospira interrogans*. We developed a reporter plasmid that replicates in *L. interrogans* with green fluorescent protein (GFP) as the readout of promoter activity. We demonstrated an application of the new reporter plasmid by identifying an upstream element responsible for the poor basal expression of the *sph2* sphingomyelinase gene in an *L. interrogans* serovar Lai strain. This new tool is useful for the discovery of the molecular determinants of *L. interrogans* gene expression.

KEYWORDS *Leptospira*, gene expression, leptospirosis, reporter gene, sphingomyelinase, transcription

Most pathogenic members of the genus *Leptospira* have a bimodal lifestyle. In one phase, leptospires live in moist soil and freshwater bodies located throughout the world (1). At other times, they invade vertebrate hosts to permanently colonize the

Received 23 August 2018 Accepted 25 September 2018

Accepted manuscript posted online 28 September 2018

Citation Matsunaga J, Haake DA. 2018. *cis*-acting determinant limiting expression of sphingomyelinase gene *sph2* in *Leptospira interrogans*, identified with a *gfp* reporter plasmid. *Appl Environ Microbiol* 84:e02068-18. <https://doi.org/10.1128/AEM.02068-18>.

Editor Marie A. Elliot, McMaster University
This is a work of the U.S. Government and is not subject to copyright protection in the United States. Foreign copyrights may apply.
Address correspondence to James Matsunaga, jamesm@ucla.edu.

renal tubules, resulting in long-term shedding into the environment by these carriers via their urine (2). The adaptation to these diverse conditions requires the regulation of large numbers of specialized genes, consistent with the results from whole-genome transcriptional array and sequencing studies of *L. interrogans* incubated under conditions simulating those encountered during its life cycle (3–8). Additionally, different serovars of *Leptospira* are generally associated with specific hosts (9), which may reflect adaptation by differential expression of the same genes in different serovars of *Leptospira*.

One example of a gene expressed at dissimilar levels in different serovars is *sph2*, which encodes an enzyme responsible for most, if not all, of the sphingomyelinase activity of *L. interrogans* (10). *L. interrogans* expresses *sph2* in dialysis membrane chambers implanted into the peritoneal cavities of rats. Additionally, antisera raised against Sph2 detect the antigen in kidney sections from acutely infected hamsters (11) and in urine obtained from leptospirosis patients (12). Sph2 may contribute to the pathogenesis of leptospirosis by adhering to fibronectin, inducing apoptosis of host cells, and triggering the production of proinflammatory cytokines (13–16). Most members of *L. interrogans* serovar Pomona examined to date produce large amounts of *sph2* transcript and Sph2 protein, whereas a strain of *L. interrogans* serovar Lai produces smaller amounts of transcript and undetectable levels of Sph2 during growth in conventional culture medium (10, 11). Strains of *L. interrogans* serovars Copenhageni and Manilae also produce negligible amounts of Sph2 (10).

The genetic basis for the increased expression of *sph2* in Pomona strains is unknown. A molecular epidemiology study with PCR primers encompassing an insertion sequence (IS)-like element located upstream of *sph2* found that the element is widespread among Pomona subtype kennewicki strains (17). The IS-like element is missing from the *sph2* upstream sequences of the four *L. interrogans* strains shown to produce low basal levels of Sph2 (10). For these reasons, we hypothesized that this element is responsible for the high levels of Sph2 produced in the Pomona strains (10). However, it should be noted that this hypothesis was based on observations of a single Pomona strain (10), and Sph2 was not detected in another Pomona strain that has the IS-like element upstream of the coding region (13). Whether other Pomona strains that produce high levels of Sph2 harbor the IS-like element is unknown (11).

Several reporter constructs have been developed for studying the expression of *L. interrogans* genes such as *sph2*. Osmotic induction of the *ligA-ligB* and *sph2* *L. interrogans* promoters was demonstrated by genetic fusions to *gfp* in the nonpathogen *Leptospira biflexa* (18). The *kdp* promoter was fused to the 5' end of the *L. biflexa* β -galactosidase gene *bgaL* and integrated by homologous recombination into the endogenous *bgaL* gene to create a model system for demonstrating positive regulation by KdpE of *L. interrogans* (19). The β -galactosidase gene from *Geobacillus stearothermophilus*, *bgaB*, was fused downstream of the *lig* promoter and 5' untranslated region (UTR) to examine expression in *Escherichia coli*. A mutational analysis of the 5' UTR showed that the secondary structure sequesters the ribosome-binding site (20). In each of these cases, the regulation of pathogenic gene expression was examined in a surrogate host, because a plasmid able to replicate in *L. interrogans* was not available. Promoter fusions to *gfp* or luciferase genes have been introduced into *L. interrogans* by transposition (21–23). However, a mutational analysis of promoters fused to the reporter is not possible with the transposon-based system; the random insertion of the transposon carrying the fusion complicates the interpretation of expression levels, because the location of the transposon in the chromosome may influence fusion expression. To overcome this limitation, we took advantage of the fact that several replicative plasmids for *L. interrogans* have recently been developed (24, 25). For example, pMaORI was constructed by cloning the *rep* and partition loci from phage-like sequences integrated in the *Leptospira mayottensis* genome (24). pMaORI has been shown to be stable in the absence of antibiotic selection in seven strains of *L. interrogans*, the nonpathogen *L. biflexa*, and the intermediate species *Leptospira fainei* and *Leptospira licerasiae*. Here, we adapted the pMaORI plasmid to create a new *gfp*




FIG 1 Western blot analysis of Sph2 production from three strains of *L. interrogans* serovar Pomona. Western blots from three *L. interrogans* Pomona strains grown in EMJH medium were probed with Sph2 and LipL41 antisera. The Sph2 antiserum cross-reacts with SphH. *L. interrogans* serovar Manilae strain L495 was grown in EMJH medium supplemented with 120 mM sodium chloride, and its lysate was included for the comparison of Sph2 levels. Sph2 produced by the L495 strain has a slightly lower molecular weight than Sph2 produced by Pomona strains.

reporter vector that can be used to examine gene expression in *L. interrogans*. We used the *gfp* reporter construct to identify a sequence responsible for the low basal levels of *sph2* expression in a Lai strain of *L. interrogans* compared to that in a Pomona strain that produces large amounts of Sph2.

RESULTS

Previously, strain LC82-25 was the only *L. interrogans* serovar Pomona strain with an IS-like element upstream of *sph2* shown to produce high levels of Sph2 (10). For this reason, we tested two additional Pomona strains with IS-like elements for Sph2 production by Western blotting (Fig. 1). Both the RM211 and P10637-46 strains were isolated from pigs. The immunoblot shows that both strains, like the LC82-25 strain, produced high levels of Sph2 when incubated in EMJH medium. The levels of Sph2 were much higher than that produced by the *L. interrogans* serovar Manilae strain L495 grown in EMJH medium with sodium chloride added to achieve a physiologic osmolarity.

To determine whether transcription from within or adjacent to the IS-like element accounted for the high basal expression of *sph2* in the Pomona LC82-25 strain, we mapped the major 5' end of the *sph2* transcript by 5' rapid amplification of cDNA ends (RACE) (Fig. 2A). The trace shows a single 5' end located 28 nucleotides from the *sph2* start codon in both *L. interrogans* strain LC82-25 grown in EMJH medium and strain Fiocruz L1-130 grown at physiologic osmolarity (Fig. 2A). The mixture of the four nucleotides positioned next to the 5' end likely stems from the terminal transferase activity of some reverse transcriptases (26). The 5' end of the *sph2* transcript was found to be located downstream of putative -10 and -35 *E. coli*-like promoter sequences (Fig. 2B).

We next constructed a *gfp* reporter plasmid to examine *sph2* gene expression in *L. interrogans*. We selected the *gfp* allele demonstrated previously to produce functional green fluorescent protein (GFP) in *L. biflexa* when fused to leptospiral promoters (18, 21). The *gfp* allele includes the S65T "red-shift" mutation and the F64L mutation that increases protein solubility (27), as well as a Shine-Dalgarno sequence complementary to the 3' end of the *L. interrogans* 16S rRNA. To construct the reporter plasmid, we cloned the *gfp* gene, along with its translation initiation region, into the plasmid




FIG 2 Mapping the major 5' end of *sph2* by 5' RACE. (A) The 5' ends of the *sph2* transcripts from *L. interrogans* LC82-25 grown in EMJH medium (top) and *L. interrogans* Fiocruz L1-130 grown in EMJH medium with 80 mM sodium chloride (bottom) were mapped by 5' RACE. Electropherograms from sequencing reactions of the amplicons are shown. The peaks representing the major 5' ends of the *sph2* transcripts are bisected by a vertical line. The sequence complementary to the start codon is boxed. The direction of transcription is indicated by the long arrow. (B) The nucleotide sequence around the major 5' end of *sph2* is shown. The start codon is shown in bold, and the predicted -10 and -35 promoter elements are underlined.

pMaORI, an RP4-based mobilizable plasmid that replicates in *L. interrogans* (24). Four unique restriction sites were available for cloning promoters upstream of *gfp* (Fig. 3). A T7 transcription terminator was inserted downstream of *gfp* to create pRAT723 (Fig. 3).

To minimize the transcription of *gfp* initiating from cryptic vector promoters upstream of *gfp*, an *rrnD* transcription terminator was placed upstream of the multicloning site (Fig. 3). *E. coli* transformed with pRAT723, which lacked the transcription terminator, or pRAT724, which possessed the terminator, was collected by centrifugation of liquid cultures to examine the color of the cell pellet. To quantify the terminator's activity in *L. interrogans*, pRAT723 and pRAT724 were transformed into an *E. coli* strain harboring the RP4 conjugation machinery and transferred by conjugation into the *L. interrogans* serovar Lai 56601 strain. Cultures were grown from four colonies of each transconjugant and were processed for fluorescence measurements. The *rrnD* terminator reduced GFP production by $\sim 50\%$ in both *L. interrogans* strains (Fig. 4). The GFP levels in the absence of the *rrnD* terminator were similar to the levels produced from the *L. interrogans* *lipL41* promoter (Fig. 4).

The *sph2* transcript synthesized by the Lai 56601 strain is likely to have the same 5' end as the *sph2* transcript produced by the Fiocruz L1-130 strain, as the sequences upstream of the coding regions in the two strains are nearly identical. On the basis of the 5' RACE result, we cloned the promoter regions of *sph2* from the Lai 56601 and Pomona LC82-25 strains upstream of *gfp* in pRAT724. The *sph2* sequence from the Lai strain (*sph2*_{Lai}) comprised nucleotides -298 through $+3$ relative to the major 5' end of the transcript. The *sph2* promoter fragment cloned from the Pomona strain (*sph2*_{Pom}) had 5' and 3' ends identical to those of the Lai fragment and included the IS-like element. The *lipL41* promoter was also cloned into pRAT724. The resulting




FIG 3 Construction of *L. interrogans* *gfp* reporter plasmid. The thick arrows denote important open reading frames. The restriction sites shown are unique except for the XmaI sites of pGreenTIR. The numbers next to restriction enzyme names refer to the nucleotide positions after the cleavage site. The transcription terminators and a multicloning site were inserted as double-stranded oligonucleotides with overhangs compatible with the ends of the restriction-digested parent plasmid DNA. *aadA*, spectinomycin resistance; *parA* and *parB*, partition loci; *rep*, replication initiator; ENH, T7 gene 10 translational enhancer; SD, Shine-Dalgarno sequence; T7t, early T7 transcription terminator; *rrnDt*, *E. coli* *rrnD* transcription terminator; RP4 *oriT*, RP4 origin of transfer; R6K *ori*, R6K origin of replication; MCS, multicloning site; *, restriction site subject to *dam* methylation.




FIG 4 Effect of a transcriptional terminator on background *gfp* expression. Fluorescence measurements (arbitrary units) of *L. interrogans* containing plasmids with and without the *E. coli rrnD* transcription terminator upstream of *gfp* are shown. The level of GFP produced from a *lipL41p-gfp* fusion with an upstream *rrnD* transcriptional terminator is shown for comparison. The dashed line indicates the mean background reading obtained with *L. interrogans* carrying the empty *gfp* fusion vector pRAT724. Means \pm standard deviations are plotted ($n = 4$). *P* values were calculated by one-way analysis of variance (ANOVA) with Tukey's honestly significant difference (HSD) *post hoc* test.

plasmids carrying the *sph2p-gfp* and *lipL41p-gfp* fusions, along with pRAT724, were introduced into the Lai and Pomona strains by conjugation. The four plasmids were also introduced into the nonpathogen *L. biflexa*, which lacks the *sph2* gene.

As shown in Fig. 5A, the *gfp* expression from the fusions in the Lai strain reflected the expression observed with native *sph2* in the Lai and Pomona strains in a previous study (10). The GFP levels from the *sph2p_{Lai}-gfp* fusion in the Lai strain incubated in EMJH medium were near background levels. The expression from the *sph2p_{Pom}-gfp* fusion was much higher than from the *sph2p_{Lai}-gfp* fusion (Fig. 5A). The same two fusions were also examined in the Pomona strain. In contrast to its activity in the Lai strain, the *sph2p_{Lai}-gfp* fusion produced large amounts of GFP in the Pomona strain, although not as much as the *sph2p_{Pom}-gfp* fusion (Fig. 5B). These observations suggest that the activity of the (Lai) *sph2* promoter was being suppressed in the Lai strain. Neither of the *sph2* promoter fusions produced GFP above background levels in *L. biflexa* (Fig. 5C). GFP fluorescence was detected with the control *lipL41p-gfp* fusion in all three strains (Fig. 5).

We further examined the activities of the promoters in the Lai strain under different conditions of growth. To examine fusion expression at different cell densities, cultures of the four Lai fusion strains were started at a low cell density. Once the density reached 1×10^7 cells/ml, the GFP production from the fusion strains was examined daily until early stationary phase (2×10^9 cells/ml) was reached (Fig. 6). Prior to fluorescence measurement, the densities of all cultures were adjusted to a uniform optical density. The GFP readings for all fusions, including the promoterless control, rose as the cultures reached stationary phase, although the rise was slightly greater for the *sph2* promoter from the Pomona strain and the *lipL41* promoter (Fig. 6). Microscopic observations revealed that cells taken at early stationary phase were noticeably shorter than cells sampled at earlier time points. Notably, the GFP level from the Lai *sph2* promoter fusion never rose above background levels (Fig. 6).

We next determined whether the increase in native Sph2 levels observed when *L. interrogans* is shifted from EMJH medium to EMJH medium with sodium chloride added to achieve physiologic osmolarity (10) is also observed with the *sph2* fusions. The fusion expression was higher when the transconjugants harboring the *sph2p-gfp* constructs were grown in EMJH medium with sodium chloride added to achieve physiologic osmolarity (Fig. 7). The *sph2p_{Pom}-gfp* fusion generated high levels of GFP in the Lai strain, and the expression was even higher at physiologic osmolarity (Fig. 7). As expected, the expression from the control *lipL41p-gfp* fusion was not affected by the sodium chloride supplement (Fig. 7).




FIG 5 GFP production (arbitrary units) from *sph2p-gfp* fusions in *L. interrogans* and *L. biflexa*. Fusion plasmids were introduced into *L. interrogans* Lai strain 56601 (A), *L. interrogans* Pomona strain LC82-25 (B), and *L. biflexa* strain Patoc I (C) and incubated in EMJH medium with spectinomycin. The dashed lines indicate the mean background readings obtained with transconjugants carrying the promoterless *gfp* fusion vector pRAT724 (none). *, $P < 0.05$; **, $P < 0.01$; NS, not significant by Welch's *t* tests.

To determine whether the IS-like element in the promoter region of *sph2* of the Pomona strain is responsible for its high basal level of expression, the IS-like element (Fig. 8A) was precisely deleted from the *sph2* promoter region in the Pomona strain, and the promoter variant was fused to *gfp*. Surprisingly, the GFP assay showed that the removal of the IS-like element had no significant effect on the expression of the *sph2* promoter fusion when examined in the Lai strain incubated in EMJH medium and in EMJH supplemented with sodium chloride (Fig. 8B).




FIG 6 GFP production (arbitrary units) from *sph2p-gfp* fusions in *L. interrogans* at different cell densities. Three cultures of each of the transconjugants harboring the fusion plasmids were sampled daily starting from a cell density of 1×10^7 cells/ml. The cells were collected by centrifugation and resuspended in PBS to an OD_{420} of 0.4 prior to fluorescence measurement. Means \pm standard deviations ($n = 3$) of cell counts (horizontal) and fluorescence readings (vertical) are shown.

To search for sequences that might account for the differences of basal *sph2* expression levels between the Pomona LC82-25 strain and non-Pomona strains, we aligned the sequences upstream of the *sph2* coding regions from the Pomona LC82-25, Lai 56601, Copenhageni Fiocruz L1-130, and Manilae L495 strains. The 5' ends of the aligned sequences correspond to the 5' ends of the *sph2* promoter fragments cloned into pRAT724. As expected, the multisequence alignment revealed a 321-bp IS-like sequence in the Pomona strain, located from position -216 through -536 relative to the major 5' end of the *sph2* transcript of the Pomona strain (Fig. 8A) (13). Sequence elements characteristic of transposable elements were present: a 17-bp inverted repeat at the ends of the element and a 9-bp direct repeat of host sequences immediately adjacent to the inverted repeat (Fig. 8A).

An additional unique feature of the sequence in the Pomona strain is its lack of a 17-bp segment between positions -178 and -179 relative to the major 5' end of its *sph2* transcript (Fig. 8A). To determine whether this 17-bp sequence was responsible for the low basal *sph2* expression in the Lai strain, the 17-bp sequence was removed from




FIG 7 Regulation of *sph2p-gfp* fusion expression by sodium chloride. Fusion plasmids were introduced into *L. interrogans* Lai strain 56601. Fluorescence (arbitrary units) was measured from transconjugants maintained in EMJH medium (EMJH) or EMJH medium with 120 mM sodium chloride (NaCl) for 4 h. Means \pm standard deviations are plotted ($n = 3$). *, $P < 0.05$; ****, $P < 0.0001$; NS, not significant by Welch's *t* tests.






FIG 8 Effect of mutations introduced upstream of the *sph2* promoter. (A) Multisequence alignment. Sequences upstream of *sph2* from *L. interrogans* strains Fiocruz L1-130 (Copenhageni), LC82-25 (Pomona), 56601 (Lai), and L495 (Manilae) were aligned with Kalign. The aligned sequences extend from the 5' ends of the promoter fragments cloned into pRAT724 to the start codon of *sph2*. Most of the IS-like sequences and sequences further upstream were omitted from the figure for brevity. The sequence and nucleotides unique to the Pomona LC82-25 are shown in red. An outside end of the IS-like element containing an inverted repeat is underlined with a double arrowhead at the end. The duplicated host sequence adjacent to the insertion site of the IS-like element is shaded gray. The key 17-nucleotide sequence is boxed, and the imperfect inverted repeats (IR) overlapping with the key sequence are marked with arrows. The proposed transcription start site is indicated by an arrow at position +1. The *E. coli*-like -10 and -35 promoter sequences, Shine-Dalgarno sequence (SD), and start codons are demarked with dashed boxes. Fusion plasmids were introduced into *L. interrogans* serovar Lai strain 56601. Transconjugants were incubated in EMJH medium (EMJH) or EMJH with 120 mM sodium chloride (NaCl) for 4 h

(Continued on next page)

the upstream region of the Lai *sph2* promoter in the *gfp* fusion. The deletion mutation restored 58% (Fig. 8B), 63%, and 81% of the expression observed with the Pomona *sph2p-gfp* fusion over three experiments. The converse experiment was also performed by inserting the 17-bp segment upstream of the Pomona *sph2* promoter in the fusion. The insertion reduced fusion expression by 68% (Fig. 8B), 70%, and 72%. The expression from all *sph2* fusions was elevated at physiologic osmolarity, indicating that the 17-bp sequence is not a major contributor to osmotic induction of *sph2* expression.

To rule out the possibility that removing the 17-bp sequence altered the spacing between transcriptional control elements, an additional *sph2* promoter variant was constructed by scrambling the 17-bp sequence. The scrambled sequence restored 44% of the expression observed with the Pomona fusion (Fig. 8C), indicating that at least a portion of the 17-bp sequence plays a direct role in limiting the transcription from the *sph2* promoter.

DISCUSSION

Differential expression of the *sph2* sphingomyelinase gene from a Pomona and a non-Pomona strain of *L. interrogans* was examined with a new *gfp* fusion plasmid. This *gfp* fusion plasmid enabled the identification of a segment of the *sph2* promoter in the Lai strain that is primarily responsible for its low levels of *sph2* expression. The Copenhageni and Manilae strains, which also express low basal levels of *sph2*, also possess the 17-bp sequence. The three Pomona strains, which express high levels of Sph2, lack the 17-bp segment. Interestingly, the 17-bp sequence contains one arm of an inverted repeat (Fig. 8A), which may be a binding site for a *trans*-acting factor responsible for the low basal expression of *sph2*. In contrast, an upstream IS-like sequence hypothesized to increase *sph2* expression did not contribute to basal expression from the Pomona *sph2* promoter when examined in the Lai strain.

It is anticipated that this *gfp* reporter strategy can be used to study gene expression in other strains and species of *Leptospira*. The parent of pRAT724, pMaORI, stably replicates in at least five other strains of *L. interrogans* and in *L. mayottensis*, *L. licerasiae*, *L. fainei*, and *L. biflexa* (24). pMaORI carrying the *fliM* gene also restored the virulence of a spontaneous motility mutant of *L. interrogans*, indicating that pMaORI is stable *in vivo* in the absence of antibiotic selection (28). Nevertheless, future efforts to examine the expression of promoter fusions to *gfp in vivo* will require additional experiments to verify the stability of the fusion plasmid in the absence of antibiotics.

One limitation of the new fusion plasmid is that variability and irreproducibility in the GFP levels were observed at high levels of expression of the *sph2* promoter fusions in the Pomona strain. For this reason, although the IS-like element did not contribute to *sph2* expression in the Lai strain, we cannot firmly exclude the possibility that the IS-like element contributes to basal *sph2* expression in the Pomona strain. Additionally, we observed increases in GFP from all fusions in cells entering stationary phase (Fig. 6). Because cell densities across cultures were normalized on the basis of optical density rather than cell number prior to fluorescent measurement, the cell shrinkage we observed in cultures entering stationary phase may have contributed to the increased GFP readings. Further studies are needed to determine whether an increase in plasmid copy number per cell during entry into stationary phase also contributed to the increase. Nevertheless, the development of this *gfp* fusion plasmid will facilitate the identification of *cis*-acting sequences critical for the expression of *L. interrogans* genes. The recent publication of the genome-wide transcription start sites for *L. interrogans* will aid the selection of the correct junction for promoter fusions to *gfp* (29). A transcription start site was

FIG 8 Legend (Continued)

(B) or in EMJH medium only (C). Fluorescence is shown in arbitrary units. The dashed lines indicate the mean background readings obtained with *L. interrogans* carrying the empty *gfp* fusion vector pRAT724. Means \pm standard deviations are plotted ($n = 3$). *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$; NS, not significant by Welch's *t* test. S17, 17-bp sequence scrambled.

not detected for *sph2* in the study, consistent with the limited expression of *sph2* in non-Pomona strains under standard culture conditions.

The elucidation of the mechanism for differential *sph2* gene expression among *L. interrogans* strains raises the interesting question of whether the 17-bp promoter deletion provides a selective benefit to these strains. Increased *sph2* expression appears to be unique among Pomona strains (10, 11). In contrast, *sph2* is expressed poorly in strains from serovars Copenhageni, Lai, and Manilae, whose reservoir hosts are small rodents (30–32). The association of *L. interrogans* serovar Pomona infection with sheep and pigs (9) suggests that high levels of *sph2* expression confer an adaptive benefit to the Pomona serovar for the infection of specific maintenance hosts. Additional studies comparing infection in different hosts with leptospiral strains with and without the 17-bp deletion would be helpful in understanding the role of Sph2, assessing whether there is a benefit of increased *sph2* expression in some hosts, and determining whether that benefit is host specific.

The *sph2* fusion constructs described here may be helpful in identifying the *cis*-acting site(s) involved in the osmotic regulation of *sph2*. In a previous study, the addition of 120 mM sodium chloride to cultures increased *sph2* transcript levels by over 100-fold in both *L. interrogans* serovar Manilae and *L. interrogans* serovar Pomona (10). The fact that these increases were similar in both low and high Sph2 producers suggests that the 17-bp *cis*-acting element is not involved in the osmotic regulation of transcription. A previous study with the 600 bp immediately upstream of the *sph2* coding region from the Fiocruz L1-130 strain of *L. interrogans* fused to *gfp* demonstrated the osmotic induction of the fusion in *L. biflexa* (18). We observed osmotic induction with a fragment extending from 24 to 325 nucleotides upstream of the *sph2* coding region fused to *gfp*. Therefore, the 302-bp region serves as a starting point to locate the regulatory target. The observation that osmotic induction occurs in the nonpathogen *L. biflexa* suggests that a *trans*-acting determinant of *sph2* osmotic regulation is conserved between at least some pathogenic and nonpathogenic *Leptospira* species.

Little is known about how *cis*- and *trans*-acting factors establish the level of gene expression in *Leptospira*. Several *cis*-acting sequences affecting the expression of *Leptospira* genes involving adaptation to different environmental stresses have been identified. The *L. interrogans* genome encodes two *lexA* genes (33). DNase I footprinting and gel shift assays with the *recA* upstream sequence revealed an SOS box bound by LexA1 (34). Additional gel shift assays demonstrated that similar sequences upstream of other SOS-induced *L. interrogans* genes bound to LexA1 (33). LexA2 bound only to its own promoter (33). In another study, a high-affinity binding site for the phosphorylated form of the *L. biflexa* HemR response regulator, which regulates heme metabolism, was enriched from a highly randomized oligonucleotide library by several rounds of affinity selection with the phosphorylated protein (35). In a recent study, the *L. interrogans* enhancer-binding protein EBP and the alternative sigma factor RpoN were shown to bind to various promoters by gel shift analysis (36). For *L. interrogans* genes such as *sph2* for which the *trans*-acting regulatory factor remains to be identified, the pRAT724 *gfp* reporter plasmid will facilitate the identification of additional *cis*-acting sequences affecting gene expression.

MATERIALS AND METHODS

Bacterial strains. *Leptospira interrogans* serovar Lai strain 56601 (37), *L. interrogans* serovar Copenhageni strain Fiocruz L1-130 (38), and *L. interrogans* serovar Pomona subtype kennewicki strains LC82-25, RM211, and P10637-46 (National Animal Disease Center, Ames, IA) were grown in EMJH medium (Probumin vaccine-grade solution, catalog number 840665, lot 103; Millipore) at 30°C with 40 µg/ml spectinomycin for plasmid selection. *E. coli* strains DH5 α , π 1, and β 2163 (39) were incubated in LB with the appropriate antibiotic added to select for plasmids. The LB medium for *E. coli* π 1 and β 2163 was supplemented with 0.3 mM thymidine and 0.3 mM diaminopimelic acid, respectively (39).

Western blots. Immunoblot analyses of *L. interrogans* lysates were performed as described previously (20). The blots were probed with 1:1,000 dilution of Sph2 antiserum and 1:5,000 dilution of LipL41 antiserum (40, 41).

Plasmid construction. The plasmids used in the study are listed in Table 1. Plasmid DNA was extracted from 4 ml of overnight *E. coli* cultures with the Qiagen QIAprep Spin Miniprep kit (Valencia, CA) and eluted with 50 µl of 10 mM Tris-HCl, 0.1 mM EDTA, pH 8.5. Genomic DNA was purified from 5 ml of saturated *L. interrogans* cultures with the Promega Wizard genomic DNA purification kit and resuspended

TABLE 1 Plasmids used in this study

Plasmid	Description	Source or reference
pMaORI	Replicative plasmid for <i>L. interrogans</i>	24
pGreenTIR	<i>gfp</i> vector	27
pBluescript SK ⁻	Cloning vector	Stratagene
pRAT720	pMaORI with multicloning site	This study
pRAT722	pGreenTIR with T7 early transcription terminator downstream of <i>gfp</i>	This study
pRAT723	<i>gfp</i> vector in pMaORI, no <i>rrnD</i> transcription terminator	This study
pRAT724	<i>gfp</i> vector in pMaORI	This study
pRAT725	<i>lipL41p-gfp</i> in pRAT724	This study
pRAT727	<i>sph2p_{Lai}-gfp</i> in pRAT724	This study
pRAT728	<i>sph2p_{Pom}-gfp</i> in pRAT724	This study
pRAT729	<i>sph2p_{Pom}ΔIS-gfp</i> in pRAT724	This study
pRAT735	<i>sph2p_{Lai}</i> in pBluescript SK ⁻	This study
pRAT736	<i>sph2p_{Pom}</i> in pBluescript SK ⁻	This study
pRAT739	<i>sph2p_{Pom}(+17)-gfp</i> in pRAT724	This study
pRAT740	<i>sph2p_{Lai}(Δ17)-gfp</i> in pRAT724	This study

in 50 μ l of 1 mM Tris-HCl, 0.1 mM EDTA, pH 7.4. Restriction enzymes and Quick ligase were supplied by New England BioLabs (Ipswich, MA). Synthetic oligonucleotides were obtained from Invitrogen, and the nucleotide sequences are provided in Table 2. PCRs were performed with *Pfu* DNA polymerase according to the manufacturer's instructions (Thermo Fisher Scientific).

The *gfp* reporter plasmid was generated as follows (Fig. 1). A double-stranded oligonucleotide harboring the T7 early transcription terminator with 5' KpnI and 3' SphI single-stranded overhangs (Table 2) was inserted downstream of *gfp* in pGreenTIR (27) to create pRAT722. Additionally, a multicloning site was introduced into the plasmid pMaORI (24) by replacing its small BamHI-SalI fragment with the double-stranded DNA 5'-GATCCGGTACCTAGTCTCGAGCCCGGGCATGCG-3'/5'-TCGACGCATGCCCGGGCTCGAGACTAGTGGTACCG-3' (BamHI and SalI overhangs underlined), resulting in pRAT720. The *gfp* ribosome-binding site and coding region along with the 3' T7 early terminator were excised from pRAT722 and inserted into XmaI-SphI-cut pRAT720 to generate pRAT723. Finally, a double-stranded oligonucleotide carrying the *E. coli rrnD* transcription terminator with 5' BamHI and 3' SpeI cohesive ends and BamHI and KpnI sites 3' of the terminator was inserted upstream of *gfp* sequences in pRAT723 to generate pRAT724.

To clone the *lipL41* and *sph2* promoters, PCR primers were designed with BamHI and XhoI restriction sites near the 5' ends of the upstream and downstream primers, respectively (Table 2). The *lipL41* promoter was PCR amplified using *L. interrogans* strain Fiocruz L1-130 genomic DNA as the template. The *sph2* promoters were amplified using *L. interrogans* serovar Lai strain 56601 and *L. interrogans* serovar Pomona strain LC82-25 genomic DNA as the templates. The PCR products were digested with BamHI and XhoI and inserted into the corresponding site in pRAT724 to create the promoter fusions to *gfp*.

Mutagenesis. The template for oligonucleotide-directed mutagenesis of the sequence upstream of the *sph2* promoter was generated by transferring the BamHI-XhoI fragment containing the promoter sequence from the wild-type *sph2-gfp* fusion plasmids into pBluescript SK⁻ (Stratagene) to generate pRAT735 and pRAT736 (Table 1), which harbor the *sph2* promoters from the Lai 56601 and Pomona LC82-25 strains, respectively. The desired mutations upstream of the *sph2* promoter were generated as described previously (20, 42). In brief, overlapping oligonucleotides with the desired changes directed toward each strand (Table 3) were used to PCR amplify the plasmid with *Pfu* DNA polymerase. The template DNA was depleted by digestion with DpnI, and the amplicon was transformed into *E. coli* DH5 α . The transformation culture was plated on LB plates containing 100 μ g/ml ampicillin.

TABLE 2 Synthetic oligonucleotides used for plasmid constructions

Oligonucleotide	Sequence (5'→3') ^a	Description
T7Te(Kp)-5F	TAATCACACTGGCTCACCTTCGGGTGGGCCCTTTCTGCGTTTATAAGGAGCATG	T7 early transcription terminator
T7Te(Sp)-5R	CTCCTTATAAACGCAGAAAGGCCACCCGAAGGTGAGCCAGTGTGATTAGTAC	
<i>rrnD</i> (Bm)-2F	GATCTCAAATAAAAACAAAAGGCTCAGTCGGAAGACTGGGCCCTTTGTTTTATCTGTTGGATCCGGTACCA	<i>E. coli rrnD</i> transcription terminator
<i>rrnD</i> (Se)-2R	CTAGTGGTACCGGATCCAACAGATAAAAACAAAAGGCCAGTCTCCGACTGAGCCTTTGTTTTATTGA	
<i>lipL41p</i> (Bm)-3F	GTTCCAGGATCCCTTGTAAATTCAGTATCTGTATGAGAAGT	<i>lipL41</i> promoter
<i>lipL41p</i> (Xh)-4R	CACCAA CTCGAG TTGATTTTGGGGAATAAGG	
la1029(Bm)-2F	GACCATGGATCCAGCGAGACGCTTGAGTCTGA	<i>sph2_{Lai}</i> promoter
lic12631(Bm)-32F	GACCATGGATCCAGCGAGACGCTTGAGTCTGA	<i>sph2_{Pom}</i> promoter
lic12631p(Xh)-2R	TCCAAT CTCGAG GTTTGTATTTTATGTTATCATATAACATAAAAATG	<i>sph2</i> promoter

^aOverhangs are underlined; restriction sites are in bold font.

TABLE 3 Oligonucleotides used for site-directed mutagenesis

Oligonucleotide	Sequence (5'→3') ^a
lip0980p(−206i)-1F	TATTTAAAATAACATTCAATTTAAATAATATTATAATAAAAATAAAATTTTATAAAAAATACA
lip0980p(−206i)-1R	AATTGAATGTTATTTTAAATAATTTTAGAGAAAGCGGATGATAG
la1029p(−222/−206d)-1F	CCTCAAATTA TTTAAAGAATATTATAATAAAAATAAAATTTTATAAAAAATACA
la1029p(−222/−206d)-1R	ATTCITTTAAA TAATTTGAGGACAAGTGGATG
la1029p(−222/−206s)-2F	CCTCAAATTA TGATTTTTAGAATAATTTTAAAGAATATTATAATAAAAATAAAATTTTATAAAAAATACA
la1029p(−222/−206s)-2R	ATTCITTTAAA ATTATTCTAAAAATCATAATTTGAGGACAAGTGGATG

^aInserted nucleotides are underlined; deletions introduced at vertical lines; scrambled sequences are in bold font.

Plasmid DNA was purified from overnight cultures of colonies, and the presence of the desired mutations was confirmed by Sanger sequencing with the T7 and T3 sequencing primers (Laragen, Culver City, CA).

Conjugation. The *gfp* fusion plasmids were transferred into *L. interrogans* by conjugation from a donor *E. coli* strain as described (43). Briefly, *E. coli* β 2163 was transformed with the *gfp* fusion plasmids and grown overnight in LB with 0.3 mM diaminopimelic acid (DAP) and 40 μ g/ml spectinomycin at 37°C. The next day, 40 μ l of the overnight culture was transferred to 4 ml of LB with DAP and grown to late exponential phase. *L. interrogans* was grown to a density of $\approx 10^8$ cells/ml. Five milliliters of *L. interrogans* and 0.5 ml *E. coli* were mixed and filtered with a 0.1- μ m filter (Fisher). The filter was placed face up on an EMJH plate containing 0.3 mM DAP and incubated overnight at 30°C. The bacteria were washed off the filter with 1 ml of EMJH medium and plated on three EMJH plates containing 40 μ g/ml spectinomycin.

5' end mapping of RNA. A culture of *L. interrogans* Fiocruz L1-130 was started at 2×10^7 cells/ml in 25 ml EMJH medium supplemented with 80 mM NaCl and incubated for 5 days to a density of 6×10^8 cells/ml. Additionally, a culture of *L. interrogans* LC82-25 was initiated in 25 ml EMJH at 1×10^7 cells/ml and incubated for 3 days to a density of 5×10^8 cells/ml. RNA was extracted from *L. interrogans* with TRIzol (Invitrogen), and DNA was removed with Turbo DNase (Ambion). The 5' end of the *sph2* transcript was determined with the Roche 5'/3' RACE kit, 2nd generation (Roche), using oligonucleotide sph2-8R (5'-CGTTTGGCTCTTTCATCGTGTC-3') to prime the reverse transcription reaction and sph2-9R (5'-GCGGAGCTGCCATTTCTG-3') to PCR amplify the deoxyribosyladenine (dA)-tailed *sph2* cDNA. The amplicon was sequenced with the sph2-9R primer to map the 5' end.

GFP assay. *L. interrogans* transconjugants carrying *gfp* fusion plasmids were grown in EMJH medium with 40 μ g/ml spectinomycin to a density of $\approx 1 \times 10^8$ to 5×10^8 cells/ml, and the optical density at 420 nm (OD_{420}) of the cultures was measured with an Ultrospec 2000 spectrophotometer (Pharmacia Biotech). Eight hundred microliters of each culture was collected by centrifugation for 5 min at $9,000 \times g$ in an Eppendorf 5424 microcentrifuge. The cells were resuspended in a volume of phosphate-buffered saline (PBS) to obtain an OD_{420} reading of 0.4 (4 ml). One hundred microliters of each cell suspension was transferred to a dark-walled clear-bottom 96-well Costar plate (Thermo Fisher Scientific), and the fluorescence emitted at 528 nm (excitation 485 nm) was measured with a Synergy2 Multi-Mode microplate reader (BioTek, Winooski, VT, USA). Because the fluorescence readings of wild-type *L. interrogans* were similar to those of PBS, PBS was used as a blank for background subtraction. Fluorescence values from PBS were subtracted from the readings of *L. interrogans* carrying the *gfp* fusion plasmids.

Multisequence alignment. The *sph2* promoter sequences from *L. interrogans* strains Fiocruz L1-130, L495, 56601, and LC82-25 were aligned with Kalign (44).

Statistics. Experiments were conducted with cultures initiated from three colonies, unless stated otherwise. Statistical analysis was conducted with R, version 3.4.2, "Short Summer" (45).

ACKNOWLEDGMENTS

We thank William G. Miller for providing pGreenTIR and Mathieu Picardeau for contributing pMaORI. We also thank Rich Zuerner for supplying the *L. interrogans* serovar Pomona strains. We thank Suneel Narayanavari for technical assistance and David Beenhouwer for use of the fluorescence microplate reader.

The study was funded by Veterans Affairs merit awards (to J.M. and D.A.H.) and a Public Health Service grant R21 AI128560-01A1 from the National Institute of Allergy and Infectious Diseases (to D.A.H.).

J.M. conceived the study, contributed materials, designed and performed the experiments, analyzed the data, and wrote the paper. D.A.H. helped analyze the data and write the paper.

REFERENCES

- Guerra MA. 2009. Leptospirosis. *J Am Vet Med Assoc* 234:472–478. <https://doi.org/10.2460/javma.234.4.472>.
- Babudieri B. 1958. Animal reservoirs of leptospire. *Ann N Y Acad Sci* 70:393–413. <https://doi.org/10.1111/j.1749-6632.1958.tb35398.x>.
- Caimano MJ, Sivasankaran SK, Allard A, Hurley D, Hokamp K, Grassmann AA, Hinton JC, Nally JE. 2014. A model system for studying the transcriptomic and physiological changes associated with mammalian host-adaptation by *Leptospira interrogans* serovar Copenhageni. *PLoS Pathog* 10:e1004004. <https://doi.org/10.1371/journal.ppat.1004004>.
- Iraola G, Spangenberg L, Lopes Bastos B, Grana M, Vasconcelos L,

- Almeida A, Greif G, Robello C, Ristow P, Naya H. 2016. Transcriptome sequencing reveals wide expression reprogramming of basal and unknown genes in *Leptospira biflexa* biofilms. *mSphere* 1:e00042-16. <https://doi.org/10.1128/mSphere.00042-16>.
5. Lo M, Bulach DM, Powell DR, Haake DA, Matsunaga J, Paustian ML, Zuerner RL, Adler B. 2006. Effects of temperature on gene expression patterns in *Leptospira interrogans* serovar Lai as assessed by whole-genome microarrays. *Infect Immun* 74:5848–5859. <https://doi.org/10.1128/IAI.00755-06>.
 6. Lo M, Murray GL, Khoo CA, Haake DA, Zuerner RL, Adler B. 2010. Transcriptional response of *Leptospira interrogans* to iron limitation and characterization of a PerR homolog. *Infect Immun* 78:4850–4859. <https://doi.org/10.1128/IAI.00435-10>.
 7. Matsunaga J, Lo M, Bulach DM, Zuerner RL, Adler B, Haake DA. 2007. Response of *Leptospira interrogans* to physiologic osmolarity: relevance in signaling the environment-to-host transition. *Infect Immun* 75:2864–2874. <https://doi.org/10.1128/IAI.01619-06>.
 8. Patarakul K, Lo M, Adler B. 2010. Global transcriptomic response of *Leptospira interrogans* serovar Copenhageni upon exposure to serum. *BMC Microbiol* 10:31. <https://doi.org/10.1186/1471-2180-10-31>.
 9. Bharti AR, Nally JE, Ricaldi JN, Matthias MA, Diaz MM, Lovett MA, Levett PN, Gilman RH, Willig MR, Gotuzzo E, Vinetz JM. 2003. Leptospirosis: a zoonotic disease of global importance. *Lancet Infect Dis* 3:757–771. [https://doi.org/10.1016/S1473-3099\(03\)00830-2](https://doi.org/10.1016/S1473-3099(03)00830-2).
 10. Narayanavari SA, Lourdault K, Sritharan M, Haake DA, Matsunaga J. 2015. Role of *sph2* gene regulation in hemolytic and sphingomyelinase activities produced by *Leptospira interrogans*. *PLoS Negl Trop Dis* 9:e0003952. <https://doi.org/10.1371/journal.pntd.0003952>.
 11. Carvalho E, Barbosa AS, Gomez RM, Oliveira ML, Romero EC, Goncalves AP, Morais ZM, Vasconcelos SA, Ho PL. 2010. Evaluation of the expression and protective potential of leptospiral sphingomyelinases. *Curr Microbiol* 60:134–142. <https://doi.org/10.1007/s00284-009-9519-3>.
 12. Chaurasia R, Thresiamma KC, Eapen CK, Zachariah BJ, Paul R, Sritharan M. 2018. Pathogen-specific leptospiral proteins in urine of patients with febrile illness aids in differential diagnosis of leptospirosis from dengue. *Eur J Clin Microbiol Infect Dis* 37:423–433. <https://doi.org/10.1007/s10096-018-3187-9>.
 13. Artiushin S, Timoney JF, Nally J, Verma A. 2004. Host-inducible immunogenic sphingomyelinase-like protein, Lk73.5, of *Leptospira interrogans*. *Infect Immun* 72:742–749. <https://doi.org/10.1128/IAI.72.2.742-749.2004>.
 14. Pinne M, Matsunaga J, Haake DA. 2012. Leptospiral outer membrane protein microarray, a novel approach to identification of host ligand-binding proteins. *J Bacteriol* 194:6074–6087. <https://doi.org/10.1128/JB.01119-12>.
 15. Wang H, Wu Y, Ojcius DM, Yang XF, Zhang C, Ding S, Lin X, Yan J. 2012. Leptospiral hemolysins induce proinflammatory cytokines through Toll-like receptor 2- and 4-mediated JNK and NF- κ B signaling pathways. *PLoS One* 7:e42266. <https://doi.org/10.1371/journal.pone.0042266>.
 16. Zhang YX, Geng Y, Yang JW, Guo XK, Zhao GP. 2008. Cytotoxic activity and probable apoptotic effect of Sph2, a sphingomyelinase hemolysin from *Leptospira interrogans* strain Lai. *BMB Rep* 41:119–125. <https://doi.org/10.5483/BMBRep.2008.41.2.119>.
 17. Timoney JF, Kalimthusamy N, Velineni S, Donahue JM, Artiushin SC, Fetting M. 2011. A unique genotype of *Leptospira interrogans* serovar Pomona type kennewicki is associated with equine abortion. *Vet Microbiol* 150:349–353. <https://doi.org/10.1016/j.vetmic.2011.02.049>.
 18. Cerqueira GM, Souza NM, Araujo ER, Barros AT, Morais ZM, Vasconcelos SA, Nascimento AL. 2011. Development of transcriptional fusions to assess *Leptospira interrogans* promoter activity. *PLoS One* 6:e17409. <https://doi.org/10.1371/journal.pone.0017409>.
 19. Matsunaga J, Coutinho ML. 2012. Positive regulation of *Leptospira interrogans kdp* expression by KdpE as demonstrated with a novel β -galactosidase reporter in *Leptospira biflexa*. *Appl Environ Microbiol* 78:5699–5707. <https://doi.org/10.1128/AEM.00713-12>.
 20. Matsunaga J, Schlax PJ, Haake DA. 2013. Role for *cis*-acting RNA sequences in the temperature-dependent expression of the multiadhesive Lig proteins in *Leptospira interrogans*. *J Bacteriol* 195:5092–5101. <https://doi.org/10.1128/JB.00663-13>.
 21. Aviat F, Slamti L, Cerqueira GM, Lourdault K, Picardeau M. 2010. Expanding the genetic toolbox for *Leptospira* species by generation of fluorescent bacteria. *Appl Environ Microbiol* 76:8135–8142. <https://doi.org/10.1128/AEM.02199-10>.
 22. Ozuru R, Saito M, Kanemaru T, Miyahara S, Villanueva SY, Murray GL, Adler B, Fujii J, Yoshida SI. 2017. Adipose tissue is the first colonization site of *Leptospira interrogans* in subcutaneously infected hamsters. *PLoS One* 12:e0172973. <https://doi.org/10.1371/journal.pone.0172973>.
 23. Ratet G, Veyrier FJ, Fanton d'Andon M, Kammerscheit X, Nicola MA, Picardeau M, Boneca IG, Werts C. 2014. Live imaging of bioluminescent *Leptospira interrogans* in mice reveals renal colonization as a stealth escape from the blood defenses and antibiotics. *PLoS Negl Trop Dis* 8:e3359. <https://doi.org/10.1371/journal.pntd.0003359>.
 24. Pappas CJ, Benaroudj N, Picardeau M. 2015. A replicative plasmid vector allows efficient complementation of pathogenic *Leptospira* strains. *Appl Environ Microbiol* 81:3176–3181. <https://doi.org/10.1128/AEM.00173-15>.
 25. Zhu W, Wang J, Zhu Y, Tang B, Zhang Y, He P, Zhang Y, Liu B, Guo X, Zhao G, Qin J. 2015. Identification of three extra-chromosomal replicons in *Leptospira* pathogenic strain and development of new shuttle vectors. *BMC Genomics* 16:90. <https://doi.org/10.1186/s12864-015-1321-y>.
 26. Chen D, Patton JT. 2001. Reverse transcriptase adds nontemplated nucleotides to cDNAs during 5'-RACE and primer extension. *Biotechniques* 30:574–582. <https://doi.org/10.2144/01303rr02>.
 27. Miller WG, Lindow SE. 1997. An improved GFP cloning cassette designed for prokaryotic transcriptional fusions. *Gene* 191:149–153. [https://doi.org/10.1016/S0378-1119\(97\)00051-6](https://doi.org/10.1016/S0378-1119(97)00051-6).
 28. Fontana C, Lambert A, Benaroudj N, Gasparini D, Gorgette O, Cachet N, Bomchil N, Picardeau M. 2016. Analysis of a spontaneous non-motile and avirulent mutant shows that FlIM is required for full endoflagella assembly in *Leptospira interrogans*. *PLoS One* 11:e0152916. <https://doi.org/10.1371/journal.pone.0152916>.
 29. Zhukova A, Fernandes LG, Hugon P, Pappas CJ, Sismeiro O, Coppee JY, Becavin C, Malabat C, Eshghi A, Zhang JJ, Yang FX, Picardeau M. 2017. Genome-wide transcriptional start site mapping and sRNA identification in the pathogen *Leptospira interrogans*. *Front Cell Infect Microbiol* 7:10. <https://doi.org/10.3389/fcimb.2017.00010>.
 30. de Faria MT, Calderwood MS, Athanasio DA, McBride AJ, Hartskeerl RA, Pereira MM, Ko AI, Reis MG. 2008. Carriage of *Leptospira interrogans* among domestic rats from an urban setting highly endemic for leptospirosis in Brazil. *Acta Trop* 108:1–5. <https://doi.org/10.1016/j.actatropica.2008.07.005>.
 31. Villanueva SY, Ezoe H, Baterna RA, Yanagihara Y, Muto M, Koizumi N, Fukui T, Okamoto Y, Masuzawa T, Cavinta LL, Gloriani NG, Yoshida S. 2010. Serologic and molecular studies of *Leptospira* and leptospirosis among rats in the Philippines. *Am J Trop Med Hyg* 82:889–898. <https://doi.org/10.4269/ajtmh.2010.09-0711>.
 32. Hu W, Lin X, Yan J. 2014. *Leptospira* and leptospirosis in China. *Curr Opin Infect Dis* 27:432–436. <https://doi.org/10.1097/QCO.000000000000097>.
 33. Fonseca LS, da Silva JB, Milanez JS, Monteiro-Vitorello CB, Momo L, de Morais ZM, Vasconcelos SA, Marques MV, Ho PL, da Costa RM. 2013. *Leptospira interrogans* serovar Copenhageni harbors two *lexA* genes involved in SOS response. *PLoS One* 8:e76419. <https://doi.org/10.1371/journal.pone.0076419>.
 34. Cuñé J, Cullen P, Mazon G, Campoy S, Adler B, Barbe J. 2005. The *Leptospira interrogans lexA* gene is not autoregulated. *J Bacteriol* 187:5841–5845. <https://doi.org/10.1128/JB.187.16.5841-5845.2005>.
 35. Morero NR, Botti H, Nitta KR, Carrion F, Obal G, Picardeau M, Buschiazzo A. 2014. HemR is an OmpR/PhoB-like response regulator from *Leptospira*, which simultaneously effects transcriptional activation and repression of key haem metabolism genes. *Mol Microbiol* 94:340–352. <https://doi.org/10.1111/mmi.12763>.
 36. Hu WL, Pappas CJ, Zhang JJ, Yang YY, Yan J, Picardeau M, Yang XF. 2017. The EbpA-RpoN regulatory pathway of the pathogen *Leptospira interrogans* is essential for survival in the environment. *Appl Environ Microbiol* 83:e02377-16. <https://doi.org/10.1128/AEM.02377-16>.
 37. Ren SX, Fu G, Jiang XG, Zeng R, Miao YG, Xu H, Zhang YX, Xiong H, Lu G, Lu LF, Jiang HQ, Jia J, Tu YF, Jiang JX, Gu WY, Zhang YQ, Cai Z, Sheng HH, Yin HF, Zhang Y, Zhu GF, Wan M, Huang HL, Qian Z, Wang SY, Ma W, Yao ZJ, Shen Y, Qiang BQ, Xia QC, Guo XK, Danchin A, Saint Girons I, Somerville RL, Wen YM, Shi MH, Chen Z, Xu JG, Zhao GP. 2003. Unique physiological and pathogenic features of *Leptospira interrogans* revealed by whole-genome sequencing. *Nature* 422:888–893. <https://doi.org/10.1038/nature01597>.
 38. Ko AI, Galvao Reis M, Ribeiro Dourado CM, Johnson WD, Jr, Riley LW, Salvador Leptospirosis Study Group. 1999. Urban epidemic of severe leptospirosis in Brazil. *Lancet* 354:820–825. [https://doi.org/10.1016/S0140-6736\(99\)80012-9](https://doi.org/10.1016/S0140-6736(99)80012-9).
 39. Demarre G, Guerout AM, Matsumoto-Mashimo C, Rowe-Magnus DA, Marliere P, Mazel D. 2005. A new family of mobilizable suicide plasmids based on broad host range R388 plasmid (IncW) and RP4 plasmid

- (IncPα) conjugative machineries and their cognate *Escherichia coli* host strains. Res Microbiol 156:245–255. <https://doi.org/10.1016/j.resmic.2004.09.007>.
40. Matsunaga J, Medeiros MA, Sanchez Y, Werneid KF, Ko AI. 2007. Osmotic regulation of expression of two extracellular matrix-binding proteins and a haemolysin of *Leptospira interrogans*: differential effects on LigA and Sph2 extracellular release. Microbiology 153:3390–3398. <https://doi.org/10.1099/mic.0.2007/007948-0>.
 41. Shang ES, Summers TA, Haake DA. 1996. Molecular cloning and sequence analysis of the gene encoding LipL41, a surface-exposed lipoprotein of pathogenic *Leptospira* species. Infect Immun 64:2322–2330.
 42. Zheng L, Baumann U, Reymond JL. 2004. An efficient one-step site-directed and site-saturation mutagenesis protocol. Nucleic Acids Res 32:e115. <https://doi.org/10.1093/nar/gnh110>.
 43. Slamti L, Picardeau M. 2012. Construction of a library of random mutants in the spirochete *Leptospira biflexa* using a mariner transposon. Methods Mol Biol 859:169–176. https://doi.org/10.1007/978-1-61779-603-6_9.
 44. Lassmann T, Sonnhammer EL. 2005. Kalign—an accurate and fast multiple sequence alignment algorithm. BMC Bioinformatics 6:298. <https://doi.org/10.1186/1471-2105-6-298>.
 45. R Core Team. 2014. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.