

UC Riverside

UC Riverside Previously Published Works

Title

Draft Genome Sequences of the Antarctic Endolithic Fungi *Rachicladosporium antarcticum* CCFEE 5527 and *Rachicladosporium* sp. CCFEE 5018

Permalink

<https://escholarship.org/uc/item/8bv632cn>

Journal

Microbiology Resource Announcements, 5(27)

ISSN

2576-098X

Authors

Coleine, Claudia
Masonjones, Sawyer
Selbmann, Laura
et al.

Publication Date

2017-07-13

DOI

10.1128/genomea.00397-17

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial-ShareAlike License, available at <https://creativecommons.org/licenses/by-nc-sa/4.0/>

Peer reviewed



Draft Genome Sequences of the Antarctic Endolithic Fungi *Rachicladosporium antarcticum* CCFEE 5527 and *Rachicladosporium* sp. CCFEE 5018

Claudia Coleine,^{a,b} Sawyer Masonjones,^b Laura Selbmann,^a Laura Zucconi,^a Silvano Onofri,^a Claudia Pacelli,^a  Jason E. Stajich^b

Department of Ecological and Biological Sciences, University of Tuscia, Viterbo, Italy^a; Department of Plant Pathology and Microbiology and Institute of Integrative Genome Biology, University of California, Riverside, Riverside, California, USA^b

ABSTRACT The draft genome sequences of *Rachicladosporium antarcticum* CCFEE 5527 and *Rachicladosporium* sp. CCFEE 5018 are the first sequenced genomes from this genus, which comprises rock-inhabiting fungi. These endolithic strains were isolated from inside rocks collected from the Antarctic Peninsula and Battleship Promontory (McMurdo Dry Valleys), Antarctica, respectively.

The past decade has revealed an unexpected fungal diversity associated with rocks, which serves as a primary substrate colonized by microorganisms in extreme dry and cold or hot environments. Under these harsh conditions, active growth is rare on exposed surfaces, and endolithism is a necessary ecological adaptation for survival (1). Black meristematic fungi are a morpho-ecological group of ascomycetes with a peculiar tendency to the extremes and are characterized by melanin pigmentation. They are typical and abundant members of Antarctic cryptoendolithic communities (2). These fungi are equally named black yeasts or microcolonial fungi and rock inhabitant fungi when found growing within rocks (3–7). We produced draft genome sequences of the Antarctic fungi *Rachicladosporium antarcticum* CCFEE 5527 Onofri & Egidi (2) and *Rachicladosporium* sp. strain CCFEE 5018 to provide genome resources to study fungal adaptation to extreme environments and endolithic lifestyles. These genomic resources may give clues for studying the evolution of extremophiles and stress adaptation in these enigmatic fungi.

Rachicladosporium strains were obtained from the Culture Collection of Fungi from Extreme Environments (Viterbo, Italy) and were cultured from inside Antarctic rocks. Species designation of *Rachicladosporium* sp. strain CCFEE 5018 is still being determined, and the internal transcribed spacer sequence is 98.6% identical to *Rachicladosporium monterosium* strain CBS 137178 Isola & Zucconi (2). Cultures were grown on 2% malt extract agar. DNA was extracted using a cetyltrimethylammonium bromide (CTAB)-based protocol (8). Two phenol-chloroform purification steps were used to eliminate melanin from the DNA. Total genomic DNA was sheared with a Covaris S220 ultrasonicator. Sequencing libraries were constructed using NeoPrep TruSeq Nano DNA sample prep (Illumina, Inc., San Diego, CA). Libraries were normalized, pooled, and sequenced on an Illumina MiSeq with 2 × 300 paired-end reads. *R. antarcticum* was sequenced to a depth of 41× and *Rachicladosporium* sp. CCFEE 5018 to 175× to improve assembly quality.

Genome assembly with MaSuRCA version 2.3.2 (9) was followed by vector sequence filtering with Sequin (<https://www.ncbi.nlm.nih.gov/Sequin/>). Redundant contigs which

Received 3 April 2017 Accepted 8 May 2017 Published 6 July 2017

Citation Coleine C, Masonjones S, Selbmann L, Zucconi L, Onofri S, Pacelli C, Stajich JE. 2017. Draft genome sequences of the Antarctic endolithic fungi *Rachicladosporium antarcticum* CCFEE 5527 and *Rachicladosporium* sp. CCFEE 5018. *Genome Announc* 5:e00397-17. <https://doi.org/10.1128/genomeA.00397-17>.

Copyright © 2017 Coleine et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Jason E. Stajich, jason.stajich@ucr.edu.

aligned by MUMmer (10) at 95% across their entire length were removed. The *R. antarcticum* genome assembly was 47.4 Mb (number of contigs, 267; N_{50} , 896 kb; L_{50} , 20). The initial *Rachicladosporium* sp. CCFEE 5018 assembly was fragmented (2,099 contigs) but was scaffolded by synteny to *R. antarcticum* with Satsuma2 (11) and Mercator (12) into 233 scaffolds (N_{50} , 1.35 Mb; L_{50} , 12). The genomes were annotated with Funannotate utilizing Augustus (13), GeneMark.hmm-ES (14), and EVM (15) and prepared for GenBank with Genome Annotation Generator (16). Gene function predictions were assigned by matches to the Pfam (17), MEROPS (18), CAZy (19), InterProScan (20), and Swiss-Prot databases (21). Product descriptions were transferred from homologs with 60% similar alignments across 60% of the protein length. A total of 18,781 protein-coding genes were predicted in *R. antarcticum* and 18,892 in *Rachicladosporium* sp. CCFEE 5018.

Accession number(s). These whole-genome shotgun projects have been deposited at DDBJ/ENA/GenBank under the accession numbers [NAJO00000000](https://doi.org/10.1093/nar/njz000) and [NAEU00000000](https://doi.org/10.1093/nar/naeu00000000). The versions described in this paper are the first versions, NAJO01000000 and NAEU01000000.

ACKNOWLEDGMENTS

We thank the Italian National Program for Antarctic Researches for funding sampling campaigns and research in the frame of project 2013/AZ-17. The Italian National Museum of Antarctica (MNA) is acknowledged for financial support to the Culture Collection of Fungi from Extreme Environments (CCFEE), the mycological section of the MNA, and for kindly providing the strains. This work was partially supported by the USDA Agriculture Experimental Station at the University of California, Riverside and NIFA Hatch project CA-R-PPA-5062-H to J.E.S., and high-performance computing resources in the Institute for Integrative Genome Biology (IIGB) at UC Riverside supported by grants NSF DBI-1429826 and NIH S10-OD016290.

Genome sequencing was performed in the Genomics Core Facility of the IIGB at University of California, Riverside.

REFERENCES

- Zucconi L, Onofri S, Cecchini C, Isola D, Ripa C, Fenice M, Madonna S, Reboleiro-Rivas P, Selbmann L. 2016. Mapping the lithic colonization at the boundaries of life in northern Victoria Land, Antarctica. *Polar Biol* 39:91–102. <https://doi.org/10.1007/s00300-014-1624-5>.
- Egidi E, de Hoog GS, Isola D, Onofri S, Quaedvlieg W, de Vries M, Verkley GJM, Stielow JB, Zucconi L, Selbmann L. 2014. Phylogeny and taxonomy of meristematic rock-inhabiting black fungi in the Dothideomycetes based on multi-locus phylogenies. *Fungal Divers* 65:127–165. <https://doi.org/10.1007/s13225-013-0277-y>.
- Sterflinger K. 1998. Temperature and NaCl-tolerance of rock-inhabiting meristematic fungi. *Antonie Van Leeuwenhoek* 74:271–281. <https://doi.org/10.1023/A:1001753131034>.
- Ruibal C, Gueidan C, Selbmann L, Gorbushina AA, Crous PW, Groenewald JZ, Muggia L, Grube M, Isola D, Schoch CL, Staley JT, Lutzoni F, de Hoog GS. 2009. Phylogeny of rock-inhabiting fungi related to Dothideomycetes. *Stud Mycol* 64:123–133S7 123–133S7. <https://doi.org/10.3114/sim.2009.64.06>.
- Selbmann L, de Hoog GS, Zucconi L, Isola D, Ruisi S, van den Ende AHGG, Ruibal C, De Leo F, Urzi C, Onofri S. 2008. Drought meets acid: three new genera in a dothidealean clade of extremotolerant fungi. *Stud Mycol* 61:1–20. <https://doi.org/10.3114/sim.2008.61.01>.
- Selbmann L, de Hoog GS, Mazzaglia A, Friedmann EI, Onofri S. 2005. Fungi at the edge of life: cryptendolithic black fungi from Antarctic desert. *Stud Mycol* 51:1–32.
- Selbmann L, de Hoog GS, Zucconi L, Isola D, Onofri S. 2014. Black yeasts in cold habitats, p 173–189. In Buzzini P, Margesin R (ed), *Cold-adapted yeasts*. Springer, Berlin, Germany.
- Fulton TM, Chunwongse J, Tanksley SD. 1995. Microprep protocol for extraction of DNA from tomato and other herbaceous plants. *Plant Mol Biol Rep* 13:207–209. <https://doi.org/10.1007/BF02670897>.
- Zimin AV, Marçais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. 2013. The MaSuRCA genome assembler. *Bioinformatics* 29:2669–2677. <https://doi.org/10.1093/bioinformatics/btt476>.
- Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004. Versatile and open software for comparing large genomes. *Genome Biol* 5:R12. <https://doi.org/10.1186/gb-2004-5-2-r12>.
- Grabherr MG, Russell P, Meyer M, Mauceli E, Alföldi J, Di Palma F, Lindblad-Toh K. 2010. Genome-wide synteny through highly sensitive sequence alignment: Satsuma. *Bioinformatics* 26:1145–1151. <https://doi.org/10.1093/bioinformatics/btq102>.
- Dewey CN. 2006. Whole-genome alignments and polytopes for comparative genomics. Ph.D. thesis. University of California, Berkeley, Berkeley, CA.
- Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B. 2006. AUGUSTUS: ab initio prediction of alternative transcripts. *Nucleic Acids Res* 34:W435–W439. <https://doi.org/10.1093/nar/gkl200>.
- Ter-Hovhannisyan V, Lomsadze A, Chernoff YO, Borodovsky M. 2008. Gene prediction in novel fungal genomes using an ab initio algorithm with unsupervised training. *Genome Res* 18:1979–1990. <https://doi.org/10.1101/gr.081612.108>.
- Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, White O, Buell CR, Wortman JR. 2008. Automated eukaryotic gene structure annotation using EVIDENCEModeler and the program to assemble spliced alignments. *Genome Biol* 9:R7. <https://doi.org/10.1186/gb-2008-9-1-r7>.
- Hall B, DeRego T, Geib S. 2014. GAG: the genome annotation generator. <https://genomeannotation.github.io/GAG/>.
- Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, Heger A, Hetherington K, Holm L, Mistry J, Sonnhammer ELL, Tate J, Punta M. 2014. Pfam: the protein families database. *Nucleic Acids Res* 42:D222–D230. <https://doi.org/10.1093/nar/gkt1223>.
- Rawlings ND, Barrett AJ, Bateman A. 2014. Using the MEROPS database

- for proteolytic enzymes and their inhibitors and substrates. *Curr Protoc Bioinformatics* 48:1.25.1–1.2533. <https://doi.org/10.1002/0471250953.bi0125s48>.
19. Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B. 2014. The carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Res* 42:D490–D495. <https://doi.org/10.1093/nar/gkt1178>.
 20. Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong SY, Lopez R, Hunter S. 2014. InterProScan 5: Genome-scale protein function classification. *Bioinformatics* 30:1236–1240. <https://doi.org/10.1093/bioinformatics/btu031>.
 21. Boutet E, Lieberherr D, Tognolli M, Schneider M, Bansal P, Bridge AJ, Poux S, Bougueleret L, Xenarios I. 2016. UniProtKB/Swiss-Prot, the manually annotated section of the UniProt KnowledgeBase: how to use the entry view. *Methods Mol Biol* 1374:23–54. https://doi.org/10.1007/978-1-4939-3167-5_2.