

UC San Diego

UC San Diego Previously Published Works

Title

Microbiota Modulates Cardiac Transcriptional Responses to Intermittent Hypoxia and Hypercapnia

Permalink

<https://escholarship.org/uc/item/8n69k5kt>

Authors

Zhou, Dan
Xue, Jin
Miyamoto, Yukiko
et al.

Publication Date

2021

DOI

10.3389/fphys.2021.680275

Peer reviewed



Microbiota Modulates Cardiac Transcriptional Responses to Intermittent Hypoxia and Hypercapnia

Dan Zhou^{††}, Jin Xue^{††}, Yukiko Miyamoto², Orit Poulsen¹, Lars Eckmann² and Gabriel G. Haddad^{1,3,4*}

OPEN ACCESS

Edited by:

Rodrigo Iturriaga,
Pontificia Universidad Católica
de Chile, Chile

Reviewed by:

Lena Lavie,
Technion – Israel Institute
of Technology, Israel
Bin Liu,
University of Texas MD Anderson
Cancer Center, United States

*Correspondence:

Gabriel G. Haddad
ghaddad@health.ucsd.edu

^{††} These authors have contributed
equally to this work and share first
authorship

Specialty section:

This article was submitted to
Integrative Physiology,
a section of the journal
Frontiers in Physiology

Received: 13 March 2021

Accepted: 20 May 2021

Published: 25 June 2021

Citation:

Zhou D, Xue J, Miyamoto Y,
Poulsen O, Eckmann L and
Haddad GG (2021) Microbiota
Modulates Cardiac Transcriptional
Responses to Intermittent Hypoxia
and Hypercapnia.
Front. Physiol. 12:680275.
doi: 10.3389/fphys.2021.680275

¹ Division of Respiratory Medicine, Department of Pediatrics, University of California, San Diego, La Jolla, CA, United States, ² Department of Medicine, University of California, San Diego, La Jolla, CA, United States, ³ Department of Neurosciences, University of California, San Diego, La Jolla, CA, United States, ⁴ Rady Children's Hospital-San Diego, San Diego, CA, United States

The microbiota plays a critical role in regulating organismal health and response to environmental stresses. Intermittent hypoxia and hypercapnia, a condition that represents the main hallmark of obstructive sleep apnea in humans, is known to induce significant alterations in the gut microbiome and metabolism, and promotes the progression of atherosclerosis in mouse models. To further understand the role of the microbiome in the cardiovascular response to intermittent hypoxia and hypercapnia, we developed a new rodent cage system that allows exposure of mice to controlled levels of O₂ and CO₂ under gnotobiotic conditions. Using this experimental setup, we determined the impact of the microbiome on the transcriptional response to intermittent hypoxia and hypercapnia in the left ventricle of the mouse heart. We identified significant changes in gene expression in both conventionally reared and germ-free mice. Following intermittent hypoxia and hypercapnia exposure, we detected 192 significant changes in conventionally reared mice (96 upregulated and 96 downregulated) and 161 significant changes (70 upregulated and 91 downregulated) in germ-free mice. Only 19 of these differentially expressed transcripts (~10%) were common to conventionally reared and germ-free mice. Such distinct transcriptional responses imply that the host microbiota plays an important role in regulating the host transcriptional response to intermittent hypoxia and hypercapnia in the mouse heart.

Keywords: gnotobiotics, transcriptome, intermittent hypoxia and hypercapnia, heart, mice

Abbreviations: OSA, obstructive sleep apnea; IHH, intermittent hypoxia and hypercapnia; GF, germ-free; CONV, conventional; LV, left ventricle; HEPA, high-efficiency particulate air; DAVID, Database for Annotation, Visualization, and Integrated Discovery; IPA, Ingenuity Pathway Analysis; KEGG, Kyoto Encyclopedia of Genes and Genomes.

INTRODUCTION

Obstructive sleep apnea (OSA) is the most common type of sleep disorder in humans, and OSA is very prevalent in adults and children. It is characterized by repeated episodes of complete or partial obstructions of the upper airways during sleep, which is associated with a reduction in blood oxygen saturation and an increase in blood CO₂. It has been demonstrated by experimental and clinical studies that OSA may lead to a number of health consequences including cardiovascular diseases ranging from cardiometabolic disorders and arrhythmogenesis to heart failure (Al Lawati et al., 2009; Baguet et al., 2012; Levy et al., 2013; Drager et al., 2015; Lyons and Bradley, 2015; Hoyos et al., 2017; May et al., 2017).

In addition, evidence from both animal and human studies supports the notion that the gut microbiota plays a critical role in the onset and progression of various cardiovascular diseases (Karbach et al., 2016; Koch et al., 2017; Weis, 2018). Previous studies by others and us have shown that intermittent hypoxia with/without hypercapnia alters the gut microbiome (Moreno-Indias et al., 2015, 2016; Lucking et al., 2018; Tripathi et al., 2018, 2019). In turn, such alterations may induce developmental defects and functional dysregulations. For example, the human oral and intestinal microbiota has been recognized as a significant determinant of cardiovascular disease risk (Ordovas and Mooser, 2006), and suppression of the gut microbiome ameliorates age-dependent arterial dysfunction (Brunt et al., 2019). Recent studies also revealed a potential contribution of gut microbes to certain human cardiometabolic diseases (Vinje et al., 2014) and heart failure (Sandek et al., 2007; Nagatomo and Tang, 2015; Nemet et al., 2020).

Besides impacting cellular signaling, the microbiome has been demonstrated to modify the epigenetic landscape of the host genome, which may influence cellular differentiation and proliferation, leading to new phenotypic traits throughout the age spectrum (Takahashi et al., 2011; Pevsner-Fischer et al., 2017; Qin and Wade, 2018). For example, several studies have demonstrated a direct link between the gut microbiome and host epigenetic states in the context of inflammatory bowel disease, obesity, cancer, and behavioral abnormalities (Kellermayer et al., 2010; Manichanh et al., 2012; Dejea et al., 2014; Paul et al., 2015; Farhana et al., 2018; Sfanos et al., 2018; Chu et al., 2019).

Since gut microbiota plays an important role in regulating host gene expression (Patrignani et al., 2014; Chu et al., 2019; Richards et al., 2019; Weger et al., 2019), in the current study, we tested the hypothesis that the microbiome regulates the transcriptional response to intermittent hypoxia and hypercapnia (IHH) exposure in the heart. Using a newly developed experimental setup to impose IHH under gnotobiotic conditions, we performed 150-bp paired-end RNA-seq and utilized multiple bioinformatic approaches to analyze the transcriptional alterations following IHH treatment in the left ventricle (LV) of conventionally reared (CONV) and germ-free (GF) C57BL/6 mice. In addition, we compared our results with those reported for other organs in male

GF mice of similar genetic background and age (Weger et al., 2019) (dataset GSE77221). We found significant and distinct differences in transcriptional responses between CONV and GF mice, demonstrating that, indeed, the microbiome plays an important role in regulating IHH response in the heart.

MATERIALS AND METHODS

Experimental Animals


GF and CONV C57BL/6 mice that were 8–12 weeks old were used for this experiment. GF status was confirmed weekly by standard microbiological cultures of fecal extracts, using thioglycollate broth and potato dextrose broth for facultative anaerobes including fungi, and cooked meat broth, heart infusion broth, and Columbia blood agar for obligate anaerobes (Moody et al., 2019). All animal protocols were approved by the Animal Care Committee of the University of California, San Diego and followed the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health.

IHH Treatment

All mice were housed in specialized cages for the 2-week treatments (IHH or normoxia; GF or CONV) (**Figure 1**). These specialized cages were developed by us in the Gnotobiotic Animal Facility at the University of California, San Diego. Each sealed single cage (Sealed Positive Pressure System, Allentown, NJ, United States) was ventilated by a single, HEPA (high-efficiency particulate air)-filtered gas supply pump. The cages were opened and closed for stocking and servicing inside a Class IIa laminar-flow biosafety cabinet. The system allows complete control of gas inflow and outflow, as has been confirmed by measurements of exhaust gases using an O₂/CO₂ analyzer (Quantek Instruments, Grafton, MA, United States). The IHH condition was maintained in a computer-controlled atmosphere chamber system (OxyCycler, BioSpherix, Redfield, NY, United States) as previously described (Fan et al., 2005; Zhou et al., 2008). IHH exposure was introduced to the mice in short periods of ~4 min of synchronized reduction in (O₂) (from 21 to 8%) and increase in (CO₂) (from ~0.5 to 8%) separated by alternating ~4-min periods of normoxic room air [(O₂) = 21% and (CO₂) ≈ 0.5%] with 2 ramp intervals (~1 min each) for a total of 10 IHH events per hour for 10 h per day during the light cycle. Controls received a flow of room air throughout the entire experimental period. Prior works have shown that the oxygen saturation level (SaO₂) ranged from 60 to 80% when mice were subjected to 5–10% hypoxic treatments (Sforza and Roche, 2016; Xue et al., 2017).

RNA Sequencing and Data Analysis

At the end of the IHH exposure, mice were sacrificed, and the heart was harvested and separated into the right and left ventricle under sterile conditions and then placed into separate tubes containing RNeasy Lysis Solution (Thermo Fisher Scientific, Ashville, NC, United States). RNA from the LV was extracted using the RNeasy Mini Kit



(QIAGEN, Hilden, Germany) and sent to Novogene (Chula Vista, CA, United States) for QC and sequencing ($n = 3/\text{group}$). A 150 nt paired-end sequencing on the Illumina HiSeq 4000 platform (Illumina, San Diego, CA, United States) was performed. Rigorous quality controls of the paired-end reads were assessed using FastQC tools, and the adapter sequences and low-quality bases were trimmed using Cutadapt (Didion et al., 2017). The resulting reads were processed using the Illumina BaseSpace software package using the STAR aligner (Dobin et al., 2013) to align reads against the *Mus musculus* UCSC mm9 reference genome. Differential expression was calculated using DESeq 2 with default settings (Anders and Huber, 2010), and differentially expressed genes (DEGs) were identified with a threshold of $|\text{Fold-Change}| > 1.5$ and adjusted p -value < 0.05 . Bioinformatic data mining of the differentially expressed transcripts was carried out with the tools provided by the GWAS Catalog¹ (Buniello et al., 2019), the Database for Annotation, Visualization, and Integrated Discovery (DAVID)² (Dennis et al., 2003; Huang et al., 2007), the Ingenuity Pathway Analysis (IPA) package (QIAGEN, Redwood City, CA, United States), as well as KEGG³ (Ogata et al.,

1999; Kanehisa and Goto, 2000) and the Reactome⁴ (Fabregat et al., 2017). Multiple testing correction was applied, and the FDR q -values were estimated to correct the p -values for the multiple testing issue.

Data Availability

The paired-end RNA-seq data of $n = 12$ samples are available at <https://www.ncbi.nlm.nih.gov/bioproject/692551> (BioProject ID: PRJNA692551).

RESULTS AND DISCUSSION

Experimental Design


The microbiome, particularly in the gut, plays an important role in maintaining systemic homeostasis and contributes to a number of human diseases [for selected reviews, see Lynch and Pedersen (2016), Gilbert et al. (2018), Dominguez-Bello et al. (2019), Harikrishnan (2019), Helmink et al. (2019), Team (2019)]. However, its potential involvement in regulating many physiologic and pathophysiologic responses under stressful conditions remains to be explored. In the current study, we used a prospective case-control study design to determine the

¹<https://www.ebi.ac.uk/gwas/>

²<https://david.ncifcrf.gov/>

³<https://www.genome.jp/kegg/>


⁴<https://reactome.org/>



impact of the microbiome on cardiac transcriptional responses to IHH in CONV and GF mice (Figure 1A). To maintain GF animals under controlled gas conditions, we developed a custom cage system with minimal dead space and programmable HEPA-filtered gas supply (Figure 1B). Analysis of the exhaust gases of the cage system revealed rapid (<20 s) equilibration of the cage environment to match the input gas mixture (Figure 1C). GF mice could be kept in the system for extended periods (>2 weeks) without compromising their GF state, demonstrating that the newly developed system was well suited to conduct short- and long-term experiments of constant

or intermittent hypoxia with or without hypercapnia under gnotobiotic conditions.

GF and CONV mice of the same strain, sex, and age (kept in the same cage system as the GF mice) were either exposed to the IHH condition for 2 weeks to mimic sleep apnea in humans (Kanaan et al., 2006; Xue et al., 2017; Hunyor and Cook, 2018) or kept under normoxia/normocapnia throughout as controls. For a broad characterization of the cardiac gene expression responses, we extracted total RNA from the LV and applied paired-end RNA-seq as a genome-wide discovery approach. Approximately 38–50 million total reads were obtained per sample with over 94%



of the reads uniquely mapped to the mouse reference genome (UCSC mm9) (**Supplementary Table 1**).

Microbiota Impacts Cardiac Gene Expression Under Normoxic Conditions

We first determined the impact of the microbiota on cardiac gene expression under normoxic and normocapnic conditions. A total of 117 DEGs were identified between CONV and GF mice. Of these, 73 were upregulated and 44 were downregulated (Fold change > 1.5 and adjusted p -value < 0.05) in the left

ventricle of GF mice as compared to CONV controls (**Figure 2A** and **Supplementary Table 2**). The most upregulated transcripts included *Amd1*, *Zfp950*, *Rnf125*, and *Khdc4*, which are involved in polyamine biosynthesis, transcription regulation, ubiquitination, and subsequent proteasomal degradation of target proteins, and pre-mRNA splicing, respectively. In contrast, the most downregulated transcripts included genes encoding a transcription repressor (*Zmynd11*), a Calcium Voltage-Gated Channel Subunit (*Cacna1d*), and a serine/threonine kinase (*Limk1*) (**Figure 2B**). In addition, several functional implications were suggested by data mining and bioinformatic analyses

using the DAVID and IPA tools. Results derived from DAVID functional annotation suggested alterations in transcription regulation, alpha-actinin binding, and sarcolemma organization, as well as pathways that regulate cardiac function and activity, including the GnRH pathway, the MAPK signaling pathway, the Oxytocin signaling pathway, and vascular muscle contraction (**Supplementary Table 3A**). In addition, Ingenuity analysis showed potential changes in several canonical pathways that are important in cardiac function, such as the mechanisms regulating synaptic long-term potentiation, Myc-mediated apoptosis signaling, and the role of NFAT in cardiac hypertrophy (**Figure 3** and **Supplementary Table 3B**). The different alterations on the “Myc Mediated Apoptosis Signaling” and the “Role of NFAT in Cardiac Hypertrophy” suggested that cell death and proliferation signaling pathways were differently configured in the GF mice (**Supplementary Table 3B**). Since (1) Myc-mediated apoptosis plays a critical role in response to stress in the myocardium (Hou et al., 2016; Veeraveedu et al., 2017; Guan et al., 2019), (2) NFAT plays a major role in cardiac hypertrophy [for selected reviews see Clerk et al. (2007), Rohini et al. (2010), Nakayama et al. (2013)], and (3) previous studies also shown that gut microbiome regulates the activity of Myc in mouse liver (Kolodziejczyk et al., 2020) and NFAT in colorectal cancer in humans (Peucker et al., 2016), our current findings suggest that alterations of Myc and NFAT signaling in GF mice may modify cardiac activity and stress response.

Furthermore, we found that 36% of the DEGs (42 out of 117) in GF mice encode nuclear proteins, including transcription factors (e.g., *Dach1* and *Gtf2ird2*), protein kinases (e.g., *Clk1* and *Limk1*), DNA, and histone modification enzymes (e.g., *Dnmt3b* and *Kmt5b*). Such a significant enrichment of nuclear proteins suggested differences of transcriptional regulation mechanisms between GF and CONV mice, which may lead to distinct transcriptional responses to stress in the heart.

Distinct Transcriptional Alterations in Different Organs of GF Mice

Alterations in gene expression in GF mice have been reported in several organs (Weger et al., 2019; Mills et al., 2020). In order to determine common and distinct transcriptional responses between different organs in GF mice, we compared the reported changes (GSE77221, GEO, NCBI) with those in our study. As shown in **Figure 4**, each organ displayed a distinct transcriptional profile. Among the total 5,109 significantly altered transcripts ($|\text{Fold Change}| > 1.5$ and $\text{FDR} < 0.05$) in the heart, liver, white adipose tissue, duodenum, and ileum of GF mice, 3,386 (66%) were tissue specific. The largest number of unique changes was observed in the ileum (1,144 genes), followed by duodenum (1,096 genes), liver (587 genes), white fat (467 genes), and heart (92 genes), possibly reflecting the relative vicinity of the respective organs to the abundant gut microbiota. No genes were identified that were altered in all five organs, indicating a lack of common tissue responses to the microbiota. Although the vast majority of cardiac genes with significant expression changes were unique to the heart under GF conditions (79%),




FIGURE 4 | Distinct transcriptional alterations in different organs of GF mice in normoxia. Little common changes were identified in the differentially expressed genes (DEGs) between the examined organs of the GF mice under normoxic condition, except a significant overlapping between the two gut segments (duodenum and ileum), demonstrating a distinct and organ-specific impact of microbiome on gene expression.

the most overlap was seen with the duodenum (5 genes, 4.3% of total DEGs in the heart) and the least with duodenum (or ileum) and white fat (1 gene each, <1% of total DEGs in the heart). Interestingly, the gene *cytochrome b-245, beta chain* (*Cybb*) was commonly altered in heart and gut, suggesting that the lack of a microbiome alters the activity of immune cells, especially neutrophils, which may lead to immune dysregulation in these organs.

GF Condition Alters Transcriptional Responses to IHH in the Heart

Previous studies have shown that stress can alter the composition of the gut microbiome, and in turn, such changes in gut microbiota may regulate stress response in various organs and tissues, including the central nervous system, the digestive system, and the immune system (Lyte et al., 2020; Park et al., 2020; Rengarajan et al., 2020; Tian et al., 2020; Vuong et al., 2020; Wang et al., 2020; Xu et al., 2020). The current study provides evidence that the microbiome may also affect stress responses in the heart through regulating transcription activities. Indeed, following 2 weeks of IHH exposure, we detected 192 significant changes in CONV mice (96 upregulated and 96 downregulated; $|\text{Fold Change}| > 1.5$ and $\text{FDR} < 0.05$) (**Figure 5A** and **Supplementary Table 4**) and 161 significant changes (70 upregulated and 91 downregulated; $|\text{Fold Change}| > 1.5$ and $\text{FDR} < 0.05$) in GF mice (**Figure 5B** and **Supplementary Table 5**). As shown in **Table 1**, the top DEGs in CONV mice include *Rnf44*, *Cyb561a3*, *Cep250*,





TABLE 1 | Top transcriptional alterations in CONV and GF mice following IHH exposure.

RefSeq_ID	Symbol	Normalized read counts						Fold change	Padj
		Air-1	Air-2	Air-3	IHH-1	IHH-2	IHH-3		
CONV Mice									
NR_027395	Rnf44	272.4	260.7	211.7	0.0	0.0	0.0	-1406.2	6.5E-15
NM_201351	Cyb561a3	158.4	127.1	150.7	0.0	0.0	0.0	-827.7	1.8E-12
NM_008383	Cep250	141.4	111.9	138.3	0.0	0.0	0.0	-740.1	6.2E-12
NM_001286664	Ssbp1	93.6	88.9	82.6	0.0	0.0	0.0	-509.8	2.8E-10
NM_001199248	Kat5	123.6	107.7	17.6	0.0	0.0	0.0	-471.9	8.8E-07
NM_001163009	Tnik	62.4	73.3	97.8	0.0	0.0	0.0	-443.1	2.6E-09
NM_001204277	Hdac7	81.2	89.8	54.1	0.0	0.0	0.0	-422.7	5.2E-09
NR_130970_1	Gm14296	71.3	48.6	79.5	0.0	0.0	0.0	-381.8	1.3E-08
NM_008488	Arhgef1	27.4	81.0	85.5	0.0	0.0	0.0	-365.4	3.4E-07
NM_008642	Mttp	61.8	46.2	74.7	0.0	0.0	0.0	-349.2	3.0E-08
NM_001347201	Gucd1	0.0	0.0	0.0	14.4	28.5	43.9	150.7	2.3E-04
NM_001012363	Slc2a9	0.0	0.0	0.0	28.6	39.1	24.6	159.8	2.7E-05
NM_001358355	Zfx	0.0	0.0	0.0	34.6	50.0	27.8	191.7	7.3E-06
NM_178722	Zfp438	0.0	0.0	0.0	38.2	42.6	55.3	234.8	6.2E-07
NM_001167886	Kmt5b	0.0	0.0	0.0	48.3	70.0	21.8	241.4	7.3E-06
NM_001284506	Plxnb2	0.0	0.0	0.0	40.1	70.8	40.4	263.5	5.6E-07
NM_001270495	Tmem254a	0.0	0.0	0.0	51.6	54.4	48.3	266.2	1.3E-07
NM_001291122	Epb411l	0.0	0.0	0.0	100.8	47.7	33.0	314.3	8.4E-07
NR_130970	Gm14296	0.0	0.0	0.0	67.2	61.0	85.7	367.4	8.8E-09
NM_001290367	Asph	0.0	0.0	0.0	168.1	41.5	83.9	502.3	3.9E-08
GF Mice									
NM_001310725	Kmt5a	332.6	616.9	544.7	0.0	0.0	0.0	-2582.6	1.9E-17
NM_009665	Amd1	321.6	421.3	295.8	0.0	0.0	0.0	-1800.0	3.6E-16
NM_001130180	Tnnt2	0.0	69299.9	70470.5	61.6	13.2	12.4	-1602.1	2.1E-03
NM_134125	Trip10	92.3	146.7	489.2	0.0	0.0	0.0	-1251.8	1.0E-07
NM_001102438	Acbd5	107.8	132.9	183.7	0.0	0.0	0.0	-734.7	1.1E-11
NM_001360080	Herc2	151.3	143.0	85.8	0.0	0.0	0.0	-657.8	4.6E-11
NM_001329143	Adams10	94.8	120.1	117.7	0.0	0.0	0.0	-571.0	4.8E-11
NM_001361696	Slc38a10	170.4	58.5	88.2	10.3	29.6	12.4	-6.1	9.7E-04
NM_001039087	Rapgef1	767.1	518.2	501.2	156.1	160.6	283.6	-3.0	2.4E-06
NM_001243837	C7	290.4	258.7	163.4	80.0	95.4	69.2	-2.9	3.2E-06
NM_011638	Tfrc	1034.7	887.6	1268.0	2834.2	3037.6	3108.3	2.8	1.7E-28
NM_001304810	Kcnj3	37.4	11.6	29.9	79.4	115.0	66.8	3.3	1.7E-02
NM_001277328	Sox6	57.1	31.0	27.7	163.1	117.7	123.1	3.5	3.5E-05
NM_001357298	Tfrc	11.3	65.4	25.6	136.5	133.8	147.1	4.1	1.3E-02
NM_001313974	Prkar1a	26.0	24.4	23.2	256.7	24.7	170.0	6.1	1.1E-02
NM_001355067	Ctnnd1	0.0	0.0	0.0	145.9	152.4	18.6	600.8	1.7E-07
NM_001199308	Gm14440	0.0	0.0	0.0	133.6	156.4	63.1	670.1	2.0E-10
NM_198300	Cpeb3	0.0	0.0	0.0	185.3	184.8	41.5	785.4	1.5E-09
NM_013747	Golga5	0.0	0.0	0.0	151.5	128.8	200.1	912.2	9.2E-13
NM_001199186	Slc44a2	0.0	0.0	0.0	222.6	265.2	248.5	1400.1	3.3E-15

Asph, Gm14296, and Epb411l, whereas Kmt5a, Amd1, Tnnt2, Slc44a2, Golga5, and Cpeb3 were among the top changes in GF mice. Interestingly, only 19 transcripts were altered in both CONV and GF mice, representing around 10% of the changes in CONV and GF, respectively, demonstrating that IHH treatment induced distinct and specific transcriptional responses between CONV and GF mice. The majority of these common altered transcripts


were downregulated, and the top three downregulated transcripts were Acta1 (Alpha actin 1), Rn45s (45S pre-ribosomal RNA), and Rhobtb1 (Rho-related BTB domain containing 1). On the other hand, Tfrc (transferrin receptor), Rgs2 (regulator of G-protein signaling 2), and Prkar1a (protein kinase, cAMP-dependent regulatory, type I, alpha) were the top three genes that were commonly upregulated in both CONV and GF mice. In contrast, two transcripts, Nadk2



(mitochondrial NAD kinase 2) and Ssbp1 (single-stranded DNA binding protein 1), were downregulated in CONV mice but upregulated in GF mice following IHH exposure (Figure 5C). Such dramatically different transcriptional responses suggested that IHH exposure may lead to different physiological and functional alterations in the heart of CONV and GF mice.

Distinct Alterations in Cell Signaling in Response to IHH Exposure Between CONV and GF Mice

We identified distinct cell signaling mechanisms in the hearts of CONV and GF mice in response to IHH. In the CONV mice, the top significantly altered transcripts



encoding a group of nuclear proteins [66/192 (34%), $FDR < 0.05$] that are involved in “DNA-templated transcription” [GO:0006351, 42/192 (22%), $FDR < 0.05$] and “regulation of DNA-templated transcription” [GO:0006355, 46/192 (24%), $FDR < 0.05$] (**Supplementary Table 6A**). The top altered signaling mechanisms were Myc mediated apoptosis signaling, coronavirus pathogenesis pathway, and IL-9 signaling (**Figure 6A** and **Supplementary Table 6B**). In GF mice, on the other hand, the significantly altered genes were involved in the biological processes that regulate cell cycle (GO:0051726, including some cyclins, e.g., *Cdkn1a*, *Ccnd1*, and *Cables1*, $FDR < 0.05$), biological rhythms ($FDR < 0.05$),

and MHC class II protein complex (GO:0042613, $FDR < 0.05$) (**Supplementary Table 7A**). The top representative pathways were “regulation of actin-based motility by Rho,” “integrin signaling,” “HGF signaling,” and “circadian rhythm signaling” (**Figure 6B** and **Supplementary Table 7B**).

Following IHH exposure, “regulation of DNA-templated transcription” was one of the most significantly altered biological processes represented by 24% of the significantly altered transcripts in CONV mice. These results suggested a fine-tuning of the transcriptome, at least in part, through epigenetic modifications and rebalance of transcription regulators. Indeed, as shown in **Figure 7**, transcripts encoding several epigenetic modifiers were significantly changed in the CONV mice, but not GF mice, following IHH treatment. In contrast, in the heart of the GF mouse, IHH-induced transcriptional alterations suggested an array of functional/activity adjustments, such as the circadian rhythm and cell cycle genes (e.g., *Per2*, *Clock*, *Cdkn1a*, and *Stat3*) that regulate the heart rate (Yamamura et al., 2010) or the metabolism, hypertrophy, and apoptosis of cardiac cells (Bonney et al., 2013; Hauck et al., 2016; Harhous et al., 2019).

Transcriptomic Indications of Potential Cardiac Disorders in GF and CONV Mice Following IHH Treatment

Substantial evidence has been reported that OSA significantly increases the risk of stroke, coronary heart disease, and heart failure in humans (McEvoy et al., 2016; Drager and Lee, 2018; Benjamin et al., 2019). Interestingly, we found that the IHH-induced transcriptional alterations in the GF and CONV mice suggested distinct risks of cardiac disorders, i.e., IHH exposure evoked different sets of well-characterized pathogenetic genes in GF and CONV mice. As shown in **Table 2** and **Figure 8**, in CONV mice, IHH-induced transcriptional alterations mainly affect cardiac necrosis and cell death. In contrast, in the GF mice, the DEGs included several key regulators of cardiac hypertrophy. In the CONV mice, changes of both well-characterized cell

TABLE 2 | Transcriptome indication of potential IHH-induced cardiac disorders in GF and CONV mice.

Categories	Diseases/Function annotation	<i>p</i> -val	Gene transcripts
CONV/IHH			
Cardiac Necrosis/Cell Death	Cell death of heart cells	0.004	Abcc9, Bcl2l11, Cryab, Gsn, Ikbkb, Nr4a1, Ppp1r10, Stat1, Txnip
Cardiac Necrosis/Cell Death	Cell death of cardiomyocytes	0.012	Abcc9, Bcl2l11, Cryab, Gsn, Nr4a1, Ppp1r10, Stat1, Txnip
Heart Failure	Onset of heart failure	0.013	Cryab
Hepatocellular Peroxisome Proliferation	Formation of peroxisomes	0.013	Pex26
Cardiac Necrosis/Cell Death	Apoptosis of heart cells	0.020	Bcl2l11, Cryab, Gsn, Ikbkb, Nr4a1, Ppp1r10, Stat1, Txnip
Cardiac Arterioopathy	Vasospasm of coronary artery	0.050	Abcc9
Cardiac Necrosis/Cell Death	Apoptosis of cardiomyocytes	0.050	Bcl2l11, Cryab, Gsn, Ikbkb, Nr4a1, Ppp1r10, Txnip
GF/IHH			
Cardiac Arrhythmia	Ventricular extrasystole	0.025	Tnnt2
Cardiac Enlargement	Hypertrophy of heart	0.036	Ace, Ankrd1, Aplnr, Cdkn1a, Fhl2, Nos2, Rgs2, Stat3, Tnnt2
Cardiac Dilatation and Enlargement	Idiopathic dilated cardiomyopathy	0.037	Aplnr
Cardiac Arrhythmia	Atrial fibrillation	0.037	Ace

Red font: Upregulation. Blue font: Downregulation.






FIGURE 8 | Different set of cardiovascular diseases-related genes was altered in the heart of CONV and GF mice following IHH exposure. Prediction of clinical pathology endpoints driven by IHH-induced differentially expressed genes (DEGs) suggested a significant alteration of cardiac cell death in CONV (A) and cardiac hypertrophy in GF mice (B) under IHH condition (right-tailed Fisher’s exact test, p -value < 0.05). (C,D) Expressional abundance of the related DEGs in CONV and GF mice, respectively (* [Fold Change] > 1.5, Benjamini–Hochberg adjusted p < 0.05).

death stimulators and inhibitors were detected, which include BCL2-like 1 (Bcl2l1), Crystallin alpha B (Cryab), Gelsolin (Gsn), Inhibitor of nuclear factor kappa B kinase subunit beta (Ikbkb), Nuclear receptor subfamily 4 group A member 1 (Nr4a1), Protein phosphatase 1 regulatory subunit 10 (Ppp1r10), and Signal transducer and activator of transcription 1 (Stat1), as well as the downregulation of Thioredoxin interacting protein (Txnip) [for selected reviews see Wagner and Siddiqui (2009), Li et al. (2012), Ganguly et al. (2014), Wang and Yoshioka (2017), Opferman and Kothari (2018), Herring et al. (2019), Tang et al. (2019)]. In the GF mice, however, most of the disease-related DEGs were cardiac hypertrophy regulators, such as the downregulation of Angiotensin I converting enzyme (Ace), Ankyrin repeat domain 1 (Ankrd1), Apelin receptor (Aplnr), Nitric oxide synthase 2 (Nos2), and Cardiac type Troponin T2 (Tnnt2), as well as the upregulation of Cyclin-dependent kinase inhibitor 1A (Cdkn1a), Four and a half LIM domains 2 (Fhl2), Regulator of G protein signaling 2 (Rgs2), and Signal transducer and activator of transcription 3 (Stat3) [for selected reviews, see Brooks et al. (1998), Drexler (1999), Takeda et al. (2005), Tsang et al. (2010), Haghikia et al. (2011), Liang et al. (2018), Yotti et al. (2019)]. Since several genes in this group are key regulators of hypertension (e.g., Ace, Nos2, and Rgs2) (Li and Forstermann, 2000; Prichard et al., 2001; Tsang et al., 2010), it is reasonable to hypothesize that dysregulation of blood pressure might be an important mechanism regulating IHH-induced hypertrophy in GF mice.

In summary, our study demonstrated that the microbiome regulates the cardiac transcriptome at baseline and in response to IHH stress. A total of 117 DEGs were identified in the heart of GF mice under normoxic condition. These DEGs encode proteins involved in transcriptional regulation, alpha-actinin binding, and sarcolemma organization. Lack of microbiome also altered cardiac transcriptional responses to IHH stress, since we found 192 IHH-induced DEGs in CONV mice and 161 DEGs in GF mice with little overlap, suggesting that IHH exposure leads to different cardiac responses between CONV and GF mice. Indeed, 24% of the DEGs detected in CONV mice are involved in the “regulation of DNA-templated transcription,” while in GF mice, IHH-induced cardiac DEGs are mostly involved in functional or activity adjustments, such as circadian rhythm and cell cycle. Future functional studies will be needed to define the functional implications of the observed DEGs. For example, IHH-induced DEGs suggest distinct risks of cardiac disorders in GF and CONV mice, since disease-related DEGs mainly affected cardiac necrosis and cell death in CONV mice, while regulators of cardiac hypertrophy were mostly observed in GF mice. Additional studies are also warranted to explore the function of specific microbial species and their metabolites on regulating cardiac gene expression under baseline and stress conditions. Gnotobiotic mouse models are uniquely suited to unravel such microbial contributions to specific physiologic responses.

CONCLUSION

Even though it has been shown that OSA or IHH exposure may lead to the development of various types of cardiac disorders

[for selected reviews, see Baguet et al. (2012), Dumitrascu et al. (2013), Levy et al. (2015), Turnbull (2018)], any contribution of the microbiome to this process is still largely uncharacterized. In the present study, we found that the microbiome plays an important role in the transcriptional responses to IHH stress and identified distinct responsive mechanisms in the heart of GF and CONV mice, suggesting that the microbiome modulates the cardiac function during stressful and disease conditions like OSA. Our study provides insight into the unique and common transcriptional response to IHH that is related to microbial colonization in mice. These transcriptional alterations suggest changes in cellular signaling and interaction networks regulating cardiac function, such as cardiac hypertrophy and cell death. Overall, our study is a step toward a better understanding of the role of microbiota in regulating cardiac health and disease.

DATA AVAILABILITY STATEMENT

The datasets presented in this study can be found in online repositories. The names of the repository/repositories and accession number(s) can be found in the article/**Supplementary Material**.

ETHICS STATEMENT

The animal study was reviewed and approved by the Animal Care Committee of the University of California, San Diego.

AUTHOR CONTRIBUTIONS

DZ, JX, LE, and GH conceptualized the study, designed the experiments, and wrote the manuscript. DZ, JX, YM, and OP performed the experiments. DZ and JX performed data curation and analysis. LE and GH acquired funding and facilitated access to resources for the study. GH supervised the study. All authors contributed to the article and approved the submitted version.

FUNDING

This work was partially supported by the NIH grant P30 DK120515 (LE).

ACKNOWLEDGMENTS

We thank Iain Hartley and Travis Smith for technical support.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fphys.2021.680275/full#supplementary-material>

REFERENCES

- Al Lawati, N. M., Patel, S. R., and Ayas, N. T. (2009). Epidemiology, risk factors, and consequences of obstructive sleep apnea and short sleep duration. *Prog. Cardiovasc. Dis.* 51, 285–293. doi: 10.1016/j.pcad.2008.08.001
- Anders, S., and Huber, W. (2010). Differential expression analysis for sequence count data. *Genome Biol.* 11:R106. doi: 10.1186/gb-2010-11-10-r106
- Baguet, J. P., Barone-Rochette, G., Tamisier, R., Levy, P., and Pepin, J. L. (2012). Mechanisms of cardiac dysfunction in obstructive sleep apnea. *Nat. Rev. Cardiol.* 9, 679–688. doi: 10.1038/nrcardio.2012.141
- Benjamin, E. J., Muntner, P., Alonso, A., Bittencourt, M. S., Callaway, C. W., Carson, A. P., et al. (2019). Heart Disease and Stroke Statistics-2019 Update: a Report From the American Heart Association. *Circulation* 139, e56–e528. doi: 10.1161/CIR.0000000000000659
- Bonney, S., Kominsky, D., Brodsky, K., Eltzschig, H., Walker, L., and Eckle, T. (2013). Cardiac Per2 functions as novel link between fatty acid metabolism and myocardial inflammation during ischemia and reperfusion injury of the heart. *PLoS One* 8:e71493. doi: 10.1371/journal.pone.0071493
- Brooks, G., Poolman, R. A., and Li, J. M. (1998). Arresting developments in the cardiac myocyte cell cycle: role of cyclin-dependent kinase inhibitors. *Cardiovasc. Res.* 39, 301–311. doi: 10.1016/s0008-6363(98)00125-4
- Brunt, V. E., Gioscia-Ryan, R. A., Richey, J. J., Zigler, M. C., Cuevas, L. M., Gonzalez, A., et al. (2019). Suppression of the gut microbiome ameliorates age-related arterial dysfunction and oxidative stress in mice. *J. Physiol.* 597, 2361–2378. doi: 10.1113/jp277336
- Buniello, A., MacArthur, J. A. L., Cerezo, M., Harris, L. W., Hayhurst, J., Malangone, C., et al. (2019). The NHGRI-EBI GWAS Catalog of published genome-wide association studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res.* 47, D1005–D1012. doi: 10.1093/nar/gky1120
- Chu, C., Murdock, M. H., Jing, D., Won, T. H., Chung, H., Kressel, A. M., et al. (2019). The microbiota regulate neuronal function and fear extinction learning. *Nature* 574, 543–548. doi: 10.1038/s41586-019-1644-y
- Clerk, A., Cullingford, T. E., Fuller, S. J., Giraldo, A., Markou, T., Pikkariainen, S., et al. (2007). Signaling pathways mediating cardiac myocyte gene expression in physiological and stress responses. *J. Cell Physiol.* 212, 311–322. doi: 10.1002/jcp.21094
- Dejea, C. M., Wick, E. C., Hechenbleikner, E. M., White, J. R., Mark Welch, J. L., Rossetti, B. J., et al. (2014). Microbiota organization is a distinct feature of proximal colorectal cancers. *Proc. Natl. Acad. Sci. U. S. A.* 111, 18321–18326. doi: 10.1073/pnas.1406199111
- Dennis, G. Jr., Sherman, B. T., Hosack, D. A., Yang, J., Gao, W., Lane, H. C., et al. (2003). DAVID: database for Annotation, Visualization, and Integrated Discovery. *Genome Biol.* 4:P3.
- Didion, J. P., Martin, M., and Collins, F. S. (2017). Atropos: specific, sensitive, and speedy trimming of sequencing reads. *PeerJ* 5:e3720. doi: 10.7717/peerj.3720
- Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., et al. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29, 15–21. doi: 10.1093/bioinformatics/bts635
- Dominguez-Bello, M. G., Godoy-Vitorino, F., Knight, R., and Blaser, M. J. (2019). Role of the microbiome in human development. *Gut* 68, 1108–1114. doi: 10.1136/gutjnl-2018-317503
- Drager, L. F., and Lee, C. H. (2018). Treatment of obstructive sleep apnoea as primary or secondary prevention of cardiovascular disease: where do we stand now? *Curr. Opin. Pulm. Med.* 24, 537–542. doi: 10.1097/MCP.0000000000000523
- Drager, L. F., Polotsky, V. Y., O'donnell, C. P., Cravo, S. L., Lorenzi-Filho, G., and Machado, B. H. (2015). Translational approaches to understanding metabolic dysfunction and cardiovascular consequences of obstructive sleep apnea. *Am. J. Physiol. Heart Circ. Physiol.* 309, H1101–H1111. doi: 10.1152/ajpheart.00094.2015
- Drexler, H. (1999). Nitric oxide synthases in the failing human heart: a doubled-edged sword? *Circulation* 99, 2972–2975. doi: 10.1161/01.cir.99.23.2972
- Dumitrascu, R., Heitmann, J., Seeger, W., Weissmann, N., and Schulz, R. (2013). Obstructive sleep apnea, oxidative stress and cardiovascular disease: lessons from animal studies. *Oxid. Med. Cell. Longev.* 2013:234631. doi: 10.1155/2013/234631
- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V., et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC Bioinformatics* 18:142. doi: 10.1186/s12859-017-1559-2
- Fan, C., Iacobas, D. A., Zhou, D., Chen, Q., Lai, J. K., Gavrialov, O., et al. (2005). Gene expression and phenotypic characterization of mouse heart after chronic constant or intermittent hypoxia. *Physiol. Genomics* 22, 292–307. doi: 10.1152/physiolgenomics.00217.2004
- Farhana, L., Banerjee, H. N., Verma, M., and Majumdar, A. P. N. (2018). Role of Microbiome in Carcinogenesis Process and Epigenetic Regulation of Colorectal Cancer. *Methods Mol. Biol.* 1856, 35–55. doi: 10.1007/978-1-4939-8751-1_3
- Ganguly, S., Mitra, A., and Sarkar, S. (2014). Role of alpha-crystallin B in regulation of stress induced cardiomyocyte apoptosis. *Cardiovasc. Hematol. Agents Med. Chem.* 12, 60–65. doi: 10.2174/1871525713666150123151731
- Gilbert, J. A., Blaser, M. J., Caporaso, J. G., Jansson, J. K., Lynch, S. V., and Knight, R. (2018). Current understanding of the human microbiome. *Nat. Med.* 24, 392–400. doi: 10.1038/nm.4517
- Guan, F., Yang, X., Li, J., Dong, W., Zhang, X., Liu, N., et al. (2019). New Molecular Mechanism Underlying Myc-Mediated Cytochrome P450 2E1 Upregulation in Apoptosis and Energy Metabolism in the Myocardium. *J. Am. Heart Assoc.* 8:e009871. doi: 10.1161/JAHA.118.009871
- Haghikia, A., Stapel, B., Hoch, M., and Hilfiker-Kleiner, D. (2011). STAT3 and cardiac remodeling. *Heart Fail. Rev.* 16, 35–47. doi: 10.1007/s10741-010-9170-x
- Harhous, Z., Booz, G. W., Ovize, M., Bidaux, G., and Kurdi, M. (2019). An Update on the Multifaceted Roles of STAT3 in the Heart. *Front. Cardiovasc. Med.* 6:150. doi: 10.3389/fcvm.2019.00150
- Harikrishnan, S. (2019). Diet, the Gut Microbiome and Heart Failure. *Card. Fail. Rev.* 5, 119–122. doi: 10.15420/cfr.2018.39.2
- Hauck, L., Grothe, D., and Billia, F. (2016). p21(CIP1/WAF1)-dependent inhibition of cardiac hypertrophy in response to Angiotensin II involves Akt/Myc and pRb signaling. *Peptides* 83, 38–48. doi: 10.1016/j.peptides.2016.07.003
- Helmink, B. A., Khan, M. A. W., Hermann, A., Gopalakrishnan, V., and Wargo, J. A. (2019). The microbiome, cancer, and cancer therapy. *Nat. Med.* 25, 377–388. doi: 10.1038/s41591-019-0377-7
- Herring, J. A., Elison, W. S., and Tessem, J. S. (2019). Function of Nr4a Orphan Nuclear Receptors in Proliferation, Apoptosis and Fuel Utilization Across Tissues. *Cells* 8:1373. doi: 10.3390/cells8111373
- Hou, N., Ye, B., Li, X., Margulies, K. B., Xu, H., Wang, X., et al. (2016). Transcription Factor 7-like 2 Mediates Canonical Wnt/beta-Catenin Signaling and c-Myc Upregulation in Heart Failure. *Circ. Heart Fail.* 9:e003010. doi: 10.1161/CIRCHEARTFAILURE.116.003010
- Hoyos, C. M., Drager, L. F., and Patel, S. R. (2017). OSA and cardiometabolic risk: what's the bottom line? *Respirology* 22, 420–429. doi: 10.1111/resp.12984
- Huang, D. W., Sherman, B. T., Tan, Q., Collins, J. R., Alvord, W. G., Roayaei, J., et al. (2007). The DAVID Gene Functional Classification Tool: a novel biological module-centric algorithm to functionally analyze large gene lists. *Genome Biol.* 8:R183. doi: 10.1186/gb-2007-8-9-r183
- Hunyor, I., and Cook, K. M. (2018). Models of intermittent hypoxia and obstructive sleep apnea: molecular pathways and their contribution to cancer. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 315, R669–R687. doi: 10.1152/ajpregu.00036.2018
- Kanaan, A., Farhani, R., Douglas, R. M., Lamanna, J. C., and Haddad, G. G. (2006). Effect of chronic continuous or intermittent hypoxia and reoxygenation on cerebral capillary density and myelination. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 290, R1105–R1114. doi: 10.1152/ajpregu.00535.2005
- Kanehisa, M., and Goto, S. (2000). KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* 28, 27–30. doi: 10.1093/nar/28.1.27
- Karbach, S. H., Schonfelder, T., Brandao, I., Wilms, E., Hormann, N., Jackel, S., et al. (2016). Gut Microbiota Promote Angiotensin II-Induced Arterial Hypertension and Vascular Dysfunction. *J. Am. Heart Assoc.* 5:e003698. doi: 10.1161/JAHA.116.003698
- Kellermayer, R., Balasa, A., Zhang, W., Lee, S., Mirza, S., Chakravarty, A., et al. (2010). Epigenetic maturation in colonic mucosa continues beyond infancy in mice. *Hum. Mol. Genet.* 19, 2168–2176. doi: 10.1093/hmg/ddq095
- Koch, C. D., Gladwin, M. T., Freeman, B. A., Lundberg, J. O., Weitzberg, E., and Morris, A. (2017). Enterosalivary nitrate metabolism and the microbiome:

- intersection of microbial metabolism, nitric oxide and diet in cardiac and pulmonary vascular health. *Free Radic. Biol. Med.* 105, 48–67. doi: 10.1016/j.freeradbiomed.2016.12.015
- Kolodziejczyk, A. A., Federici, S., Zmora, N., Mohapatra, G., Dori-Bachash, M., Hornstein, S., et al. (2020). Acute liver failure is regulated by MYC- and microbiome-dependent programs. *Nat. Med.* 26, 1899–1911. doi: 10.1038/s41591-020-1102-2
- Levy, P., Kohler, M., McNicholas, W. T., Barbe, F., Mcevoy, R. D., Somers, V. K., et al. (2015). Obstructive sleep apnoea syndrome. *Nat. Rev. Dis. Primers* 1:15015. doi: 10.1038/nrdp.2015.15
- Levy, P., Ryan, S., Oldenburg, O., and Parati, G. (2013). Sleep apnoea and the heart. *Eur. Respir. Rev.* 22, 333–352. doi: 10.1183/09059180.00004513
- Li, G. H., Arora, P. D., Chen, Y., McCulloch, C. A., and Liu, P. (2012). Multifunctional roles of gelsolin in health and diseases. *Med. Res. Rev.* 32, 999–1025. doi: 10.1002/med.20231
- Li, H., and Forstermann, U. (2000). Nitric oxide in the pathogenesis of vascular disease. *J. Pathol.* 190, 244–254. doi: 10.1002/(SICI)1096-9896(200002)190:3<244::AID-PATH575<3.0.CO;2-8
- Liang, Y., Bradford, W. H., Zhang, J., and Sheikh, F. (2018). Four and a half LIM domain protein signaling and cardiomyopathy. *Biophys. Rev.* 10, 1073–1085. doi: 10.1007/s12551-018-0434-3
- Lucking, E. F., O'Connor, K. M., Strain, C. R., Fouhy, F., Bastiaanssen, T. F. S., Burns, D. P., et al. (2018). Chronic intermittent hypoxia disrupts cardiorespiratory homeostasis and gut microbiota composition in adult male guinea-pigs. *EBioMedicine* 38, 191–205. doi: 10.1016/j.ebiom.2018.11.010
- Lynch, S. V., and Pedersen, O. (2016). The Human Intestinal Microbiome in Health and Disease. *N. Engl. J. Med.* 375, 2369–2379. doi: 10.1056/NEJMra1600266
- Lyons, O. D., and Bradley, T. D. (2015). Heart Failure and Sleep Apnea. *Can. J. Cardiol.* 31, 898–908. doi: 10.1016/j.cjca.2015.04.017
- Lyte, J. M., Gheorghe, C. E., Goodson, M. S., Kelley-Loughnane, N., Dinan, T. G., Cryan, J. F., et al. (2020). Gut-brain axis serotonergic responses to acute stress exposure are microbiome-dependent. *Neurogastroenterol. Motil.* 32:e13881. doi: 10.1111/nmo.13881
- Manichanh, C., Borrueal, N., Casellas, F., and Guarner, F. (2012). The gut microbiota in IBD. *Nat. Rev. Gastroenterol. Hepatol.* 9, 599–608. doi: 10.1038/nrgastro.2012.152
- May, A. M., Van Wagoner, D. R., and Mehra, R. (2017). OSA and Cardiac Arrhythmogenesis: mechanistic Insights. *Chest* 151, 225–241. doi: 10.1016/j.chest.2016.09.014
- McEvoy, R. D., Antic, N. A., Heeley, E., Luo, Y., Ou, Q., Zhang, X., et al. (2016). CPAP for Prevention of Cardiovascular Events in Obstructive Sleep Apnea. *N. Engl. J. Med.* 375, 919–931. doi: 10.1056/NEJMoa1606599
- Mills, R. H., Wozniak, J. M., Vrbanc, A., Campeau, A., Chassaing, B., Gewirtz, A., et al. (2020). Organ-level protein networks as a reference for the host effects of the microbiome. *Genome Res.* 30, 276–286. doi: 10.1101/gr.256875.119
- Moody, L. V., Miyamoto, Y., Ang, J., Richter, P. J., and Eckmann, L. (2019). Evaluation of Peroxides and Chlorine Oxides as Disinfectants for Chemical Sterilization of Gnotobiotic Rodent Isolators. *J. Am. Assoc. Lab. Anim. Sci.* 58, 558–568. doi: 10.30802/AALAS-JAALAS-18-000130
- Moreno-Indias, I., Torres, M., Montserrat, J. M., Sanchez-Alcoholado, L., Cardona, F., Tinahones, F. J., et al. (2015). Intermittent hypoxia alters gut microbiota diversity in a mouse model of sleep apnoea. *Eur. Respir. J.* 45, 1055–1065. doi: 10.1183/09031936.00184314
- Moreno-Indias, I., Torres, M., Sanchez-Alcoholado, L., Cardona, F., Almendros, I., Gozal, D., et al. (2016). Normoxic Recovery Mimicking Treatment of Sleep Apnea Does Not Reverse Intermittent Hypoxia-Induced Bacterial Dysbiosis and Low-Grade Endotoxemia in Mice. *Sleep* 39, 1891–1897. doi: 10.5665/sleep.6176
- Nagatomo, Y., and Tang, W. H. (2015). Intersections Between Microbiome and Heart Failure: revisiting the Gut Hypothesis. *J. Card. Fail.* 21, 973–980. doi: 10.1016/j.cardfail.2015.09.017
- Nakayama, H., Fujio, Y., and Yamaguchi, O. (2013). [Calcium dependent signaling in cardiac hypertrophy and cell death]. *Clin. Calcium* 23, 505–515. doi: 10.1007/978-4-431-55515-5
- Nemet, I., Saha, P. P., Gupta, N., Zhu, W., Romano, K. A., Skye, S. M., et al. (2020). A Cardiovascular Disease-Linked Gut Microbial Metabolite Acts via Adrenergic Receptors. *Cell* 180, 862–877.e22. doi: 10.1016/j.cell.2020.02.016
- Ogata, H., Goto, S., Sato, K., Fujibuchi, W., Bono, H., and Kanehisa, M. (1999). KEGG: kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res.* 27, 29–34. doi: 10.1093/nar/27.1.29
- Opferman, J. T., and Kothari, A. (2018). Anti-apoptotic BCL-2 family members in development. *Cell Death Differ.* 25, 37–45. doi: 10.1038/cdd.2017.170
- Ordovas, J. M., and Mooser, V. (2006). Metagenomics: the role of the microbiome in cardiovascular diseases. *Curr. Opin. Lipidol.* 17, 157–161. doi: 10.1097/01.mol.0000217897.75068.ba
- Park, Y. S., Kim, S. H., Park, J. W., Kho, Y., Seok, P. R., Shin, J. H., et al. (2020). Melatonin in the colon modulates intestinal microbiota in response to stress and sleep deprivation. *Intest. Res.* 18, 325–336. doi: 10.5217/ir.2019.00093
- Patrignani, P., Tacconelli, S., and Bruno, A. (2014). Gut microbiota, host gene expression, and aging. *J. Clin. Gastroenterol.* 48, S28–S31. doi: 10.1097/MCG.0000000000000229
- Paul, B., Barnes, S., Demark-Wahnefried, W., Morrow, C., Salvador, C., Skibola, C., et al. (2015). Influences of diet and the gut microbiome on epigenetic modulation in cancer and other diseases. *Clin. Epigenetics* 7:112. doi: 10.1186/s13148-015-0144-7
- Peucker, K., Muff, S., Wang, J., Kunzel, S., Bosse, E., Zeissig, Y., et al. (2016). Epithelial calcineurin controls microbiota-dependent intestinal tumor development. *Nat. Med.* 22, 506–515. doi: 10.1038/nm.4072
- Pevsner-Fischer, M., Zmora, N., Braverman, S., and Elinav, E. (2017). “Epigenetics and the Microbiome,” in *Handbook of Nutrition, Diet, and Epigenetics*, eds V. Patel and V. Preedy (Cham: Springer International Publishing), 1–25. doi: 10.1007/978-3-319-31143-2_35-1
- Prichard, B. N., Cruickshank, J. M., and Graham, B. R. (2001). Beta-adrenergic blocking drugs in the treatment of hypertension. *Blood Press.* 10, 366–386. doi: 10.1080/080370501753400665
- Qin, Y., and Wade, P. A. (2018). Crosstalk between the microbiome and epigenome: messages from bugs. *J. Biochem.* 163, 105–112. doi: 10.1093/jb/mvx080
- Rengarajan, S., Knoop, K. A., Rengarajan, A., Chai, J. N., Grajales-Reyes, J. G., Samineni, V. K., et al. (2020). A Potential Role for Stress-Induced Microbial Alterations in IgA-Associated Irritable Bowel Syndrome with Diarrhea. *Cell Rep. Med.* 1:100124. doi: 10.1016/j.xcrm.2020.100124
- Richards, A. L., Muehlbauer, A. L., Alazizi, A., Burns, M. B., Findley, A., Messina, F., et al. (2019). Gut Microbiota Has a Widespread and Modifiable Effect on Host Gene Regulation. *mSystems* 4:e00323. doi: 10.1128/mSystems.00323-18
- Rohini, A., Agrawal, N., Koyani, C. N., and Singh, R. (2010). Molecular targets and regulators of cardiac hypertrophy. *Pharmacol. Res.* 61, 269–280. doi: 10.1016/j.phrs.2009.11.012
- Sandek, A., Bauditz, J., Swidsinski, A., Buhner, S., Weber-Eibel, J., Von Haehling, S., et al. (2007). Altered intestinal function in patients with chronic heart failure. *J. Am. Coll. Cardiol.* 50, 1561–1569. doi: 10.1016/j.jacc.2007.07.016
- Sfanos, K. S., Yegnasubramanian, S., Nelson, W. G., and De Marzo, A. M. (2018). The inflammatory microenvironment and microbiome in prostate cancer development. *Nat. Rev. Urol.* 15, 11–24. doi: 10.1038/nrurol.2017.167
- Sforza, E., and Roche, F. (2016). Chronic intermittent hypoxia and obstructive sleep apnea: an experimental and clinical approach. *Hypoxia* 4, 99–108. doi: 10.2147/HP.S103091
- Takahashi, K., Sugi, Y., Nakano, K., Tsuda, M., Kurihara, K., Hosono, A., et al. (2011). Epigenetic control of the host gene by commensal bacteria in large intestinal epithelial cells. *J. Biol. Chem.* 286, 35755–35762. doi: 10.1074/jbc.M111.271007
- Takeda, T., Otsu, K., and Hori, M. (2005). [Calcium channel blocker and cardiac hypertrophy]. *Clin. Calcium* 15, 1645–1649. doi: 10.1007/s1016451649
- Tang, D., Kang, R., Berghie, T. V., Vandenabeele, P., and Kroemer, G. (2019). The molecular machinery of regulated cell death. *Cell Res.* 29, 347–364. doi: 10.1038/s41422-019-0164-5
- Team, N. I. H. H. M. P. A. (2019). A review of 10 years of human microbiome research activities at the US National Institutes of Health, Fiscal Years 2007–2016. *Microbiome* 7:31. doi: 10.1186/s40168-019-0620-y
- Tian, P., O’riordan, K. J., Lee, Y. K., Wang, G., Zhao, J., Zhang, H., et al. (2020). Towards a psychobiotic therapy for depression: bifidobacterium breve CCFM1025 reverses chronic stress-induced depressive symptoms and gut microbial abnormalities in mice. *Neurobiol. Stress* 12:100216. doi: 10.1016/j.ynstr.2020.100216

- Tripathi, A., Melnik, A. V., Xue, J., Poulsen, O., Meehan, M. J., Humphrey, G., et al. (2018). Intermittent Hypoxia and Hypercapnia, a Hallmark of Obstructive Sleep Apnea, Alters the Gut Microbiome and Metabolome. *mSystems* 3, e00020–18. doi: 10.1128/mSystems.00020-18
- Tripathi, A., Xu, Z. Z., Xue, J., Poulsen, O., Gonzalez, A., Humphrey, G., et al. (2019). Intermittent Hypoxia and Hypercapnia Reproducibly Change the Gut Microbiome and Metabolome across Rodent Model Systems. *mSystems* 4, e00058–19. doi: 10.1128/mSystems.00058-19
- Tsang, S., Woo, A. Y., Zhu, W., and Xiao, R. P. (2010). Deregulation of RGS2 in cardiovascular diseases. *Front. Biosci.* 2:547–557. doi: 10.2741/s84
- Turnbull, C. D. (2018). Intermittent hypoxia, cardiovascular disease and obstructive sleep apnoea. *J. Thorac. Dis.* 10, S33–S39. doi: 10.21037/jtd.2017.10.33
- Veeraveedu, P. T., Sanada, S., Okuda, K., Fu, H. Y., Matsuzaki, T., Araki, R., et al. (2017). Ablation of IL-33 gene exacerbate myocardial remodeling in mice with heart failure induced by mechanical stress. *Biochem. Pharmacol.* 138, 73–80. doi: 10.1016/j.bcp.2017.04.022
- Vinje, S., Stroes, E., Nieuwdorp, M., and Hazen, S. L. (2014). The gut microbiome as novel cardio-metabolic target: the time has come! *Eur. Heart J.* 35, 883–887. doi: 10.1093/eurheartj/eh467
- Vuong, H. E., Pronovost, G. N., Williams, D. W., Coley, E. J. L., Siegler, E. L., Qiu, A., et al. (2020). The maternal microbiome modulates fetal neurodevelopment in mice. *Nature* 586, 281–286. doi: 10.1038/s41586-020-2745-3
- Wagner, M., and Siddiqui, M. A. (2009). Signaling networks regulating cardiac myocyte survival and death. *Curr. Opin. Investig. Drugs* 10, 928–937.
- Wang, B. F., and Yoshioka, J. (2017). The Emerging Role of Thioredoxin-Interacting Protein in Myocardial Ischemia/Reperfusion Injury. *J. Cardiovasc. Pharmacol. Ther.* 22, 219–229. doi: 10.1177/1074248416675731
- Wang, R. X., Lee, J. S., Campbell, E. L., and Colgan, S. P. (2020). Microbiota-derived butyrate dynamically regulates intestinal homeostasis through regulation of actin-associated protein synaptopodin. *Proc. Natl. Acad. Sci. U. S. A.* 117, 11648–11657. doi: 10.1073/pnas.1917597117
- Weger, B. D., Gobet, C., Yeung, J., Martin, E., Jimenez, S., Betrisey, B., et al. (2019). The Mouse Microbiome Is Required for Sex-Specific Diurnal Rhythms of Gene Expression and Metabolism. *Cell Metab.* 29, 362–382.e8. doi: 10.1016/j.cmet.2018.09.023
- Weis, M. (2018). Impact of the gut microbiome in cardiovascular and autoimmune diseases. *Clin. Sci.* 132, 2387–2389. doi: 10.1042/CS20180410
- Xu, C., Lee, S. K., Zhang, D., and Frenette, P. S. (2020). The Gut Microbiome Regulates Psychological-Stress-Induced Inflammation. *Immunity* 53, 417–428.e4. doi: 10.1016/j.immuni.2020.06.025
- Xue, J., Zhou, D., Poulsen, O., Imamura, T., Hsiao, Y. H., Smith, T. H., et al. (2017). Intermittent Hypoxia and Hypercapnia Accelerate Atherosclerosis, Partially via Trimethylamine-Oxide. *Am. J. Respir. Cell Mol. Biol.* 57, 581–588. doi: 10.1165/rcmb.2017-0086OC
- Yamamura, Y., Yano, I., Kudo, T., and Shibata, S. (2010). Time-dependent inhibitory effect of lipopolysaccharide injection on Per1 and Per2 gene expression in the mouse heart and liver. *Chronobiol. Int.* 27, 213–232. doi: 10.3109/07420521003769111
- Yotti, R., Seidman, C. E., and Seidman, J. G. (2019). Advances in the Genetic Basis and Pathogenesis of Sarcomere Cardiomyopathies. *Annu. Rev. Genomics Hum. Genet.* 20, 129–153. doi: 10.1146/annurev-genom-083118-015306
- Zhou, D., Wang, J., Zapala, M. A., Xue, J., Schork, N. J., and Haddad, G. G. (2008). Gene expression in mouse brain following chronic hypoxia: role of sarcospan in glial cell death. *Physiol. Genomics* 32, 370–379. doi: 10.1152/physiolgenomics.00147.2007

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Zhou, Xue, Miyamoto, Poulsen, Eckmann and Haddad. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.