UC Santa Cruz UC Santa Cruz Electronic Theses and Dissertations

Title

Parallel evolution of the summer steelhead ecotype in multiple populations from Oregon and Northern California

Permalink https://escholarship.org/uc/item/2z65n7mb

Author Arciniega, Martha

Publication Date 2016

Peer reviewed|Thesis/dissertation

UNIVERSITY OF CALIFORNIA

SANTA CRUZ

PARALLEL EVOLUTION OF THE SUMMER STEELHEAD ECOTYPE IN MULTIPLE POPULATIONS FROM OREGON AND NORTHERN CALIFORNIA

A thesis submitted in partial satisfaction of the requirements for the degree of

MASTER OF SCIENCE

in

OCEAN SCIENCES

by Martha Angélica Arciniega Hernández June 2017

> The Thesis of Martha Angélica Arciniega Hernández is approved:

Carlos Garza, Ph.D., Chair

Professor Jonathan Zehr

Professor Grant Pogson

Devon Pearse, Ph.D.

Tyrus Miller Vice Provost and Dean of Graduate Studies Copyright © by

Martha A. Arciniega

2017

Table of Contents

iv
v
vi
vii
ix
X
s in
1
1
6
6
6
7
9
9
9
10
11
11
12
12
13
16
18

List of Tables

- 1. Summary statistics for all populations included in the present study. N = Sample size, Obs Hz = Observed Heterozygosity, A_r Mic = allelic richness for microsatellite loci only. Summer-run steelhead populations designated with **(S)**.
- 2. Pairwise F_{ST} values for the eight focal winter- and summer-run steelhead population pairs. * = significant (P < 0.05). S = summer ecotype
- 3. AMOVA results evaluating different hypotheses for groupings of the focal populations. Nb = number of groups, Var = covariance component, % = percent overall genetic variance, and F-statistics (F_{CT}, F_{SC}, F_{ST}) appropriate for each level of comparison. Bold values are significantly different from zero (P < 0.05)

List of Figures

- 1. Map showing sample locations for 25 populations, including pairs of winterand summer-run steelhead within the same watershed. Circles = winter ecotype, squares = summer ecotype, triangle = winter and summer ecotypes sampled in the same location (Winchester weir, Umpqua River, Oregon).
- 2. Unrooted neighbor-joining tree constructed with chord distances estimated from the allele frequency data. Edge labels are percentages from 1,000 bootstrap replicates. The length of the internal branches is in proportion to the number of trees in which that branch was found. Only bootstrap proportions above 80% are shown. An (S) denotes summer steelhead. An * denotes juvenile samples.
- 3. Regression of genetic distance (F_{sT}) on geographic distance (km) for all pairwise comparisons of populations in this study.

List of Supplemental Material

- 1. Plots made by the program *Distruct* representing results of STRUCTURE analyses for the winter- and summer-run steelhead populations from the Columbia, Umpqua, Klamath, and Eel rivers, with each fish represented by a thin, vertical bar. K = number of genetic clusters.
- **2.** Assay IDs, Assay targets, and references for microsatellite genetic markers and single nucleotide polymorphisms (SNPs) used in this study.

Parallel evolution of the summer steelhead ecotype from multiple populations in Oregon and Northern California

Martha A. Arciniega-Hernández

Abstract

Parallel adaptive divergence of migratory and reproductive behavior can occur in multiple populations when similar selection is acting on these traits. Timing of migration, sexual maturity, and reproduction can have major impacts on the dynamics and viability of a population. Life-history variation in steelhead, Oncorhynchus mykiss, including variation in anadromous run timing, reproductive maturity and spawn timing, represents an important aspect of their biology and adaptation to local habitats. Here we present a genetic analysis of naturally spawning steelhead to evaluate the genetic relationships and ancestry of summer- and winter-run reproductive ecotypes from multiple river basins in Oregon and Northern California. We infer the phylogeographic relationships among populations of both summer- and winter-run steelhead ecotypes using 12 microsatellite loci and 90 single nucleotide polymorphisms. Phylogenetic trees and analysis of molecular genetic variance revealed that pairs of phenotypically and genetically distinct reproductive ecotypes within rivers were each other's closest relatives. Isolation by distance was also observed, confirming that genetic relatedness was strongly associated with geographic distance, and indicating limited migration or gene flow among river basins. These patterns support the hypothesis that the summer-run steelhead ecotype has repeatedly evolved through parallel evolution in multiple river basins. These results, together with further investigation of the underlying molecular basis for the divergence of winter- and summer-run steelhead life-history traits, will inform management and conservation efforts for these ecotypes and improve our understanding of the role of adaptive variation in conservation genetics.

To my amazing parents

Laura H. Arciniega and Miguel Arciniega C.

To my awesome sisters

M. Bety Arciniega, Laura P. A. Lara, Elena C.S.A. McMillan

To my dear grandparents

Martha C. Arciniega, Juan Hernández, Elena A. Hernández

Acknowledgments

The text of this dissertation includes a reprint of the following previously published material: Arciniega M, Clemento AJ, Miller MR, Peterson M, Garza JC, Pearse DE (2016) Parallel evolution of the summer steelhead ecotype in multiple populations from Oregon and California. Conservation Genetics (17) 165-175. The first author of this publication, Martha Arciniega, planned and performed all laboratory experiments, collected the data, managed the data, performed the data analyses, interpreted the results, created the figures, and wrote the publication. The co-authors Anthony J Clemento, Devon E. Pearse and J. Carlos Garza listed in this publication supervised the research that forms the basis for the thesis. Micheal R. Miller contributed summer and winter-run steelhead samples from the Umpqua River used in this study. Matt R. Peterson contributed steelhead samples from the Klamath basin including the South Fork Trinity and New Rivers. This study would not have been possible without the collaboration of many agencies and biologists who participated in the collection of tissue samples from the various sampling locations. MA was partially supported by the Research Mentoring Institute at the University of California, Santa Cruz.

It has been a long road to this point and there are many people who have been a part of my academic career. Thank you, Carlos Garza for the opportunity to pursue this degree. I am thankful to my co-advisor Devon Pearse for being there (down the hall) ready to offer any advice and guidance. I'd also like to thank my committee members, Jon Zehr and Grant Pogson, for taking the time to read and comment on my thesis. Thank you to the UCSC Research Mentoring Institute Program for providing financial support, community, and mentorship.

Jon Zehr, thank you for hosting my undergraduate senior thesis. The experience I got in working in the Marine Microbial Ecology lab inspired me to pursue my graduate studies in molecular ecology. I can't go without mentioning the dedication and support that Shellie Bench offered me when I was an undergraduate researcher. Thank you Shellie for taking me under your wing for 2 years and teaching me lab techniques, writing skills, organizational skills, commitment, and dedication.

Many thanks to the entire Molecular Ecology team at the Santa Cruz Lab.. I'd like to thank Libby Gilbert-Hovarth, Eric Crandall, Eric Anderson, Hilary Starks, Alicia Abadía-Cardoso, Cassie Columbus, Vanessa Apkenas, and Thomas Ngyen for their enthusiasm, guidance, and training. I want to especially thank Anthony Clemento for his continuous help, insightful input, and discussions. Thank you, Hayley Nutzel and Diana Baestcher for being awesome office mates in the Grad Grotto and creating a great work environment. I wish you both the best as you continue pursuing your graduate degrees; stay strong.

I also want to give a huge thanks to Nancy Cox-Konolpeski, Zia Isola, Malika Bell, and Yuli Ortíz-Ortéga for their amazing guidance, mentorship, and support both in my undergraduate and graduate career.

Many thanks to my awesome, intelligent, and super strong friend Jo Sisdiskis for the advice, hugs, laughs, and snacks. Thank you, Victor Wolfgang Castro for shining your light and love on me throughout this journey. Your unconditional support, your jokes, and friendship mean the world.

xi

Thank you, Angela Quiros for being there during the late night work sessions, surf sessions, and Warriors basketball games. You helped me maintain a healthier work/life balance.

Thank you, Marina Brand for your mentorship, wisdom, and support during my time as a California Sea Grant Science Policy Fellow working with the Delta Science Program. I am grateful for the friendship, support, and words of encouragement I received from Lauren Yamane, Monica Oey, Annie Adelson, Sara Worden, Kari Eckdhal, Maren Farnum, and Alicia Amerson.

Lastly, I want to thank my family for their words of encouragement, unconditional love, support, and jokes.

Introduction

Variability in life-history traits can allow species to exploit diverse natural resources and may confer resilience to the potentially negative impacts of environmental change (eg. Price et al. 1984; Winemiller 1989; Dionne et al. 2008). Understanding the evolutionary basis of such variability is important for management and conservation purposes. However, changes in the expression of morphological, phenotypic, and behavioral traits can result from both plastic and evolutionary responses, making it difficult to identify the underlying genetic basis of phenotypic expression in an uncontrolled environment. Parallel evolution of phenotypic traits often occurs in association with similar selective pressures in distant habitats and can have a similar heritable component in closely related species (Haldane 1932). Similarities in the selective pressures experienced by populations or species can lead to convergent phenotypes (Pearse and Pogson 2000; Steiner et al. 2009; Rosenblum et al. 2010), although the underlying genetic change associated with the phenotypic trait may be the same (Cresko et al. 2004; Mundy et al. 2004; Colosimo et al. 2005; Gross et al. 2009; Miller et al. 2012; Hohenlohe et al. 2012; Protejo-Garcia et al. 2013; Pearse et al. 2014), or different (Jessen et al. 1991; Chen et al. 1997; Hoekstra and Nachman 2003). Thus, a first step in understanding the evolution of phenotypic variation in life-history traits is to compare the genetic relationships among populations with the same, and different, phenotypes. These relationships can then

1

provide a framework with which to test hypotheses about the evolutionary history of a specific phenotype (Harvey and Pagel 1991).

Species in the family Salmonidae are widely studied due to their importance in fisheries, aquaculture, and conservation. The complex spectrum of life-history traits in salmonid fishes (Quinn and Myers 2004) can be highly heritable (Bentzen et al. 2001; Nichols et al. 2008; Haidle et al. 2008; Miller et al. 2012; Abadía-Cardoso et al. 2013) and can change rapidly under strong selection (Hendry et al. 2000; Martínez et al. 2011; Pearse et al. 2014). Among the most variable and widely distributed of these species is *Oncorhynchus mykiss*, which is termed steelhead if it migrates to sea before returning to freshwater to spawn, and rainbow or redband trout when it does not. In California and Oregon, steelhead populations are categorized into distinct population segments (DPSs) for management and conservation purposes. The DPS boundaries are based on many factors, including geography, migratory behavior, spawn timing, and genetic structure (Busby et al. 1996; Garza et al. 2014). Because of spatial differences in temperature, water flow, sediment type, sedimentation rate, and other characteristics among rivers, divergent selection may drive the adaptation of steelhead to their local environment (Taylor 1991; Narum et al. 2008b; Narum et al. 2013; Matala et al. 2014). Instances of local adaptation in salmonid fishes have been found to occur at spatial scales of a few kilometers (Fraser et al. 2011; Bond et al. 2014) and have been shown to affect migratory timing (Major and Mighell 1966; Quinn and Adams 1996; Quinn et al. 1997; Robards and Quinn 2002). Thus, in order to manage O. mykiss effectively, it is important to understand both the geographic and genetic structure of its life-history variation and the adaptive evolutionary processes that give rise to it.

Anadromous migratory behavior can also differ by the season in which adult fish return to freshwater to spawn, or run timing. Run timing is commonly used to describe populations of anadromous fish, and such temporal differences in migration and reproduction have been shown to reduce gene flow and increase divergence among populations within a basin (Quinn et al. 2000; Hendry et al. 2002; Waples et al. 2004; Clemento 2006). This temporal divergence in patterns of life-history traits may reflect either the effects of local adaptation or phenotypic plasticity, or some combination of the two. Winter-run steelhead typically return to their natal river already sexually mature during the months of November to May and spawn shortly after entry into freshwater (Shapovalov and Taft 1954; Withler 1966; Busby et al. 1996). In contrast, summer-run steelhead return sexually immature from April to June, hold in cool pools during summer and fall, then mature and spawn in the winter or the following spring, often moving upstream into headwater habitat as they do so (Everest 1973; Moyle 2002). Timing of migration is therefore decoupled from timing of sexual maturity for the summer-run ecotype, but not for the winter-run ecotype. Although these summer- and winter-run phenotypes usually occur within the same river, they may be effectively allopatric, due to spatial and temporal separation in migration timing and spawning habitat. Such spatial and temporal differences have been found to exist between winter- and summer-run steelhead populations in British Columbia (Smith 1968) and the Rogue River (Everest 1973), with earlier returning

fish spawning farther upstream than later returning fish (Briggs 1953). In the summer, stream conditions tend to restrict steelhead to the lower part of the river and, when water flow increases in the fall or winter, summer-run fish move further upstream (Behnke 1992). Although there is overlap in spawn time, winter-run fish are just entering the river when the summer-run fish are moving further upstream, creating spatial separation. In Capilano River, British Columbia, the two races of summer-and winter-run steelhead are mutually viable, however, their behavior would indicate that they do not interbreed and there are large differences in the fat and gonad weights between them (Smith 1969). The morphological appearance of summer steelhead, such as pronounced hooking of the snout and lower jaw, also suggests something about their behavior. It is possible that summer steelhead may get territorial and defend their spawning sites against winter steelhead (Smith 1969). Thus, pairs of populations of these two ecotypes from the same river offer the opportunity to investigate the genetic basis of migratory/reproductive strategies.

In California, summer-run steelhead are currently found primarily in the tributaries of the Eel and Klamath river basins, although historically they were more widespread (Bjorkstedt et al. 2005). Further to the north, populations of summer-run steelhead are more common, particularly in the Columbia River basin, where the inland populations are derived from a distinct lineage of the species (Allendorf 1975). If these ecotypes evolved independently multiple times, then the summer- and winter-run steelhead from the same river would be expected to be each other's closest relative. In the Middle Fork Eel (MFE) River, winter-run steelhead populations are

4

more closely related to the sympatric summer-run populations than to winter-run populations in neighboring tributaries (Clemento 2006), and the time of divergence between ecotypes has been estimated as 16,000-28,000 years (Nielsen and Fountain 1999). Additionally, Clemento (2006) suggested that the summer phenotype has a heritable component because multiple generations of summer steelhead were each other's closest relatives when compared to neighboring *O.mykiss* populations. Although this evidence suggests that the summer ecotype has independently evolved within the Eel River, the population genetic relationships among sympatric summer-and winter-run steelhead populations in other basins remain unknown.

Microsatellite markers are highly polymorphic and informative, have a long history of use in population genetics (Sunnucks 2000), and have been used extensively to study *O. mykiss* populations (e.g Heath et al. 2002; Hauser et al. 2006; Pearse et al. 2007, 2009; Clemento et al. 2009; Martínez et al. 2011; Garza et al. 2014). However, because of the ease of genotyping and portability of the resulting data, single nucleotide polymorphism (SNP) markers are increasingly used in population genetic investigations (Morin et al. 2004), including to assess population demography, estimate phylogenetic relationships and for parentage analysis (e.g. Anderson and Garza 2006; Morin et al. 2009; Abadía-Cardoso et al. 2011, 2013; Hansen et al. 2011; Helyar et al. 2011; Anderson 2012). However, Narum et al. (2008a) concluded that although microsatellites had a higher accuracy than SNPs alone, using both types of markers in combination can yield higher assignment accuracy and provide the most power to determine population genetic relationships. Here we use 102 polymorphic markers, 12 microsatellites and 90 SNPs, to investigate the relationships between and evolution of summer- and winter-run ecotypes of naturally spawning steelhead. We examine the genetic structure of steelhead from summer- and winter-run populations in multiple rivers in Oregon and Northern California and place their relationships in a geographic context to test the hypothesis that the summer-run steelhead ecotype has evolved in parallel in multiple river basins from distinct local winter-run steelhead populations.

Methods

Sampling

Samples were obtained from four pairs of summer- and winter-run steelhead populations in the Columbia, Umpqua, Klamath, and Eel rivers (Table 1; Figure 1) in Oregon and Northern California through trapping and net capture of adults in holding pools or at weirs. Sampling at weirs was done during the peak time of river entry during each run, while summer-run adults were sampled in summer holding pools. For comparison of genetic distance and diversity, samples from winter-run steelhead from 17 additional locations in California were used (Table 1), as a way of visualizing how sympatric populations clustered relative to other coastal winter-run populations that are not in sympatry with summer steelhead.

Genetic Data Collection

DNA was extracted from dried fin clips using the DNeasy 96 filter-based nucleic acid extraction system on a BioRobot 3000 (Qiagen, Inc.), following the manufacturer's protocols. Extracted DNA was diluted 10:1 (microsatellites) or 2:1 (SNPs) with distilled water and used for polymerase chain reaction (PCR) amplification of 12 microsatellites and 90 SNPs (Supplemental Table 1). For microsatellites, PCR products of samples from the Columbia, Umpqua, Rogue and Trinity rivers were electrophoresed on ABI 377 sequencers (Applied Biosystems, Inc.). The same protocol and equipment was used to collect microsatellite genotypes for the other populations that were previously published based on data collected in the same laboratory (Clemento 2006; Garza et al. 2014). Genotypes were determined using Genescan 3.0 and Genotyper 2.1 software (Applied Biosystems). Two people scored all microsatellite genotypes independently, and discrepancies in the scores were resolved either by consensus or by re-genotyping. Otherwise, that genotype was deleted from the data set. SNPs were assayed with high throughput 96.96 dynamic genotyping arrays on an EP1 instrument (Fluidigm, Inc.), and genotypes called using Fluidigm SNP Genotyping Analysis software.

Data analysis

The microsatellite and SNP data were combined for all analyses. Basic population genetic statistics, and test for departures from Hardy-Weinberg and linkage equilibria were estimated using the GENEPOP program (v4.2; Rousset 2008). Allelic richness for microsatellites was calculated using FSTAT 2.9.3 (Goudet 2001). Individual fish were assigned to their most likely population of origin and individual migration between populations assessed using the Bayesian method for estimating population allele frequencies (Rannala and Mountain 1997) implemented in the program GeneClass2 (Piry et al. 2004). Pairwise differentiation between all pairs of populations was evaluated using F_{ST}, as estimated by Weir and Cockerham (1984) in ARLEQUIN (v3.11, Excoffier et al. 2007). Cavalli-Sforza and Edwards (1967) chord distances were calculated and used to generate a neighbor joining phylogram of populations, with the statistical support for population relationships evaluated by taking 1000 bootstrap samples from the dataset, all with the software package PHYLIP (Felsenstein 2005). The resulting phylogram was visualized using Dendroscope 3 (Huson and Scornavacca 2012). Isolation by distance (IBD) analysis was performed using GENEPOP with a Mantel test (1000 iterations) used to evaluate significance of relationships between FST and distance (km). To evaluate fractional ancestry of individuals, clustering analyses were conducted using the program STRUCTURE (v2.0; Pritchard et al. 2000), which allows migrants and individuals of mixed ancestry to be identified without a priori designation of defined populations (Pritchard 2000; Pearse and Crandall 2004). Clustering analyses were done with a range of values for the number of genetic clusters (K = 2 - 10), with 10 runs of each K value to evaluate consistency of patterns of genetic association. Results from the STRUCTURE runs were compiled and visualized using CLUMPP (Rosenberg 2004) and DISTRUCT (Jakobsson and Rosenberg 2007). ARLEQUIN was used to perform an analysis of molecular variance (AMOVA; Excoffier et al. 1992) for different a

8

priori groupings of population samples to evaluate the distribution of genetic variation among and between basins and ecotypes. Finally, to evaluate the neutrality of our loci, F_{ST} outlier tests were done using FDIST2 (Beaumont & Nichols 1996) as implemented in LOSITAN (Antão et al. 2008).

Results

Genotyping and Population Statistics

A total of 730 individual fish were successfully genotyped, with samples sizes from each population ranged from 18 to 47, and averaged 29 individuals (Table 1). The missing data rate over all loci was <1.3%, and all individuals were genotyped at a minimum of 90 loci. No consistent, significant deviations from Hardy-Weinberg or linkage equilibria were found. Observed heterozygosity ranged from 0.263 to 0.448 across all loci, and allelic richness for the microsatellites ranged from 5.9 to 8.1 (Table 1).

Genetic structure

A combined dataset with 12 microsatellite loci and 88 SNPs was used to construct an unrooted neighbor-joining phylogram (Figure 2). Two SNP markers were excluded from this analysis because they failed entirely in more than one population. Phylograms were also constructed using the microsatellite and SNP loci separately, and revealed no significant differences in topology (results not shown). The population clustering pattern in all analyses was consistent with the geographic locations of river basins (Figure 2). Summer- and winter-run steelhead from the same river basin were each other's closest relatives in all basins, and all steelhead populations generally clustered according to geographic proximity. For example, within the Eel River, winter-run steelhead from the upper mainstem (Van Arsdale) grouped more closely with summer-run steelhead from the Middle Fork Eel River than they did with winter-run steelhead from the Lawrence Creek and Hollow Tree Creek tributaries, which have their confluences with the mainstem Eel River much closer to its mouth. Similarly, winter-run steelhead from the South Fork Trinity River grouped more closely with summer-run steelhead from the South Fork Trinity River than they did with winter-run steelhead from the South Fork Trinity River grouped more closely with summer-run steelhead from the South Fork Trinity River grouped more closely with summer-run steelhead from the New River tributary of the Trinity River than with winter-run steelhead in other basins. This pattern was also consistent for the Umpqua and Columbia River pairs of steelhead ecotypes.

Individual assignments

Assignment tests reliably distinguished individuals sampled from various river locations in Oregon and California. Using a 90% probability confidence criteria, the accuracy with which fish were assigned to their population and basin of origin was 90.7% and 95.8%, respectively (data not shown). Misassignments between ecotypes were reflective of the population divergence, with 11, 6, and 1 summer/winter cross-assignments in the Umpqua, Klamath, and Eel population pairs, respectively. When no probability criterion was used, mean self-assignment percentage to population was 87% and ranged from 42-100% (Table 1). In addition, 66 of 730 fish(9.0%) were

misassigned to a population location other than the one where they were sampled and of those, 38 (5.2%) were assigned to a location outside their basin of origin.

Genetic differentiation

The proportion of genetic variation partitioned between population samples, pairwise F_{ST} , was significantly different from zero (p = 0.000) for all comparisons except that between the Umpqua summer- and winter-run fish. The samples from the Umpqua River were collected at the same weir, so were separated only temporally. Pairwise F_{ST} values between all population pairs ranged from 0.037 - 0.312, while F_{ST} values between ecotypes within basins were generally lower (0 – 0.069; Table 2). A pattern of isolation by distance was indicated by a strong relationship between genetic and geographic distance (F_{ST} vs. distance, $r^2 = 0.479$, Mantel test: P < 0.000; Figure 3).

STRUCTURE analyses

Model-based clustering analyses with STRUCTURE of the four regional ecotype pairs revealed distinct genetic groupings (Figure 4), concordant with the groupings in the phylogenetic tree; in every case, clusters were defined geographically by basin and not by ecotype. Multiple iterations for each value of K =2 -10, showed consistent patterns of population genetic division. At low hypothesized K (i.e., < 5), geographic proximity was the primary determinant of genetic similarity (Figure 4). As K increased to K > 5, subdivision and fractional ancestry that no longer made biological sense was apparent.

Analysis of molecular variance

AMOVA was used to partition and assess the significance of the variance at different levels of genetic structure with two hierarchal groupings. When populations were grouped by ecotype, the variance partitioned among groups was not significantly different from zero (p = 0.943), whereas when they were grouped by basin, it was 12.82% of the total genetic variance among groups, and highly significant (p=0.009; Table 3). Similarly, the variance among populations within groups categorized by ecotype and by basin were both significantly different from zero, representing 15.74 and 2.25% of the total variance, respectively (Table 3). Finally, the balance of the variance was partitioned within populations and accounted for 88.16 and 84.93% of the total variance for the ecotype and basin groups, respectively.

F_{ST} -outlier test

In order to identify loci that may not conform to neutral expectations, we conducted F_{ST} outlier tests on all genetic markers in this study. No consistent outlier loci were detected across multiple pairs of ecotype populations, confirming the

apparent selective neutrality of these loci and validating their use for population genetic inference (Supplemental table 2).

Discussion

In this study, the genetic structure of naturally spawning summer- and winterrun steelhead in multiple rivers of Oregon and Northern California was investigated using genotype data from 102 SNP and microsatellite loci. All analyses supported the hypothesis that populations of the two ecotypes within the same river basin are each other's closest relatives and that the summer-run life-history strategy has arisen multiple times independently in these basins. The close relationships of geographically proximate populations of both ecotypes indicate that local adaptation and genetic drift are more important for the evolution of the summer-run strategy than migration among populations of the same ecotype from different river basins. For example, the summer- and winter-run steelhead populations in the Klamath River basin are more closely related to each other than either one is to summer- and winterrun steelhead populations in the Eel River. These results suggest that isolation by distance (IBD) is a more important factor than ecotype in the genetic relationships of steelhead. IBD analysis found that almost 50% of the variation in genetic differentiation among populations from all sampling sites was explained by geographic distance (Figure 3). This pattern is consistent with previous studies of

coastal steelhead, which found similar signals of IBD (Pearse et al. 2011; Garza et al. 2014). A similar pattern of structuring by geography and not phenotype has also been observed in other anadromous and resident *O.mykiss* populations (Docker et al. 2003; Olsen et al. 2006; Pearse et al. 2007; Clemento et al. 2009).

The STRUCTURE analyses were concordant with the phylogram, with populations partitioned geographically by basin and not by ecotype. STRUCTURE can be used to identify recent migrants through individual assignment, but there was no evidence of recent migration of summer-run steelhead between rivers. The consistent grouping of the winter- and summer-run steelhead ecotypes within rivers with multiple iterations at different values of K is further evidence for a lack of gene flow among populations of the same ecotype from different rivers. The AMOVA analysis found that none of the total variation was distributed among groups when they are defined by ecotype, whereas when groups were defined by basins, a significant proportion was, concordant with the IBD and STRUCTURE results. In addition, approximately 85% of the molecular variance was contained within individual populations (Table 3). Finally, individual assignment tests yielded a high accuracy of assignment of individual samples to their corresponding populations of origin, with ~91% of fish accurately assigned to population and ~96% to basin of origin when a probability criterion was used. Additionally, the neutrality of the genetic markers used in this study was confirmed with the F_{ST} outlier test. Taken together, these results strongly support the hypothesis that the summer-run steelhead populations studied were more genetically similar to winter-run populations within

the same basin than to more geographically distant summer-run populations. Similarly, winter-run populations grouped with other populations of either ecotype primarily on the basis of geographic distance.

The mechanisms by which parallel evolution of the summer steelhead ecotype has occurred are not known. However, as these summer, or "early", returning steelhead populations colonized new habitat in the headwaters of these rivers, selection on migration timing, sexual maturity, and breeding date could potentially have been quite strong, with offspring viability increasing the earlier they returned from sea. Instances of adaptation of migration timing due to environmental change(s) are not uncommon in other salmonids. For example, pink salmon from Auke Creek, AK, have been observed to adapt to warmer water conditions by migrating earlier in the season in combination with a decrease in the number of very late returning fish (Kovach et al. 2012). Because of the tight coupling of run and spawn timing in Pacific salmonids (Flagg et al. 1995, Quinn et al. 2002), such divergent selection on run timing in may help facilitate differentiation between ecotypes, and the extent of reproductive isolation may be increased if individuals who disperse or reproduce at an intermediate time, especially hybrids, are maladapted and therefore have lower fitness (Crispo et al. 2006). Further work will help elucidate the underlying genetic basis of the observed differences in timing of return to freshwater and reproductive maturity.

Although summer- and winter-run steelhead appear to be discrete populations because of differences in timing of migration and spawning (Shapolov & Taft 1954; Royal 1972; Burgner et al. 1992; Hendry et al. 2005), as well as spawning location (Everest 1973), the extent of gene flow between ecotypes in the same basin is unknown and the ecotypes have been found to be genetically very similar in some cases (Allendorf 1975; Nielsen & Fountain 1999; Clemento 2006). Chilcote et al. (1980) found that genetic heterogeneity did not differ significantly between summerand winter-run steelhead in the Kalama River. In contrast, Leider et al. (1984) found genetic exchange was restricted between these populations because of spatial and temporal factors. In the Middle Fork Eel River, a landslide that formed a dam and paleolake during the Pleistocene era has been suggested to have facilitated gene flow between the two ecotypes by limiting upstream migration of summer-run steelhead and forcing both ecotypes to spawn in the same habitat (Mackey et al. 2011). However, the genetic divergence between summer- and winter-run steelhead populations in the other basins is similarly shallow, even though no such geologic explanation has been proposed, so it is not necessary to invoke a physical barrier to explain the contemporary genetic structure of steelhead ecotypes in the Eel River, even though gene flow may have been influenced by geologic events in the past.

Conservation Implications

Thus, the relationships between summer- and winter-run steelhead populations in the same basin have important management implications. In the United States, steelhead populations are administratively designated and managed as distinct population segments (DPSs), which are delineated on the basis of geographical, ecological, and genetic variation (Busby et al. 1996; Garza et al. 2014). Habitat loss, poor water quality, and other factors have led to the decline of many steelhead populations and some populations are now listed as threatened species under the US Endangered Species Act (NOAA 2006). However, ecotypic run-timing differentiation is not specifically accounted for in considering DPS designations, despite the fact that access to summer-run habitat has been dramatically reduced by dam construction. Such habitat modifications may increase the potential for gene flow among ecotypes, leading to changes in their relationships. Finally, hatchery supplementation programs that support wild populations of summer or winter steelhead need to consider if individuals from different run timings represent distinct sub populations in order to effectively maintain this phenotypic variation in captive broodstock.

References

- Abadía-Cardoso A, Clemento AJ, Garza JC (2011) Discovery and characterization of single-nucleotide polymorphisms in steelhead/rainbow trout, Oncorhynchus mykiss. Molecular Ecology Resources 11 31-49.
- Abadía-Cardoso A, Anderson EC, Pearse DE, Garza JC (2013) Large-scale parentage analysis reveals reproductive patterns and heritability of spawn timing in a hatchery population of steelhead (*Oncorhynchus mykiss*). *Molecular Ecology* 22 4733-4746.
- Aguilar A, Garza JC (2008) Isolation of 15 single nucleotide polymorphisms from coastal steelhead, *Oncorhynchus mykiss* (Salmonidae). *Molecular Ecology Resources* **8** 659-662.
- Antão T, Lopes A, Lopes RJ, Beja-Pereira A, Luikart G (2008) LOSITAN: A workbench to detect molecular adaptation based on a F_{st}-outlier method. *BMC Bioninformatics* 9 323-328.
- Anderson EC (2012) Large-scale parentage inference with SNPs: an efficient algorithm for statistical confidence of parent pair allocations. *Statistical Applications in Genetics and Molecular Biology* **11** 1081-1090.
- Allendorf FW (1975) Genetic variability in a species possessing extensive gene duplication: genetic interpretation of duplicate loci and examination of genetic variation in populations of rainbow trout. Ph.D. Dissertation, University of Washington, Seattle.
- Anderson EC, Garza JC (2006) The power of single-nucleotide polymorphisms for large-scale parentage inference. *Genetics* **172** 2567-2582.
- Beaumont MA & Nichols RA (1996) Evaluating loci for use in the genetic analysis of population structure. Proceedings of the Royal Society of London, Series B: Biological Sciences 263: 1619-1626.
- Behnke RJ (1992) Native Trout of Western North America. American Fisheries Society Monographs, 6, 275 p. American Fisheries Society, Bethesda, MD.

- Bentzen P, Olsen JB, McLean JE, Seamons TR, Quinn TP (2001) Kinship analysis of Pacific salmon: insight into mating, homing, and timing of reproduction. *Journal of Heredity* 92 344 127-136.
- Bjorkstedt EP, Spence BC, Garza JC, Hankin DG, Fuller D, Smith JJ, Macedo R (2005) An analysis of historical population structure for evolutionarily significant units of Chinook salmon, Coho salmon, and steelhead in the northcentral California coast recovery domain. NOAA Technical Memorandum NMFS-SWFSC-382. 210 p.
- Bond MH, Crane PA, Larson WA, Quinn TP (2014) Is isolation by adaptation driving genetic divergence among proximate Dolly Varden char populations? *Ecology and Evolution* **4** 2515-2532
- Briggs JC (1953) The behavior and reproduction of salmonid fishes in a small coastal stream. California Fish and Game, Fish Bulletin 94.
- Burgner RL, Light JT, Margolis L (1992) Distribution and origins of steelhead trout (Oncorhynchus mykiss) in offshore waters of the North Pacific Ocean. International North Pacific Fisheries Commission Bulletin 5151.
- Busby PJ, Wainwright TC, Bryant GJ, Lierheimer LJ, Waples RS, Waknitz FW, Lagomarsino IV (1996) Status review of West Coast steelhead from Washington, Idaho, Oregon, and California. NOAA Technical Memorandum NMFS-NWFSC-27.
- Cavalli-Sforza LL, Edwards AWF (1967) Phylogenetic analysis. Models and estimation procedures. *American Journal of Human Genetics* **19** 233-257.
- Chen L, DeVries AL, Cheng CC (1997) Convergent evolution of antifreeze glycoproteins in Antarctic notothenioid fish and Arctic cod. *Proceedings of the National Academy of Sciences* **94** 3817-3822.
- Chilcote MW, Crawford BA, Leider SA (1980) A genetic comparison of sympatric populations of summer and winter steelheads. *Transactions of the American Fisheries Society* **109** 203-206.
- Clemento AJ (2006) Subpopulation structure of steelhead trout (*Oncorhynchus mykiss*) in the Middle Fork Eel River as determined by microsatellite DNA polymorphisms. Masters Thesis, Humboldt State University.
- Clemento AJ, Anderson EC, Boughton D, Girman D, Garza JC (2009) Population genetic structure and ancestry of *Oncorhynchus mykiss* populations above and below dams in south-central California. *Conservation Genetics* **10** 1321-1336.

- Colosimo PF, Hosemann KE, Balabhadra S, Villarreal G, Dickson M, Grimwood J, Schmutz J, Myers RM, Schluter D, Kingsley DM (2005) Widespread parallel evolution in sticklebacks by repeated fixation of Ectodysplasin alleles. *Science* **307** 1928-1933.
- Cresko WA, Amores A, Wilson C, Murphy J, Currey M, Phillips P, Bell MA, Kimmel CB, Postlethwait JH (2004) Parallel genetic basis for repeated evolution of armor loss in Alaskan threespine stickleback populations. *Proceedings of the National Academy of Sciences* **101** 6050-6055.
- Crispo E, Bentzen P, Reznick DN, Kinnison MT, Hendry AP (2006) The relative influence of natural selection and geography on gene flow in guppies. *Molecular Ecology* **15** 49-62.
- Dionne M, Caron F, Dodson JJ, Bernatchez L (2008) Landscape genetics and hierarchical genetic structure in Atlantic salmon: the interaction of gene flow and local adaptation. *Molecular Ecology* **17** 2382-2396.
- Docker MF, Heath DD (2003) Genetic comparison between sympatric anadromous steelhead and freshwater resident rainbow trout in British Columbia, Canada. *Conservation Genetics* **4** 227-231.
- Everest FH (1973) Ecology and management of summer steelhead in the Rogue River. Oregon State Game Commission, Fishery Research Report 7, Corvallis, Oregon.
- Excoffier L, Smouse PE, Quattro JM (1992) Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* **131** 479-491.
- Excoffier L, Laval G, Schneider S (2007) ARLEQUIN, version 3.11: an integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* **1** 47-50.
- Felsenstein J (2005) PHYLIP v. 3.69 c Phylogeny Inference Package. (http://evolution.genetics.washington.edu/phylip.html) Department of Genetics,University of Washington, Seattle, WA.
- Flagg TA, Waknitz FW, Maynard DJ, Milner GB, Mahnken CVW (1995) The effects of hatcheries on native Coho salmon populations in the lower Columbia River. in: Schramm HL Jr. and Peper RG (ed) Uses and effects of cultured fishes in aquatic ecosystems, Symposium 15 Bethesda, Maryland: American Fisheries Society, pp 366-375

- Fraser DJ, Weir LK, Bernatchez L, Hansen MM, Taylor EB (2011) Extent and scale of local adaptation in salmonid fishes: review and meta-analysis. *Heredity* 106 404-420.
- Garza JC, Gilbert-Horvath EA, Spence BC, Williams TH, Fish H, Gough SA, Anderson JH, Hamm D, Anderson EC (2014) Population structure of steelhead in coastal California. *Transactions of the American Fisheries Society* 143 134-152.
- Goudet, J (2001) FSTAT, version 2.9.3, A program to estimate and test gene diversities and fixation indices. Lausanne University, Laussane, Switzerland.
- Gross JB, Borowsky R, Tabin CJ (2009) A novel role for Mc1r in the parallel evolution of depigmentation in independent populations of the cavefish *Astyanax mexicanus*. *PloS Genetics* **5**(1) e1000326.
- Haidle L, Janssen JE, Gharbi K, Moghadam HK, Ferguson MM, Danzmann RG (2008) Determination of quantitative trait loci (QTL) for early maturation in rainbow trout (*Oncorhynchus mykiss*). *Marine Biotechnology* **10** 579-592.

Haldane JBS (1932) The Causes of Evolution. Longman, London.

- Hansen MHH, Young S, Jørgensen HBH, Pascal C, Henryon M, Seeb J (2011)
 Assembling a dual purpose TaqMan-based panel of single-nucleotide polymorphism markers in rainbow trout and steelhead (*Oncorhynchus mykiss*) for association mapping and population genetics analysis. *Molecular Ecology Resources* 11 67-70.
- Harvey PH, Pagel MD (1991) The Comparative Method in Evolutionary Biology, Oxford Series in Ecology and Evolution, Oxford.
- Hauser L, Seamons TR, Dauer M, Naish KA, Quinn TP (2006) An empirical verification of population assignment methods by marking and parentage data: hatchery and wild steelhead (*Oncorhynchus mykiss*) in Forks Creek, Washington, USA. *Molecular Ecology* 15 3157-3173.
- Heath DD, Busch C, Kelly J, Atagi DY (2002) Temporal change in genetic structure and effective population size in steelhead trout (*Oncorhynchus mykiss*). *Molecular Ecology* **11** 197-214.
- Helyar SJ, Hemmer-Hansen J, Bekkevold D, Taylor MI, Odgen R, Limborg MT, Cariani A, Maes GE, Diopere E, Carvalho GR, Nielsen EE (2011) Application

of SNPs for population genetics of non-model organisms: new opportunities and challenges. *Molecular Ecology Resources* **11** 123-126.

- Hendry AP, Wenburg JK, Bentzen P, Volk EC, Quinn TP (2000) Rapid evolution of reproductive isolation in the wild: evidence from introduced salmon. *Science* 290 516-518.
- Hendry MA, Wenburg JK, Myers KW, Hendry AP (2002) Genetic and phenotypic variation through the migratory season provides evidence for multiple populations of wild steelhead in the Dean River, British Columbia. *Transactions of the American Fisheries Society* **131** 418-434.
- Hendry AP, Day T (2005) Population structure attributable to reproductive time: isolation by time and adaptation by time. *Molecular Ecology* **14** 901-916.
- Hoekstra HE, Nachman MW (2003) Different genes underlie adaptive melanism in different populations of rock pocket mice. *Molecular Ecology* **12** 1185-94.
- Hohenlohe PA, Bassham S, Currey M, Cresko WA (2012) Extensive linkage disequilibrium and parallel adaptive divergence across threespine stickleback genomes. *Philosophical Transactions of the Royal Society B: Biological Sciences* 367 395-408.
- Huson DH, Scornavacca C (2012) Dendroscope 3: an interactive tool for rooted phylogenetic trees and networks. *Systems Biology* **61** 1061-1067.
- Jakobsson M, Rosenberg NA (2007) CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* **23** 1801-1806.
- Jessen TH, Weber RE, Fermi G, Tame J, Braunitzer G (1991) Adaptation of bird hemoglobins to high altitudes: demonstration of molecular mechanism by protein engineering. *Proceedings of the National Academy of Sciences* **88** 6519-6522.

Kovach RP, Gharrett AJ, Tallmon DA (2012) Genetic change for earlier migration timing in a pink salmon population. *Proceedings of the Royal Society: Biological Sciences* **279** 3870-3878

Leider SA, Chilcote MW, Loch JJ (1984) Spawning characteristics of sympatric populations of steelhead trout (*Salmo gairdneri*): evidence for partial reproductive isolation. *Canadian Journal of Fisheries and Aquatic Sciences* 41 1454-1462.

- Mackey BH, Roering JJ, Lamb MP (2011) Landslide-dammed paleolake perturbs marine sedimentation and drives genetic change in anadromous fish. *Proceedings of the National Academy of Sciences* **108** 18905-18909.
- Major RL, Mighell JL (1966) Influence of Rocky Reach Dam and the temperature of the Okanogan River on the upstream migration of sockeye salmon. *Fishery Bulletin of the Fish and Wildlife Service* **66** 131-147.
- Martínez A, Garza JC, Pearse DE (2011) A microsatellite genome screen identifies chromosomal regions under differential selection in steelhead and rainbow trout. *Transactions of the American Fisheries Society* **140** 829-842.
- Matala AP, Ackerman MW, Campbell MR, Narum SR (2014) Relative contributions of neutral and non-neutral genetic differentiation to inform conservation of steelhead trout across highly variable landscapes. *Evolutionary Applications* 7 682-701.
- Miller MR, Brunelli JP, Wheeler PA, Liu SX, Rexroad CE, Palti Y, Doe CQ, Thorgaard GH (2012) A conserved haplotype controls parallel adaptation in geographically distant salmonid populations. *Molecular Ecology* **21** 237-249.
- Morin PA, Luikart G, Wayne RK, the SNP workshop group (2004) SNPs in ecology, evolution and conservation. *Trends in Ecology and Evolution* **19** 208-216.
- Morin PA, Martien KK, Taylor BL (2009) Assessing statistical power of SNPs for population structure and conservation studies. *Molecular Ecology Resources* 9 66-73.
- Moyle PB (2002) Inland Fishes of California, 2nd Edition, Berkeley, University of California Press.
- Mundy NI, Badcock NS, Hart T, Scribner K, Janssen K, Nadeau NJ (2004) Conserved genetic basis of a quantitative plumage trait involved in mate choice. *Science* **303** 1870-1873.
- Narum SR, Banks M, Beacham TD, Belling MR, Campbell MR, Dekoning J, Elz A, Guthrie III CM, Kozfkay C, Miller KM, Moran P, Phillips R, Seeb LW, Smith CT, Warheit K, Young SF, Garza JC (2008a) Differentiating salmon populations at broad and fine geographical scales with microsatellites and single nucleotide polymorphisms. *Molecular Ecology* **17** 3464-3477.
- Narum SR, Zendt JS, Graves D, Sharp WR (2008b) Influence of landscape on resident and anadromous life history types of *Oncorhynchus mykiss*. *Canadian Journal of Fisheries and Aquatic Sciences* **65** 1013-1023.

- Narum SR, Campbell NR, Meyer KA, Miller MR, Hardy RW (2013) Thermal adaptation and acclimation of ectotherms from differing aquatic climates. *Molecular Ecology* **22** 3090-3097.
- NOAA (2006) Endangered and threatened species: final listing determinations for 10 distinct population segments of West Coast steelhead, pp 833-862 US Federal Register 71.
- Nichols KM, Edo AF, Wheeler PA, Thorgaard GH (2008) The genetic basis of smoltification-related traits in *Oncorhynchus mykiss*. *Genetics* **179** 1559–1575.
- Nielsen MC, Fountain JL (1999) Microsatellite diversity in sympatric reproductive ecotypes of Pacific steelhead (*Oncorhynchus mykiss*) from the Middle Fork Eel River, California. *Ecology of Freshwater Fish* **8** 159-168.
- Olsen JB, Wuttig K, Fleming D, Kretschmer EJ, Wenburg JK (2006) Evidence of partial anadromy and resident-form dispersal bias on a fine scale in populations of *Oncorhynchus mykiss*. *Conservation Genetics* **7** 613-619.
- Pearse DE, Pogson GH (2000) Parallel evolution of the melanic form of the California legless lizard, *Anniella pulchra*, inferred from mitochondrial DNA sequence variation. *Evolution* **54** 1041-1046.
- Pearse DE, Crandall KA (2004) Beyond FST: analysis of population genetic data for conservation. *Conservation Genetics* **5** 585-602.
- Pearse DE, Donohoe C, Garza JC (2007) Population genetics of steelhead (*Oncorhynchus mykiss*) in the Klamath River. *Environmental Biology of Fishes* **80** 377-387.
- Pearse DE, Hayes SA, Bond MH, Hanson CV, Anderson EC, MacFarlane RB, Garza JC (2009) Over the falls? Rapid evolution of ecotypic differentiation in steelhead/rainbow trout (*Oncorhynchus mykiss*). Journal of Heredity 100 515-525.
- Pearse DE, Martinez E, Garza JC (2011) Disruption of patterns of isolation by distance in coastal steelhead. *Conservation Genetics* **12** 691-700.

Pearse DE, Miller MR, Abadía-Cardoso A, Garza JC (2014) Rapid parallel evolution of standing variation in a single, complex, genomic region is associated with life

history in steelhead/rainbow trout. *Proceedings of the Royal Society B: Biological Sciences* **281** 20140012.

- Price TD, Grant PR, Gibbs HL, Boag PT (1984) Recurrent patterns of natural selection in a population of Darwin's finches. *Nature* **309** 787-789.
- Piry S, Alapetite A, Cornuet J-M, Paetkau D, Baudouin L, Estoup A (2004) GeneClass2: a software for genetic assignment and first generation migrant detection. *Journal of Heredity* 95 536-539.
- Pritchard JK, Stephens M. Donnelly P (2000) Inference of population structure using multilocus genotype data. *Genetics* **155** 945-959.
- Protejo-Garcia J, Natarajan C, Moriyama H, Weber RE, Fago A et al. (2013)
 Repeated elevational transitions in hemoglobin function during the evolution of Andean hummingbirds. *Proceedings of the National Academy of Sciences* 110 20669-20674.
- Quinn TP, Adams DJ (1996) Environmental changes affecting the migratory timing of American shad and sockeye salmon. *Ecology* **77** 1151-1162.
- Quinn TP, Hodgson S, Peven C (1997) Temperature, flow, and the migration of adult sockeye salmon (*Oncorhynchus nerka*) in the Columbia River. *Canadian Journal of Fisheries and Aquatic Sciences* **54** 1349-1360.
- Quinn TP, Unwin MJ, Kinnison MT (2000) Evolution of temporal isolation in the wild: genetic divergence in timing of migration and breeding by introduced Chinook salmon populations. *Evolution* **54** 1372-1385.
- Quinn TP, Peterson JA, Gallucci VF, Hershberger WK, Brannon EL (2002) Artificial selection and environmental change: countervailing factors affecting the timing of spawning by Coho and Chinook salmon. *Transactions of the American Fisheries Society* **131** 591-598.
- Quinn TP, Myers KW (2004) Anadromy and the marine migrations of Pacific salmon and trout: Rounsefell revisited. *Reviews in Fish Biology and Fisheries* 14 421-442.
- Rannala B, Mountain JL (1997) Detecting immigration by using multilocus genotypes. *Proceedings of the National Academy of Sciences* **94** 9197-9201.
- Robards MD, Quinn TP (2002) The migratory timing of adult summer-run steelhead in the Columbia River over six decades of environmental change. *Transactions of the American Fisheries Society* **131** 523-536.

- Rosenberg NA (2004) Distruct: a program for the graphical display of population structure. *Molecular Ecology Notes* **4** 137-138.
- Rosenblum EB, Römpler H, Schöneberg T, Hoekstra HE (2010) Molecular and functional basis of phenotypic convergence in white lizards at White Sands. *Proceedings of the National Academy of Sciences* **107** 2113-2117.
- Rousset F (2008) GENEPOP'007: a complete re-implementation of the GENEPOP software for Windows and Linux. *Molecular Ecology Resources* **8** 103-106.
- Royal, LA (1972) An examination of the anadromous trout program of the Washington State Game Department. Washington Game Department, Olympia.
- Scribner KT, Gust JR, Fields RL (1996) Isolation and characterization of novel salmon microsatellite loci: cross-species amplification and population genetic applications. *Canadian Journal of Fisheries and Aquatic Sciences* 53 833-841.
- Shapovalov L, Taft AC (1954) The life histories of the steelhead rainbow trout (Salmo gairdneri gairdneri) and silver salmon (Oncorhynchus kisutch) with special reference to Waddell Creek, California, and recommendations regarding their management. California Department of Fish and Game, Fish Bulletin 98, Sacramento, California.
- Smith SB (1969) Reproductive isolation in summer and winter races of steelhead trout. Pages 21-38 in TG Northcote, editor. Symposium on Salmon and Trout in Streams. HR MacMillan Lectures in Fisheries. University of British Columbia, Vancouver, British Columbia, Canada.

Spies IB, Brasier DJ, O'Reilly PTL, Seamons TR, Bentzen P (2005) Development and characterization of novel tetra-, tri-, and dinucleotide microsatellite markers in rainbow trout (*Oncorhynchus mykiss*). *Molecular Ecology Notes* **5** 278-281.

- Steiner CC, Römpler H, Boettger LM, Schöneberg T, Hoekstra HE (2009) The genetic basis of phenotypic convergence in beach mice: similar pigment patterns but different genes. *Molecular Biology and Evolution* 26 35–45.
- Sunnucks P (2000) Efficient genetic markers for population biology. *Trends in Ecology and Evolution* **15** 199-203.
- Taylor EB (1991) A review of local adaptation in Salmonidae with particular reference to Pacific and Atlantic salmon. *Aquaculture* **98** 185-207.

- Waples RS, Teel DJ, Myers JM, Marshall AR (2004) Life-history divergence in Chinook salmon: historic contingency and parallel evolution. *Evolution* 58 386-403.
- Weir BS, Cockerham CC (1984) Estimating F-statistics for the analysis of population structure. *Evolution* **38** 1358-1370.
- Williamson KS, Cordes JF, May B (2002) Characterization of microsatellite loci in Chinook salmon (*Oncorhynchus tshawytscha*) and cross-species amplification in other salmonids. *Molecular Ecology Notes* 2 17-19.
- Winemiller KO (1989) Patterns of variation in life history among South American fishes in seasonal environments. *Oecologia* **81** 225-241.
- Withler IL (1966) Variability in life history characteristics of steelhead trout (Salmo gairdneri) along the Pacific coast of North America. Journal of Fisheries Research Board of Canada 23 365-393.

(S).	Heterozygosity, A_r Mic = allelic richness for microsatellite loci only. Summer-run steelhead populations of	Table 1. Summary statistics for all populations included in the present study. N = Sample size, Obs Hz =
	populations designated with	e, Obs $Hz = Observed$

24 South-Central CA coast 25	22 23	21 Central CA coast	20	19	16	18	16	15	14	13	12	11	10	9	∞ .		6	5 Klamath Mtn. Province	4	3 Oregon Coast	2 Upper Williamette	1 Lower Columbia	DPS
Carmel Big Sur	Redwood San Lorenzo	Miller	Gualala	Noyo				Eel	Mattole	Mad	ı	Trinity	Redwood		Klamath	Smith	(Rogue		Umpqua	Columbia	Columbia	Basin
Carmel R. Big Sur R.	Redwood Ck. San Lorenzo R.	Miller Ck.	Gualala R.	Noyo R.	Van Arsdale	Middle (S)	Hollow Tree	Lawrence Ck.	Mattole R.	Mad R.	South Fork	New R. (S)	Lost Man Ck.	Hunter Ck.	Blue Ck.	Smith R.	Lawson Ck.	Lobster Ck.	Umpqua R. (S)	Umpqua R.	Wiley Ck.	Still Ck. (S)	Population
$32 \\ 31$	$30 \\ 32$	$\frac{3}{2}$	29	31	23	24	28	30	31	28	47	47	31	28	32	32	23	18	22	23	23	24	z
102 102	102 102	102	102	102	101	101	102	102	102	102	102	102	102	102	102	102	101	101	102	102	101	101	Loci typed
$\substack{0.418\\0.427}$	$0.438 \\ 0.431$	0.407	0.448	0.415	0.377	0.397	0.390	0.402	0.401	0.410	0.340	0.349	0.387	0.384	0.364	0.361	0.361	0.360	0.313	0.328	0.263	0.298	Obs Hz
7.5 7.3	7.6 6.2	5.9	7.0	7.6	8.0	6.9	7.4	7.8	6.8	6.9	8.1	7.9	8.1	6.5	8.4	7.9	8.0	7.9	7.4	8.1	6.0	7.7	A _r Mic
93 80	88 88	100	92	97	95	95	79	88	96	88	83	90	68	92	100	97	98	42	56	75	56	100	% self assignment

Table 2. Pairwise F_{ST} values for the eight focal winter- and summer-run steell ecoty

Basin	Population	Wiley Ck.	Still Ck.(S)	Umpqua	Umpqua (S)	South Fortk Trinity	New R. (S)	Middle Fork
Columbia	Wiley Ck.	-						
	Still Ck. (S)	*0.068	-					
Umpqua	Umpqua	*0.104	*0.037	-				
	Umpqua (S)	*0.121	*0.048	0.000	-			
Klamath	South Fork Trinity	*0.156	*0.085	*0.069	*0.076	-		
	New River (S)	*0.156	*0.092	*0.073	*0.081	*0.016	-	
Eel	Middle Fork (S)	*0.291	*0.236	*0.198	*0.198	*0.166	*0.168	-
	Van Arsdale	*0.312	*0.246	*0.210	*0.212	*0.185	*0.185	*0.034

Table 3. AMOVA results evaluating different hypotheses for groupings of the focal populations. Nb = number of groups, Var = covariance component, % = percent overall genetic variance, and F-statistics (F_{CT} , F_{SC} , F_{ST}) appropriate for each level of comparison. Bold values are significantly different from zero (P < 0.05).

	Nb	Among	g groups	5	Amon	g popul	ations	Within populations within			
					group	S					
Description		Var	%	FCT	Var	%	FSC	Var	%	FST	
By ecotype	2	0.655	-3.90	-0.039	2.643	15.74	0.152	14.801	88.16	0.118	
By basin	4	2.234	12.82	0.128	0.392	2.25	0.026	14.801	84.93	0.151	









Figure 3



Supplemental Material Table 1.

Assay ID	Assay Target	Reference
Omy1011	(CAGA) ₁₁	Spies et al. 2005
OtsG243	$(TAGA)_{63}(CAGA)_{12}(GACA)_7(GA)_{22}$	Williamson et al. 2002
OtsG253B	(GACA)10(GATA) ₁₄	Williamson et al. 2002
One11	(CA) ₁₃	Scribner et al. 1996
OtsG249B	(TAGA) ₁₉	Williamson et al. 2002
OtsG43	(GACA)13(GATAGACA)2(GATA)25	Williamson et al. 2002
OtsG85	(GATA) ₁₉	Williamson et al. 2002
Oki23		Spidle et al. unpublished, GenBank AF272822
One13	$(GA)_{20}$	Scribner et al. 1996
OtsG3	(GATA) ₃₀ (TAGA) ₁	Williamson et al. 2002
OtsG409	(GA) ₉ (TAGA) ₆ -GGTA-(GATA) ₁₆	Williamson et al. 2002
Ssa289	(GT) ₁₂	McConnell et al. 1995
OMGH1PROM1-SNP1	A/T	Abadía-Cardoso et al. 2011
Omy_AldA	T/C	Aguilar & Garza 2008
Omy_arp-630	T/C	Campbell et al. 2009
Omy_aspAT-123	T/C	Campbell et al. 2009
Omy_COX1-221	T/A	Campbell et al. 2009
Omy_g12-82	G/A	WSU - J. DeKoning unpubl.
Omy_gh-475	G/A	Campbell et al. 2009
Omy_gsdf-291	T/C	WSU - J. DeKoning unpubl.
Omy_mapK3-103	T/A	CRITFC - N. Campbell unpubl.
Omy_mcsf-371	G/A	WSU - J. DeKoning unpubl.
Omy_nramp-146	T/C	Campbell et al. 2009
Omy_Ogo4-304	T/C	Campbell et al. 2009
OMY_PEPA-INT6	T/C	Aguilar & Garza 2008
ONMYCRBF_1-SNP1	T/C	Aguilar & Garza 2008
SH100771-63	T/A	Abadía-Cardoso et al. 2011
SH100974-386	T/C	Abadía-Cardoso et al. 2011
SH101554-306	T/C	Abadía-Cardoso et al. 2011
SH101770-410	T/C	Abadía-Cardoso et al. 2011

Supplemental Material Table 1 continued from last page.

Assay ID	Assay Target	Reference
SH101832-195	T/C	Abadía-Cardoso et al. 2011
SH101993-189	A/T	Abadía-Cardoso et al. 2011
SH102420-634	T/G	Abadía-Cardoso et al. 2011
SH102505-102	A/G	Abadía-Cardoso et al. 2011
SH102510-682	T/G	Abadía-Cardoso <i>et al</i> . 2011
SH102867-443	T/G	Abadía-Cardoso <i>et al</i> . 2011
SH103350-395	A/C	Abadía-Cardoso et al. 2011
SH103577-379	T/A	Abadía-Cardoso et al. 2011
SH103705-558	T/C	Abadía-Cardoso <i>et al</i> . 2011
SH104519-624	T/C	Abadía-Cardoso <i>et al</i> . 2011
SH105075-162	T/G	Abadía-Cardoso et al. 2011
SH105105-448	C/T	Abadía-Cardoso <i>et al</i> . 2011
SH105115-367	C/G	Abadía-Cardoso <i>et al</i> . 2011
SH105385-406	T/C	Abadía-Cardoso et al. 2011
SH105386-347	A/C	Abadía-Cardoso <i>et al</i> . 2011
SH105714-265	C/T	Abadía-Cardoso <i>et al</i> . 2011
SH106172-332	T/G	Abadía-Cardoso et al. 2011
SH106313-445	T/G	Abadía-Cardoso et al. 2011
SH107074-217	A/G	Abadía-Cardoso et al. 2011
SH107285-69	C/G	Abadía-Cardoso et al. 2011
SH108735-311	C/T	Abadía-Cardoso et al. 2011
SH109243-222	A/C	Abadía-Cardoso et al. 2011
SH109525-403	A/G	Abadía-Cardoso <i>et al</i> . 2011
SH109651-445	C/T	Abadía-Cardoso et al. 2011
SH109693-461	T/A	Abadía-Cardoso <i>et al</i> . 2011
SH109874-148	A/G	Abadía-Cardoso <i>et al</i> . 2011
SH110064-419	T/G	Abadía-Cardoso <i>et al</i> . 2011
SH110078-294	A/G	Abadía-Cardoso <i>et al</i> . 2011
SH110201-359	T/G	Abadía-Cardoso <i>et al</i> . 2011

Assay ID	Assay Target	Reference
SH110362-585	G/A	Abadía-Cardoso et al. 2011
SH110689-148	A/C	Abadía-Cardoso et al. 2011
SH111666-301	T/A	Abadía-Cardoso et al. 2011
SH112208-328	T/C	Abadía-Cardoso et al. 2011
SH112301-202	T/G	Abadía-Cardoso et al. 2011
SH112820-82	G/A	Abadía-Cardoso et al. 2011
SH113109-205	T/G	Abadía-Cardoso et al. 2011
SH113128-73	C/G	Abadía-Cardoso et al. 2011
SH114315-438	T/G	Abadía-Cardoso et al. 2011
SH114587-480	T/G	Abadía-Cardoso et al. 2011
SH114976-223	T/G	Abadía-Cardoso et al. 2011
SH115987-812	C/T	Abadía-Cardoso et al. 2011
SH116733-349	C/T	Abadía-Cardoso et al. 2011
SH117259-96	T/C	Abadía-Cardoso et al. 2011
SH117286-374	A/T	Abadía-Cardoso et al. 2011
SH117370-400	A/G	Abadía-Cardoso et al. 2011
SH117540-259	T/G	Abadía-Cardoso et al. 2011
SH117815-81	C/T	Abadía-Cardoso et al. 2011
SH118175-396	T/A	Abadía-Cardoso et al. 2011
SH118654-91	A/G	Abadía-Cardoso et al. 2011
SH118938-341	A/T	Abadía-Cardoso et al. 2011
SH119108-357	T/C	Abadía-Cardoso et al. 2011
SH119892-365	T/G	Abadía-Cardoso et al. 2011
SH120255-332	A/T	Abadía-Cardoso et al. 2011
SH120950-569	T/G	Abadía-Cardoso et al. 2011
SH123044-128	C/T	Abadía-Cardoso et al. 2011
SH125998-61	T/G	Abadía-Cardoso et al. 2011
SH127510-920	C/T	Abadía-Cardoso et al. 2011
SH128851-273	T/A	Abadía-Cardoso et al. 2011
SH129870-756	C/T	Abadía-Cardoso et al. 2011
SH130524-160	C/G	Abadía-Cardoso et al. 2011
SH130720-100	C/T	Abadía-Cardoso et al. 2011
SH131460-646	C/T	Abadía-Cardoso et al. 2011
SH131965-120	C/T	Abadía-Cardoso et al. 2011
SH95318-147	C/T	Abadía-Cardoso et al. 2011
SH95489-423	T/C	Abadía-Cardoso et al. 2011
SH96222-125	T/C	Abadía-Cardoso et al. 2011
SH97077-73	T/A	Abadía-Cardoso et al. 2011
SH97954-618	C/T	Abadía-Cardoso et al. 2011
SH98188-405	T/C	Abadía-Cardoso et al. 2011
SH98409-549	A/G	Abadía-Cardoso et al. 2011
SH98683-165	A/C	Abadía-Cardoso et al. 2011
SH99300-202	T/A	Abadía-Cardoso et al. 2011

Supplemental Material Table 1. Table 1 continued from last page

Supplemental Material Table 2.

Drainage	Markers	Sim. Fst	Outliers (Y/N)	Loci [He/Fst/P]
Columbia	Msats	0.0676	Ν	N/A
Umpqua	Msats	0.0041	Ν	N/A
Trinity	Msats	0.0172	Ν	N/A
Eel	Msats	0.0266	Ν	N/A
Columbia	SNPs	0.0703	Y	SH106313-445 [0.08/0.064/1], SH110064-419 [0.10/0.085/1]
Umpqua	SNPs	0.0023	Y	Omy_nramp-146 [0.04/0.021/1], SH106172-332 [0.04/0.021/1]
Trinity	SNPs	0.0137	Ν	N/A
Eel	SNPs	0.0259	Y	SH112301-202 [0.25/0.234/0.999]
Columbia	Both	0.0758	Y	SH106313-445 [0.08/0.064/1]
Umpqua	Both	0.0008	Y	Omy_nramp-146 [0.04/0.021/1], SH106172-332 [0.04/0.021/1]
Trinity	Both	0.0130	Ν	N/A
Eel	Both	0.0269	Y	SH112301-202 [0.25/0.234/0.999]

Supplemental Figure 1. Plots made by the program *Distruct* representing results of STRUCTURE analyses for the winter- and summer-run steelhead populations from the Columbia, Umpqua, Klamath, and Eel rivers, with each fish represented by a thin, vertical bar. K = number of genetic clusters.

