

UC Irvine

UC Irvine Previously Published Works

Title

Injury-induced maladaptation and dysregulation of calcium channel $\alpha 2\delta$ subunit proteins and its contribution to neuropathic pain development

Permalink

<https://escholarship.org/uc/item/6d57h68g>

Journal

British Journal of Pharmacology, 175(12)

ISSN

0007-1188

Authors

Gong, Nian
Park, John
Luo, Z David

Publication Date

2018-06-01

DOI

10.1111/bph.13930


Peer reviewed

REVIEW ARTICLE

Injury-induced maladaptation and dysregulation of calcium channel $\alpha_2\delta$ subunit proteins and its contribution to neuropathic pain development

Correspondence Dr Z. David Luo, Department of Anesthesiology & Perioperative Care, School of Medicine, University of California Irvine, Irvine 92697, CA, USA. E-mail: zluo@uci.edu

Received 28 February 2017; **Revised** 5 June 2017; **Accepted** 12 June 2017

Nian Gong¹, John Park² and Z David Luo^{1,2} 

¹Department of Anesthesiology & Perioperative Care, School of Medicine, University of California Irvine, Irvine, CA, USA, and ²Department of Pharmacology, School of Medicine, University of California Irvine, Irvine, CA, USA

Voltage-gated calcium channels (VGCCs) play important roles in physiological functions including the modulation of neurotransmitter release, neuronal network activities, intracellular signalling pathways and gene expression. Some pathological conditions, including nerve injuries, can cause the dysregulation of VGCCs and their subunits. This in turn can lead to a functional maladaptation of VGCCs and their subunits, which can contribute to the development of disorders such as pain sensations. This review has summarized recent findings related to maladaptive changes in the dysregulated VGCC $\alpha_2\delta_1$ subunit ($\text{Ca}_v\alpha_2\delta_1$) with a focus on exploring the mechanisms underlying the contribution of $\text{Ca}_v\alpha_2\delta_1$ to pain signal transduction. At least under neuropathic pain conditions, the dysregulated $\text{Ca}_v\alpha_2\delta_1$ can modulate VGCC functions as well as other plasticity changes. The latter includes abnormal excitatory synaptogenesis resulting from its interactions with injury-induced extracellular matrix glycoprotein molecule thrombospondins, which is independent of the VGCC functions. Blocking $\text{Ca}_v\alpha_2\delta_1$ with gabapentinoids can reverse neuropathic pain significantly with relatively mild side effects, but only in a small population of neuropathic pain patients due to reasons yet to be explored. There are emerging data suggesting that early preventive treatment with gabapentinoids can prevent aberrant excitatory synapse formation and the development of chronic pain. If these findings are confirmed clinically, this could be an attractive approach for neuropathic pain management.

LINKED ARTICLES

This article is part of a themed section on Recent Advances in Targeting Ion Channels to Treat Chronic Pain. To view the other articles in this section visit <http://onlinelibrary.wiley.com/doi/10.1111/bph.v175.12/issuetoc>

Abbreviations

DRG, dorsal root ganglion; HVA, high voltage-activated; LVA, low voltage-activated; TSP4, thrombospondin-4; VGCC, voltage-gated calcium channels

Introduction

Under physiological conditions, nociceptive sensations primarily serve as a protective mechanism for our body. A stimulus detected by sensory nerve endings evokes action potentials that propagate along the primary afferent nerve fibres up to the soma of dorsal root ganglion (DRG) neurons and then through their central axons to the first relay in the dorsal spinal cord. As a sensory information processing hub referring periphery sensory information to the brain, the spinal dorsal horn sensory circuitry is critically involved in processing sensory information locally and sending it through ascending tracts to the higher CNS for further processing. The brain then sends descending signals to the spinal cord through descending pathways to modulate sensory and motor neuron circuitry activities. However, under pathological conditions, damage to the nervous system can induce plasticity changes in the sensory pathway so that its activation thresholds are lowered. As a consequence, exaggerated pain perception can occur in the absence of painful stimuli or with mild painful stimulation. This kind of abnormal sensation gives rise to painful phenomena including 'allodynia' (i.e. painful perception to an innocuous stimulus) and 'hyperalgesia' (exacerbated pain perception to a mildly noxious stimulus), collectively called pain states here. Among the factors that play critical roles in pain development after nerve injuries (neuropathic pain states), voltage-gated calcium channels (VGCCs) contribute to this process through the modulation of important pathophysiological functions. These include the release of excitatory neurotransmitters (Lee, 2013), calcium-dependent enzyme activation (Park and Luo, 2010), effects on gene expression (Perret and Luo, 2009; Park and Luo, 2010; Wheeler *et al.*, 2012) and other short- and long-term plasticity changes in the spinal dorsal horn (Russo and Hounsgaard, 1994; Naka *et al.*, 2013; Youn *et al.*, 2013; Liu and Zhou, 2015). In addition, injury-induced dysregulation of VGCC subunits, such as $\alpha_2\delta_1$ ($\text{Ca}_v\alpha_2\delta$), may also contribute to pain signal transduction by modulating additional neural plasticity such as abnormal synaptogenesis (Eroglu *et al.*, 2009; Park *et al.*, 2016). These changes can alter the activities of the neural network that fundamentally contribute to the cellular underpinning of chronic pain development.

Voltage-gated calcium channels and accessory subunits

So far, 10 VGCC subtypes have been identified that are divided broadly into two categories based on their thresholds of activation by membrane depolarization: high voltage-activated (HVA) and low voltage-activated (LVA) channels (Benarroch, 2010; Simms and Zamponi, 2014; Zamponi *et al.*, 2015). The HVA channel family can be further subdivided, based on their pharmacological and functional characteristics, into L-, N-, P/Q- and R-types, and the LVA channels are mainly T-type VGCCs. Structurally, all VGCCs contain a pore-forming α_1 subunit that determines their main biophysical properties. There are three major families of α_1 subunits (Ca_v1 , Ca_v2 and Ca_v3), which can undergo alternative splicing in a tissue-, age- and pathology-dependent

manner (Lipscombe *et al.*, 2013). In addition, the HVA channels are heteromultimers with three auxiliary subunits, $\text{Ca}_v\beta$, $\text{Ca}_v\alpha_2\delta$ and $\text{Ca}_v\gamma$. Four $\text{Ca}_v\beta$ subunits ($\text{Ca}_v\beta_{1-4}$), four $\text{Ca}_v\alpha_2\delta$ subunits ($\text{Ca}_v\alpha_2\delta_{-4}$) and eight $\text{Ca}_v\gamma$ subunits ($\text{Ca}_v\gamma_{1-8}$) have been identified. Unlike the HVA channels that require co-assembly with auxiliary calcium channel subunits to function, the LVA channels appear as α_1 subunit monomers that can be functionally regulated by the auxiliary subunits (Dubel *et al.*, 2004). However, other studies reported that auxiliary subunits of calcium channels have little or no modulatory effect on LVA channel currents (Perez-Reyes, 2003; Turner and Zamponi, 2014).

The functions and details of the $\text{Ca}_v\alpha_1$, $\text{Ca}_v\beta$ and $\text{Ca}_v\gamma$ subunits are reviewed and summarized in other chapters in this issue and recent reviews (Simms and Zamponi, 2014; Zamponi *et al.*, 2015; Campiglio and Flucher, 2015). In addition, the structural feature, regulation and function of different types of VGCCs and their implications in neurological diseases including pain, migraine, epilepsy, cerebellar ataxia, Parkinson disease and hypertension have been the subjects of numerous recent reviews (Perret and Luo, 2009; Park and Luo, 2010; Cain and Snutch, 2011; Vink and Alewood, 2012; Simms and Zamponi, 2014; Zamponi *et al.*, 2015; Zamponi, 2016). In this review, we focus on recent findings related to maladaptive and dysregulation of $\text{Ca}_v\alpha_2\delta$ and its potential contribution to pain development, as well as prospective views of blocking $\text{Ca}_v\alpha_2\delta$ in neuropathic pain development and management.

$\text{Ca}_v\alpha_2\delta$ subunits

Each $\text{Ca}_v\alpha_2\delta$ subunit consists of the $\text{Ca}_v\alpha_2$ and $\text{Ca}_v\delta$ peptides that are encoded by the same gene, post-translationally cleaved and then linked by disulfide-bonds (Ellis *et al.*, 1988). Site-directed mutagenesis data have indicated that a single intermolecular disulfide bond between cysteine residue 404 in the von Willebrand factor A (VWA) domain of $\text{Ca}_v\alpha_2$ and cysteine residue 1047 in the extracellular domain of $\text{Ca}_v\delta$ is critical for the structural and functional integrity of the $\text{Ca}_v\alpha_2\delta$ protein (Calderon-Rivera *et al.*, 2012). Recently, data from cryo-electron microscopy reveal that the primary sequence of $\text{Ca}_v\alpha_2\delta$ contains four tandem cache domains and one VWA domain that are intertwined (Wu *et al.*, 2016). In addition, there are four disulfide bonds between the $\text{Ca}_v\alpha_2$ and $\text{Ca}_v\delta$ peptides and two within the $\text{Ca}_v\delta$ peptide (Wu *et al.*, 2016). In combination with the site-directed mutagenesis data, it is likely that each of these disulfide bonds is mutually critical in stabilizing the calcium channel complex and for its functions. Data from biochemical characterization of $\text{Ca}_v\alpha_2\delta$ proteins support the view that $\text{Ca}_v\alpha_2\delta$ is attached to the extracellular leaflet of the plasma membrane *via* a glycosphosphatidylinositol anchor (Davies *et al.*, 2010), which seems to be supported by structural data revealed by cryo-electron microscopy (Wu *et al.*, 2016). The known functions of $\text{Ca}_v\alpha_2\delta$ subunits include promoting and stabilizing cell surface expression and modulating functions of VGCC (Dolphin, 2012; Dolphin, 2013).

Calcium channel $\text{Ca}_v\alpha_2\delta_1$ subunit as a therapeutic target in pain management

Pharmacological and biochemical studies

Data from biochemical studies have indicated that gabapentinoids, including **gabapentin** (Neurontin; Pfizer) and **pregabalin** (Lyrica; Pfizer), bind to both $\text{Ca}_v\alpha_2\delta_1$ and $\text{Ca}_v\alpha_2\delta_2$ subunit proteins (Gee *et al.*, 1996; Gong *et al.*, 2001; Marais *et al.*, 2001b; Li *et al.*, 2011; Kukkar *et al.*, 2013; Verma *et al.*, 2014; Landmark *et al.*, 2015). Gabapentinoids are widely used in management of neuropathic pain, including that derived from diabetic neuropathy, nerve injuries, drugs and radiation-induced neuropathies, as well as post-herpetic and trigeminal neuralgia (Johnson and Rice, 2014; Moore *et al.*, 2014; Zamponi *et al.*, 2015). However, only a significant up-regulation of $\text{Ca}_v\alpha_2\delta_1$, but not $\text{Ca}_v\alpha_2\delta_2$, has been observed in DRG/spinal cord of neuropathic pain models (Luo *et al.*, 2001; 2002; Valder *et al.*, 2003; Bauer *et al.*, 2009), supporting the concept that the anti-hyperalgesic properties of gabapentinoids are mediated by their binding to the $\text{Ca}_v\alpha_2\delta_1$ subunit. This is confirmed by point mutation of the binding site for gabapentinoids on $\text{Ca}_v\alpha_2\delta_1$ in a transgenic mouse line, which diminishes $\text{Ca}_v\alpha_2\delta_1$ binding and the anti-hyperalgesic properties of gabapentinoids (Field *et al.*, 2006).

Genetic studies

One way to confirm that $\text{Ca}_v\alpha_2\delta_1$ has a role in pain processing is to use $\text{Ca}_v\alpha_2\delta_1$ knockout mice to generate pain research models, which certainly come with limitations, or correlate

abnormal sensation deficits with $\text{Ca}_v\alpha_2\delta_1$ genetic defects in patients. Table 1 summarizes findings of the studies done in $\text{Ca}_v\alpha_2\delta$ genetic-knockout mouse models or from patients with genetic defects regarding a potential role for each $\text{Ca}_v\alpha_2\delta$ subunit in pain signal transduction. Even though $\text{Ca}_v\alpha_2\delta_1$ knockout is conceived as embryonic lethal (Simms and Zamponi, 2014), it has been reported that mice with a targeted deletion of $\text{Ca}_v\alpha_2\delta_1$ (Fuller-Bicer *et al.*, 2009) display normal sensitivity to thermal stimulation but reduced sensitivity to mechanical and cold stimuli that correlates with reduced responses of dorsal horn wide dynamic range neurons to stimuli (Fuller-Bicer *et al.*, 2009; Patel *et al.*, 2013). In addition, the onset time of mechanical hypersensitivity is delayed, and the anti-hyperalgesic activity of gabapentinoids after peripheral nerve injury is absent in this line of mice (Patel *et al.*, 2013). These findings confirm that the induction of $\text{Ca}_v\alpha_2\delta_1$ is critical for the initial development of neuropathic pain states after nerve injury (Boroujerdi *et al.*, 2008) and that $\text{Ca}_v\alpha_2\delta_1$ does indeed mediate the anti-hyperalgesic properties of gabapentinoids (Field *et al.*, 2006), as reported previously. Peripherally, general $\text{Ca}_v\alpha_2\delta_1$ knockout results in a significant reduction in calcium entry per action potential, and in the duration and frequency of action potentials in DRG sensory neurons (Margas *et al.*, 2016). However, linking a direct causal role of $\text{Ca}_v\alpha_2\delta_1$ to these electrophysiological and behavioural changes in this mouse line is difficult because we cannot rule out the contributions from other compensatory factors and/or developmental changes due to general embryonic knockout of $\text{Ca}_v\alpha_2\delta_1$. It is conceivable that general knockout of $\text{Ca}_v\alpha_2\delta_1$ can affect a wide range of

Table 1

Genetic studies regarding the role of $\text{Ca}_v\alpha_2\delta$ subunits in pain signal transduction

$\text{Ca}_v\alpha_2\delta$ isoform	Genetic modification	Sensory phenotype	Reference
$\text{Ca}_v\alpha_2\delta_1$	General (embryonic) knockout in mice	↓ Mechanical/cold sensitivity Delayed mechanical hypersensitivity onset after nerve injury	Fuller-Bicer <i>et al.</i> , 2009 Patel <i>et al.</i> , 2013
	Natural aberrations/mutations in patients	Not reported	Vergult <i>et al.</i> , 2015 Hino-Fukuyo <i>et al.</i> , 2015
$\text{Ca}_v\alpha_2\delta_2$	Spontaneous mutations in mice	Not reported	Barclay <i>et al.</i> , 2001; Brodbeck <i>et al.</i> , 2002 Donato <i>et al.</i> , 2006 Brill <i>et al.</i> , 2004
	Targeted disruption in mice Homozygous mutations in patients		Ivanov <i>et al.</i> , 2004 Edvardson <i>et al.</i> , 2013 Pippucci <i>et al.</i> , 2013
$\text{Ca}_v\alpha_2\delta_3$	Spontaneous mutations in <i>Drosophila</i>	↓ Acute heat pain sensitivity	Neely <i>et al.</i> , 2010
	Deletion by homologous recombination in mice	Delayed inflammatory thermal hyperalgesia	
	Single-nucleotide polymorphisms in patients	↓ Low-back pain sensitivity	
$\text{Ca}_v\alpha_2\delta_4$	Splice site mutation	Not reported	Iossifov <i>et al.</i> , 2012
	Spontaneous homozygous frame shift mutation in mice Homozygous nucleotide substitution in patients	Not reported	Wycisk <i>et al.</i> , 2006a Wycisk <i>et al.</i> , 2006b

important organ development and physiological functions due to the importance of $\text{Ca}_v\alpha_2\delta_1$ in VGCC trafficking, membrane expression and in modulating different functions of VGCCs (Hendrich *et al.*, 2008; Cassidy *et al.*, 2014). With the application of Cre-induced recombination technology, sensory neuron-type specific deletion of $\text{Ca}_v\alpha_2\delta_1$ (Park *et al.*, 2016) allows us to study the contribution of $\text{Ca}_v\alpha_2\delta_1$ up-regulation in a subpopulation of sensory neurons to modality-specific processing of neuropathic pain states.

Genomic aberration of the CACNA2D1 gene encoding $\text{Ca}_v\alpha_2\delta_1$ has been identified in total blood DNA of three patients with epilepsy and intellectual disability, supporting a potential contribution of $\text{Ca}_v\alpha_2\delta_1$ to these clinical phenotypes (Vergult *et al.*, 2015). Similarly, another study has identified mutations in $\text{Ca}_v\alpha_2\delta_1$ from peripheral blood DNA of patients with early-onset epileptic encephalopathy (West syndrome) (Hino-Fukuyo *et al.*, 2015). Unfortunately, abnormal behavioural sensitivity has not been reported or tested in these patients. However, $\text{Ca}_v\alpha_2\delta_1$ over-expression in mouse neuronal tissues has been reported to cause brain epileptiform activity associated with behavioural arrests (Faria *et al.*, 2017) and spinal synaptic hyperexcitability associated with behavioural hypersensitivity to mechanical/thermal/cold stimuli (Li *et al.*, 2006; Zhou and Luo, 2013; Zhou and Luo, 2015). These data support a role for $\text{Ca}_v\alpha_2\delta_1$ in mediating the neural circuitry hyperexcitability associated with both epileptic and pain phenotypes.

Since gabapentinoids bind to both $\text{Ca}_v\alpha_2\delta_1$ and $\text{Ca}_v\alpha_2\delta_2$ subunits and show antiepileptic, anxiolytic and antihyperalgesic efficacy in preclinical studies, a recent study was well-designed to distinguish the role of $\text{Ca}_v\alpha_2\delta_1$ and $\text{Ca}_v\alpha_2\delta_2$ in the anticonvulsant activity of gabapentinoids. Mutations of the binding site for gabapentinoids in $\text{Ca}_v\alpha_2\delta_1$, but not in $\text{Ca}_v\alpha_2\delta_2$, result in a significant reduction in the anticonvulsant efficacy of pregabalin compared with wild-type controls and the $\text{Ca}_v\alpha_2\delta_2$ mutants (Lotarski *et al.*, 2014). Similarly, these mouse mutants were used to reveal the association of drug binding to $\text{Ca}_v\alpha_2\delta_1$, but not $\text{Ca}_v\alpha_2\delta_2$, with the anxiolytic-like activity of gabapentinoids (Lotarski *et al.*, 2011). Together with other findings indicating a role for $\text{Ca}_v\alpha_2\delta_1$ in pain signal transduction and in mediating the antihyperalgesic activity of gabapentinoids, findings from these genetic studies support the involvement of $\text{Ca}_v\alpha_2\delta_1$ in mediating pain signal transduction, epileptic and anxiolytic activities as well as the inhibitory effects of gabapentinoids on these disorders.

Genetic studies on other $\text{Ca}_v\alpha_2\delta$ isoforms

Naturally occurring mutations in the CACNA2D2 gene encoding $\text{Ca}_v\alpha_2\delta_2$ in ducky (Barclay *et al.*, 2001; Brodbeck *et al.*, 2002), ducky^{2j} (Donato *et al.*, 2006) and entla (Brill *et al.*, 2004) mice as well as targeted $\text{Ca}_v\alpha_2\delta_2$ knockout (Ivanov *et al.*, 2004) result in reduced calcium current densities, gabapentinoid binding sites and ataxia and epileptic phenotypes. Two recent clinical studies have identified different homozygous mutations in the CACNA2D2 gene that correlate with epileptic encephalopathy (Edvardson *et al.*, 2013; Pippucci *et al.*, 2013), supporting the possibility that $\text{Ca}_v\alpha_2\delta_2$

may play a role in mediating epileptic activities. Even though abnormalities in both $\text{Ca}_v\alpha_2\delta_1$ and $\text{Ca}_v\alpha_2\delta_2$ correlate with the development of epileptic activity, gabapentinoids' anti-seizure effect is likely to be mediated through their binding to $\text{Ca}_v\alpha_2\delta_1$, but not $\text{Ca}_v\alpha_2\delta_2$, because the anti-seizure efficacy of pregabalin is reduced in mice with a gabapentinoid binding site mutation in $\text{Ca}_v\alpha_2\delta_1$, but not in $\text{Ca}_v\alpha_2\delta_2$ (Lotarski *et al.*, 2014). Further mechanistic studies regarding the relative contribution of $\text{Ca}_v\alpha_2\delta_1$ and $\text{Ca}_v\alpha_2\delta_2$ to brain neural circuitry hyperexcitability in a cell type-specific manner may provide answers to this discrepancy. Abnormal nociceptive sensitivity was not reported in these studies (Table 1).

A genome-wide screen study in drosophila has identified the CACNA2D3 gene encoding $\text{Ca}_v\alpha_2\delta_3$ as an evolutionarily conserved pain gene mediating thermal hyperalgesia (Neely *et al.*, 2010). Deletion of $\text{Ca}_v\alpha_2\delta_3$ with homologous recombination in mice results in deficits in acute heat pain sensitivity and delayed thermal hyperalgesia induced by inflammation (Neely *et al.*, 2010). In addition, single-nucleotide polymorphisms within or close to the human CACNA2D3 gene are shown to be associated with reduced thermal pain sensitivity in healthy volunteers, and with chronic low-back pain in patients (Neely *et al.*, 2010). Data from functional imaging in $\text{Ca}_v\alpha_2\delta_3$ mutant mice indicate that thermal pain signal transmission from the thalamus to higher pain centres is impaired (Neely *et al.*, 2010). Since $\text{Ca}_v\alpha_2\delta_3$ is mainly expressed in the brain, and it is not a binding site for gabapentinoids (Marais *et al.*, 2001a), these findings support a role for $\text{Ca}_v\alpha_2\delta_3$ in transmitting pain signals in the brain through a mechanism that could be insensitive to gabapentinoids. Interestingly, sensory (noxious thermal and tactile) stimulation can trigger strong cross-activation of brain regions related to hearing, vision and olfaction in $\text{Ca}_v\alpha_2\delta_3$ mutant mice (Neely *et al.*, 2010), supporting a role for $\text{Ca}_v\alpha_2\delta_3$ in maintaining normal sensory signal distinction in the brain. In addition, splice site mutation of the human CACNA2D3 gene might be involved in autism spectrum disorders (Iossifov *et al.*, 2012). However, sensory testing is not reported in these studies (Table 1).

As $\text{Ca}_v\alpha_2\delta_4$ is the major $\text{Ca}_v\alpha_2\delta$ subunit expressed in mouse retina cells (Knoflach *et al.*, 2013), mutations in the CACNA2D4 gene encoding $\text{Ca}_v\alpha_2\delta_4$ result in loss of retina signalling, abnormal synapse morphology and dysfunction in mouse rods and cones (Wycisk *et al.*, 2006a), as well as autosomal recessive cone dystrophy associated with night blindness in patients (Wycisk *et al.*, 2006b). Abnormal nociceptive sensitivity was not reported or tested in these studies (Table 1).

Potential mechanisms underlying $\text{Ca}_v\alpha_2\delta_1$ dysregulation in pain signal transduction and the anti-hyperalgesic effects of gabapentinoids

$\text{Ca}_v\alpha_2\delta_1$ expression is up-regulated in DRG (and trigeminal ganglia) and spinal cord in some neuropathic pain models derived from mechanical nerve injury, spinal cord injury, diabetic or chemical neuropathies that correlate with the onset and maintenance of neuropathic pain states (Luo *et al.*, 2001; 2002; Newton *et al.*, 2001; Yusaf *et al.*, 2001; Li *et al.*, 2004; 2014b; Boroujerdi *et al.*, 2008; 2011). The $\text{Ca}_v\alpha_2\delta_1$

ligand gabapentinoids can reverse neuropathic pain states in these animal models (Hwang and Yaksh, 1997; Luo *et al.*, 2002; Boroujerdi *et al.*, 2011; Li *et al.*, 2014b). In addition, transgenic mice overexpressing $\text{Ca}_v\alpha_2\delta_1$ in neuronal tissues, including sensory and spinal cord neurons, display neuropathic pain states such as tactile allodynia and thermal hyperalgesia, in the absence of any nerve injury, that can be blocked by gabapentinoids (Li *et al.*, 2006). Together, these findings support the view that injury-induced $\text{Ca}_v\alpha_2\delta_1$ up-regulation may play a critical role in neuropathic pain development.

Data from detailed mechanistic investigations indicate that nerve injury-induced $\text{Ca}_v\alpha_2\delta_1$ expression occurs predominantly in DRG sensory neurons, then undergoes axonal translocation to presynaptic terminals in spinal dorsal horn (Figure 1), since the axonal transport can be interrupted by cutting the dorsal root connecting injured DRG and dorsal spinal cord through dorsal rhizotomy (Li *et al.*, 2004). Subsequently, this axonal transport of an elevated expression of $\text{Ca}_v\alpha_2\delta_1$ from DRG neurons to dorsal spinal cord was independently confirmed. Data from electron microscopy show directly that injury-induced $\text{Ca}_v\alpha_2\delta_1$ in DRG is indeed trafficking to pre-synaptic terminals of sensory fibres in dorsal spinal cord, and this process can be blocked by gabapentinoids (prebatalin) (Bauer *et al.*, 2009). Thus, injury-induced increased expression of $\text{Ca}_v\alpha_2\delta_1$ probably regulates sensory pathway sensitivity peripherally and centrally.

At both the peripheral and central levels, it is possible that different $\text{Ca}_v\alpha_2\delta_1$ isoforms play distinct roles in mediating pain states. Early reports have indicated that peripheral nerve injury induces the up-regulation of distinct forms of $\text{Ca}_v\alpha_2\delta_1$ in DRG and spinal cord that correlates with the development of neuropathic pain states (Luo *et al.*, 2001). In addition, injury-induced DRG $\text{Ca}_v\alpha_2\delta_1$ isoforms differ from that detected in other tissues such as spinal cord, brain and skeletal muscle as they have unique migration rates in Western blots and different glycosylation patterns (Luo, 2000; Luo *et al.*, 2001). These findings suggest that $\text{Ca}_v\alpha_2\delta_1$ may have tissue-specific functions and injury-induced $\text{Ca}_v\alpha_2\delta_1$ up-regulation may modulate pain processing in a tissue type-specific manner. Recently, it was reported that DRG $\text{Ca}_v\alpha_2\delta_1$ isoforms are indeed derived from alternative splicing (Lana *et al.*, 2014). In addition, a minor splice variant ($\Delta A + B\Delta C$) of $\text{Ca}_v\alpha_2\delta_1$ is preferentially up-regulated in small DRG neurons after spinal nerve ligation injury, which also enhances calcium currents as it is the major splice variant of DRG $\text{Ca}_v\alpha_2\delta_1$, but has significantly reduced affinity for gabapentin (Lana *et al.*, 2014). It is speculated that differential expression of specifically spliced $\text{Ca}_v\alpha_2\delta_1$ variants in neuropathic pain patients may account for the variations in efficacy of pain relief by gabapentinoids observed clinically (Lana *et al.*, 2014).

At the peripheral level, dysregulated $\text{Ca}_v\alpha_2\delta_1$ can regulate DRG neuron activities through the modulation of calcium channel activities (Li *et al.*, 2006; D'Arco *et al.*, 2015). At the spinal cord level, it can increase presynaptic excitatory input into superficial and deep dorsal horn that regulates dorsal horn neuron excitabilities (Nguyen *et al.*, 2009; Zhou and Luo, 2013, 2015). Since different VGCC blockers have distinct inhibitory effects on $\text{Ca}_v\alpha_2\delta_1$ -mediated nociception, it is likely that $\text{Ca}_v\alpha_2\delta_1$ dysregulation contributes to pain

processing at least partially through modulation of pathways associated with specific VGCCs (Chang *et al.*, 2015).

Data from *in vitro* studies indicate that $\text{Ca}_v\alpha_2\delta_1$ can increase cell-surface and synaptic expression of VGCCs, including the P/Q-type ($\text{Ca}_v2.1$) (Hendrich *et al.*, 2008; Hoppa *et al.*, 2012) and N-type ($\text{Ca}_v2.2$) (Cassidy *et al.*, 2014) calcium channels. Calcium entry through these channels predominantly drives synaptic transmission and initiates fast release of classical neurotransmitters (Catterall, 2011; Campiglio and Flucher, 2015). Mechanistically, it has been shown that increased transient $\text{Ca}_v\alpha_2\delta_1$ expression in cultured rat DRG neurons leads to increased membrane expression of N-type ($\text{Ca}_v2.2$) VGCC. This plasticity couples to a prolonged Ca^{2+} signal evoked by membrane depolarization and activity-dependent slowing of axonal mitochondrial transport, both of which can be inhibited by treatments with N-type VGCC blockers. These findings suggest that increased $\text{Ca}_v\alpha_2\delta_1$ expression modulates sensory neuron responses to stimulation through the regulation of mitochondrial calcium buffering, which is mediated by N-type VGCCs (D'Arco *et al.*, 2015). Interestingly, chronic treatment with gabapentinoids can reduce the densities of $\text{Ca}_v\alpha_2\delta_1$ and Ca_v2 ($\text{Ca}_v2.1$ and $\text{Ca}_v2.2$) subunits in plasma membrane and inhibit excitatory presynaptic transmission (Hendrich *et al.*, 2008, 2012; Bauer *et al.*, 2009, 2010; Cassidy *et al.*, 2014), presumably by interfering with $\text{Ca}_v\alpha_2\delta_1$ trafficking process through disrupting the recycling of $\text{Ca}_v\alpha_2\delta_1$ to the plasma membrane (Tran-Van-Minh and Dolphin, 2010). This is supported by *in vitro* findings that the membrane surface expression of $\text{Ca}_v2.2$ channels is reduced and become insensitive to gabapentin blockade when co-expressed with a $\text{Ca}_v\alpha_2\delta_1$ mutant lacking the binding site for gabapentinoids (Cassidy *et al.*, 2014). $\text{Ca}_v\alpha_2\delta_1$ can also enhance the expression of T-type calcium channels in plasma membrane without changing their electrophysiological properties (Dubel *et al.*, 2004). However, $\text{Ca}_v\alpha_2\delta_1$ has also been found to have little or no modulatory effect on T-type channel currents (Perez-Reyes, 2003; Turner and Zamponi, 2014), and gabapentin failed to affect VGCC trafficking or function in cultured hippocampal neurons (Brown and Randall, 2005; Hoppa *et al.*, 2012).

Recent data from biochemical studies suggest that an elevation in $\text{Ca}_v\alpha_2\delta_1$ promotes presynaptic neurotransmitter release by modulating presynaptic VGCC abundance and Ca^{2+} influx-driven exocytosis. While the former can be explained by the ability of $\text{Ca}_v\alpha_2\delta_1$ to promote VGCC trafficking, the latter is believed to be mediated through an extracellular metal ion-dependent adhesion site within the VWA domain of $\text{Ca}_v\alpha_2\delta_1$ (Hoppa *et al.*, 2012; Korber and Kuner, 2016). These findings provide mechanistic insights, revealing how $\text{Ca}_v\alpha_2\delta_1$ plays a significant pathological role in modulating the release of numerous neurotransmitters, including pain-inducing excitatory neurotransmitters and peptides at the spinal cord level. This can be one of the primary targets of gabapentinoids in pain modulation.

While alterations in VGCC trafficking may lead to changes in VGCC functions, and subsequently changes in sensory neuron excitability or presynaptic neurotransmitter release, it is unable to explain the fast actions of gabapentinoids in reversing pain states in animal models, which occur within 1 h of administration (Hwang and Yaksh, 1997; Luo *et al.*, 2002). Recently, it was reported that $\text{Ca}_v\alpha_2\delta_3$




Figure 1

Cartoon illustration of injury-induced peripheral and central plasticity contributing to sensory sensitization and pain states. Nerve injury induces an up-regulation of $Ca_v\alpha_2\delta_1$ in DRG that undergoes axonal transport to the presynaptic terminal of injured sensory fibres in the dorsal spinal cord. Nerve injury also leads to increased TSP4 expression and secretion from activated DRG neurons/non-neuronal cells and spinal cord astrocytes. Peripherally, the elevated TSP4/ $Ca_v\alpha_2\delta_1$ can interact to modulate VGCC currents and intracellular calcium levels, which may cause sensory neuron/afferent activation and sensitization of presynaptic terminals. Centrally, elevated TSP4/ $Ca_v\alpha_2\delta_1$ can interact to promote presynaptic excitatory neurotransmitter release and excitatory synapse formation, which together can lead to activation of postsynaptic dorsal horn neurons and central sensitization.

is required for the rapid induction and sustained maintenance of presynaptic homeostatic potentiation, a form of neuroplasticity in which the presynaptic neurotransmitter release apparatus is modulated in response to postsynaptic perturbations that may occur in neurological disorders

(Wang *et al.*, 2016). It appears that extracellular $Ca_v\alpha_2\delta_3$ residing in the presynaptic release sites of the synaptic cleft serves as a transsynaptic homeostatic signalling molecule that relays retrograde information to presynaptic cytoplasm to modulate active zone activities, presumably through

interactions with other proteins. This plasticity seems to be independent of its ability to regulate VGCC abundance (Wang *et al.*, 2016). If this plasticity and its sensitivity to blockade by gabapentinoids are confirmed in neuropathic pain conditions, it may underlie a new mechanism through which gabapentinoids reverse neuropathic pain states so promptly. Identification of $\text{Ca}_v\alpha_2\delta_3$ as a pain gene (Neely *et al.*, 2010) (Table 1) supports this notion. However, there is no evidence so far indicating that $\text{Ca}_v\alpha_2\delta_3$ is a binding site for gabapentinoids, or is up-regulated after nerve injuries. In fact, $\text{Ca}_v\alpha_2\delta_3$ mRNA is significantly reduced in injured DRG after nerve injury (Bauer *et al.*, 2009). More detail studies may help to distinguish the relative contribution of $\text{Ca}_v\alpha_2\delta_1$ and $\text{Ca}_v\alpha_2\delta_3$ to pain signal transduction and the anti-hyperalgesic actions of gabapentinoids.

In addition, it has been reported that gabapentin can block the interactions between $\text{Ca}_v\alpha_2\delta_1$ and synaptogenic thrombospondin-4 (TSP4) (Park *et al.*, 2016), which is also up-regulated in DRG and spinal cord after peripheral nerve injury (Kim *et al.*, 2012; Pan *et al.*, 2016). This leads to diminished dorsal horn neuron hypersensitivity and behavioural hypersensitivity within 1 h (Park *et al.*, 2016). This fast action of gabapentin may provide another possible explanation about the fast anti-hyperalgesic actions of gabapentinoids.

Current states of gabapentinoids in the management and prevention of neuropathic pain states

Gabapentinoids provide pain relief mainly in a pathophysiological state-dependent manner; they are only effective in alleviating pathological pain sensations but do not affect acute physiological noxious sensations (Field *et al.*, 1997b; Stanfa *et al.*, 1997; Dickenson *et al.*, 2005). These findings suggest that gabapentinoids most likely work on a sensitized sensory pathway related to $\text{Ca}_v\alpha_2\delta_1$ up-regulation. This may explain why gabapentinoids only provide significant pain relief in about one third of neuropathic pain patients (Moore *et al.*, 2014), since it is conceivable that pathological changes induced by $\text{Ca}_v\alpha_2\delta_1$ up-regulation may not be present in all neuropathic pain patients non-selectively. This is supported by data from animal model studies indicating that gabapentin is only effective in reversing neuropathic pain states associated with detectable $\text{Ca}_v\alpha_2\delta_1$ up-regulation in dorsal spinal cord and/or DRG (Luo *et al.*, 2002). Gabapentinoids have relatively mild side effects, which include dizziness, somnolence and ataxia. The latter is believed to be mediated through an effect on calcium channel trafficking (Jun *et al.*, 1999; Mark *et al.*, 2011) and by binding to $\text{Ca}_v\alpha_2\delta_2$ (Barclay *et al.*, 2001; Ivanov *et al.*, 2004).

Based on evidence from randomized clinical trials (RCTs), expert panels from the Neuropathic Pain Special Interest Group of the International Association for the Study of Pain (Dworkin *et al.*, 2007), European Federation of Neurological Societies (Attal *et al.*, 2006) and Canadian Pain Society (Moulin *et al.*, 2007) recommended, almost simultaneously, that gabapentinoids should be one of the first line medication classes for neuropathic pain management in 2006–2007. Recent updated reviews for using gabapentinoids in neuropathic pain management are summarized in Table 2.

Depending on the aetiology of a neuropathic pain condition, the conclusions from these reviews still confirm that gabapentinoids can be used as first-line medications or are recommended for the management of neuropathic pain derived from certain aetiologies (Dworkin *et al.*, 2010; Vargas-Espinosa *et al.*, 2012; Hershman *et al.*, 2014; Finnerup *et al.*, 2015; Loh *et al.*, 2016). Findings from these reviews also point out the need for more studies for dose-standardization, to assess their adverse effects and their suitability for neuropathic pain derived from other aetiologies such as cancer, chemotherapy-induced peripheral neuropathies, and combination therapies, so that better drug efficacy can be achieved and/or their side effects reduced (Dworkin *et al.*, 2010; Chaparro *et al.*, 2012; Vargas-Espinosa *et al.*, 2012; Hershman *et al.*, 2014; Guan *et al.*, 2016).

Even though most animal model studies confirm that gabapentinoids are effective at reversing pain states, clinical data for the efficacy of gabapentinoids in acute and persistent pain relief are not consistent. While some studies show that gabapentinoids are effective in pain relief after surgeries, this comes at the expense of increased adverse effects (Buvanendran *et al.*, 2010; Engelman and Cateloy, 2011; Clarke *et al.*, 2012; Eipe *et al.*, 2015; Mishriky *et al.*, 2015). Other studies do not show similar benefits of gabapentinoid treatment (Lunn *et al.*, 2015; Kharasch and Eisenach, 2016; Martinez *et al.*, 2017a,b). This could be due to the fact that for various reasons negative data from animal model studies are often not getting published, but that is less likely to be the case for most clinical trials. These discrepancies in clinical studies has led to the suggestion that more studies in different surgical models and specific patient populations are needed to explore the benefits of a surgical model-specific application of gabapentinoids (Fletcher and Martinez, 2015).

One way to optimize the pharmacological benefits and reduce the adverse effects of gabapentinoids in pain management is to lower the dose of gabapentinoids by using them as a combination treatment. A few studies have addressed this option with inconsistent outcomes, as summarized in the reviews presented in Table 2. Alternatively, pre-emptive treatment with gabapentinoids has been shown to be efficacious in blocking post-operative pain development in patients (Field *et al.*, 1997a; Clivatti *et al.*, 2009; Mardani-Kivi *et al.*, 2013; Ravindran, 2014; Hwang *et al.*, 2015). However, few studies have addressed the underlying mechanism of the preventive effects of gabapentinoids in chronic pain development. In another study reported in this issue, we tested the hypothesis that early low-dose gabapentin treatment can prevent injury-induced neuropathic pain development by inhibiting abnormal excitatory synaptogenesis between sensory and spinal cord neurons (Yu *et al.*, 2017).

Synaptogenic effects of $\text{Ca}_v\alpha_2\delta_1$ /TSP4 in chronic pain processing

At the dorsal spinal cord level, the interactions of the up-regulated $\text{Ca}_v\alpha_2\delta_1$ with TSP4 post injury can lead to aberrant excitatory synaptogenesis that is highly likely to contribute to the development of chronic neuropathic pain, as illustrated in Figure 1 and is supported by the following findings from animal studies. Peripheral nerve injury induces an up-

Table 2

Recommendation of gabapentinoids for neuropathic pain management

Review criteria/strength	Neuropathic pain-inducing condition	Recommendation	Reference
Randomized clinical trials (RCTs)/systemic reviews/pooled analysis/retrospective study	Post-herpetic neuralgia	Type A recommendation from consistent, good quality patient-oriented evidence More studies are needed for standardizing doses and assessing adverse effects	Vargas-Espinosa <i>et al.</i> , 2012
RCTs \geq 1 week, combination pharmacotherapy, neuropathic pain in cancer patients	Cancer	Further studies are needed	Guan <i>et al.</i> , 2016
Double-blind, RCTs comparing two or more drug combinations to placebo and/or \geq 1 other comparator	Varies	More studies are needed for two-drug combinations including comparisons with placebo and both single-agent components in neuropathic pain treatments	Chaparro <i>et al.</i> , 2012
Review established guidelines (Dworkin <i>et al.</i> , Pain 2007) and resent RCT studies for development of future recommendations	Varies	<u>Single medication therapy</u> : First-line medication <u>Combination therapy</u> : additional studies are needed to develop specific combination therapies.	Dworkin <i>et al.</i> , 2010
Recent RCTs	Chemotherapy-induced peripheral neuropathies (CIPN)	Reasonable to try for selective CIPN pain patients who should be informed about limited scientific evidence, potential benefits, harms and costs.	Hershman <i>et al.</i> , 2014
Expert opinion on clinical experience, side effect profile, effectiveness in other neuropathic pain conditions, other relevant factors	Spinal cord injury	First line therapy	Loh <i>et al.</i> , 2016
Update of evidence-based recommendations/systemic review/double-blind RCTs, including unpublished clinical trials, and meta-analysis used Numbers needed to treat for 50% pain relief and assessed publication bias	Varies	First line therapy	Finnerup <i>et al.</i> , 2015

regulation of both $Ca_v\alpha_2\delta_1$ and TSP4 in the DRG and spinal cord that precedes the onset and correlates with the duration of neuropathic pain states (Luo *et al.*, 2001; 2002; Newton *et al.*, 2001; Li *et al.*, 2004, 2014a; Kim *et al.*, 2012; Pan *et al.*, 2015). Blocking injury-induced $Ca_v\alpha_2\delta_1$ up-regulation (Boroujerdi *et al.*, 2008; Park *et al.*, 2016) or genetic ablation of TSP4 (Kim *et al.*, 2012) prevents the onset and development of injury-induced pain states. Activation of a $Ca_v\alpha_2\delta_1$ /TSP4 dependent pathway at the spinal cord level leads to sensitization of spinal dorsal horn neurons through enhanced presynaptic excitatory neurotransmission and aberrant excitatory synapse formation that are known to contribute to neuropathic pain development after peripheral nerve injuries (Zhou and Luo, 2013; 2015; Park *et al.*, 2016). Treatments that block injury-induced pain states, such as genetic ablation of either $Ca_v\alpha_2\delta_1$ or TSP4 alone, also block injury-induced aberrant excitatory synaptogenesis (Park *et al.*, 2016). Furthermore, nerve injury-induced TSP4 seems to affect VGCC subtypes in sensory neurons differently, resulting in

decreased HVA and increased LVA calcium currents, by interacting with $Ca_v\alpha_2\delta_1$ (Pan *et al.*, 2016). Together, these findings strongly support a critical role for injury-induced $Ca_v\alpha_2\delta_1$ /TSP4 in chronic neuropathic pain processing, mediated through the promotion of aberrant excitatory synapse formation and presynaptic neurotransmission in the dorsal spinal cord. The exact location of $Ca_v\alpha_2\delta_1$ /TSP4 interactions *in vivo* remains to be established since data from a recent *in vitro* co-transfection study suggest that $Ca_v\alpha_2\delta_1$ /TSP4 interactions occur intracellularly (Lana *et al.*, 2016).

New prospects of targeting $Ca_v\alpha_2\delta_1$ for neuropathic pain treatment and prevention

The translational value of these findings suggest that blocking injury-induced synapse formation may lead to the prevention or reversal of neuropathic pain states post-nerve

injuries. While genetic ablation or biochemical knockdown of $\text{Ca}_v\alpha_2\delta_1$ or TSP4 are powerful tools in mechanistic research with animal models, these approaches are not practical currently for preventing/treating neuropathic pain states in patients. Blocking injury-induced abnormal synapse formation by small molecule drugs can be an attractive approach in preventing/managing neuropathic pain development after nerve injuries. Since gabapentinoids can block the $\text{Ca}_v\alpha_2\delta_1$ /TSP4-induced aberrant excitatory synaptogenesis and development of pain states (Park *et al.*, 2016), using these drugs to block abnormal synapse formation could be a novel approach for chronic neuropathic pain management. Unfortunately, recent *in vitro* data indicate that late treatments, starting after the initiation of synapse formation, with gabapentinoids cannot reverse already formed excitatory synapses (Eroglu *et al.*, 2009; Yu *et al.*, 2017). Thus, the acute effects of gabapentinoids in blocking neuropathic pain states are unlikely to be due to reversing abnormal synapse formation.

Can early gabapentinoid treatment prevent abnormal synapse formation and pain state development? To address this question, we have tested the effects of gabapentin in synaptogenesis between cultured DRG sensory neurons and spinal cord neurons *in vitro* and in TSP4 injected mice *in vivo* (Yu *et al.*, 2017). Our data indicate that pre-emptive gabapentin treatment at a low-dose can prevent TSP4-induced excitatory synapse formation *in vitro*. However, delayed gabapentin treatment for 2 days after TSP4 addition, when excitatory synapses have been formed, has no anti-synaptogenic effects. These findings were confirmed *in vivo* by analysing synapse formation in the dorsal spinal cord of mice 4 days after bolus intrathecal TSP4 injection, which is known to cause pain-like behavioural hypersensitivity (Kim *et al.*, 2012; Park *et al.*, 2016). Early daily treatment with a sub-anti-hyperalgesic dose of gabapentin blocks TSP4-induced behavioural hypersensitivity and *in vivo* excitatory synapse formation. However, delayed gabapentin treatment starting 2 days later is not effective in blocking TSP4-induced pain states and excitatory synapse formation *in vivo* (Yu *et al.*, 2017). These findings confirm that early blocking of the activation of the TSP4/ $\text{Ca}_v\alpha_2\delta_1$ -dependent pathway by a low dose of gabapentin may prevent aberrant excitatory synapse formation in the spinal cord resulting from TSP4/ $\text{Ca}_v\alpha_2\delta_1$ up-regulation, such as in the case post-peripheral nerve injury. Clinical validation of these results may lead to this approach being used to prevent the development of pain states.

Conclusions

Maladaptive changes in the expression and functions of the VGCC $\text{Ca}_v\alpha_2\delta_1$ subunit post injuries in the peripheral and central nervous systems can contribute to chronic pain development through pathological changes associated with or independent of VGCC functions. Gabapentinoids, by acting as $\text{Ca}_v\alpha_2\delta_1$ ligands, can reverse pain states significantly in a subpopulation of neuropathic pain patients through the various potential mechanisms discussed, some of which remain to be validated. Emerging data suggest that early preventive treatment with gabapentinoids can prevent aberrant

excitatory synapse formation and pain state development, which, if confirmed clinically, could be an attractive approach for preventing neuropathic pain development.

Nomenclature of targets and ligands

Key protein targets and ligands in this article are hyperlinked to corresponding entries in <http://www.guidetopharmacology.org>, the common portal for data from the IUPHAR/BPS Guide to PHARMACOLOGY (Southan *et al.*, 2016), and are permanently archived in the Concise Guide to PHARMACOLOGY 2015/16 (Alexander *et al.*, 2015).

Acknowledgements

This work was supported in part by NIH grants NS064341 and DE021847 to Z.D.L.

Conflict of interest

The authors declare no conflicts of interest.

References

- Alexander SPH, Catterall WA, Kelly E, Marrion N, Peters JA, Benson HE *et al.* (2015). The Concise Guide to PHARMACOLOGY 2015/16: Voltage-gated ion channels. *Br J Pharmacol* 172: 5904–5941.
- Attal N, Cruccu G, Haanpaa M, Hansson P, Jensen TS, Nurmikko T *et al.* (2006). EFNS guidelines on pharmacological treatment of neuropathic pain. *Eur J Neurol* 13: 1153–1169.
- Barclay J, Balaguero N, Mione M, Ackerman SL, Letts VA, Brodbeck J *et al.* (2001). Ducky mouse phenotype of epilepsy and ataxia is associated with mutations in the *Cacna2d2* gene and decreased calcium channel current in cerebellar Purkinje cells. *J Neurosci* 21: 6095–6104.
- Bauer CS, Nieto-Rostro M, Rahman W, Tran-Van-Minh A, Ferron L, Douglas L *et al.* (2009). The increased trafficking of the calcium channel subunit $\alpha_2\delta_1$ to presynaptic terminals in neuropathic pain is inhibited by the $\alpha_2\delta_1$ ligand pregabalin. *J Neurosci* 29: 4076–4088.
- Bauer CS, Rahman W, Tran-van-Minh A, Lujan R, Dickenson AH, Dolphin AC (2010). The anti-allodynic $\alpha_2\delta_1$ ligand pregabalin inhibits the trafficking of the calcium channel $\alpha_2\delta_1$ subunit to presynaptic terminals *in vivo*. *Biochem Soc Trans* 38: 525–528.
- Benarroch EE (2010). Neuronal voltage-gated calcium channels: brief overview of their function and clinical implications in neurology. *Neurology* 74: 1310–1315.
- Boroujerdi A, Kim HK, Lyu YS, Kim DS, Figueroa KW, Chung JM *et al.* (2008). Injury discharges regulate calcium channel $\alpha_2\delta_1$ subunit upregulation in the dorsal horn that contributes to initiation of neuropathic pain. *Pain* 139: 358–366.
- Boroujerdi A, Zeng J, Sharp K, Kim D, Steward O, Luo ZD (2011). Calcium channel $\alpha_2\delta_1$ protein upregulation in dorsal spinal cord mediates spinal cord injury-induced neuropathic pain states. *Pain* 152: 649–655.

- Brill J, Klocke R, Paul D, Boison D, Gouder N, Klugbauer N *et al.* (2004). *entla*, a novel epileptic and ataxic *Cacna2d2* mutant of the mouse. *J Biol Chem* 279: 7322–7330.
- Brodbeck J, Davies A, Courtney JM, Meir A, Balaguero N, Canti C *et al.* (2002). The ducky mutation in *Cacna2d2* results in altered Purkinje cell morphology and is associated with the expression of a truncated alpha 2 delta-2 protein with abnormal function. *J Biol Chem* 277: 7684–7693.
- Brown JT, Randall A (2005). Gabapentin fails to alter P/Q-type Ca²⁺-channel-mediated synaptic transmission in the hippocampus in vitro. *Synapse* 55: 262–269.
- Buvanendran A, Kroin JS, Della Valle CJ, Kari M, Moric M, Tuman KJ (2010). Perioperative oral pregabalin reduces chronic pain after total knee arthroplasty: a prospective, randomized, controlled trial. *Anesth Analg* 110: 199–207.
- Cain SM, Snutch TP (2011). Voltage-gated calcium channels and disease. *Biofactors* 37: 197–205.
- Calderon-Rivera A, Andrade A, Hernandez-Hernandez O, Gonzalez-Ramirez R, Sandoval A, Rivera M *et al.* (2012). Identification of a disulfide bridge essential for structure and function of the voltage-gated Ca(2+) channel alpha(2)delta-1 auxiliary subunit. *Cell Calcium* 51: 22–30.
- Campiglio M, Flucher BE (2015). The role of auxiliary subunits for the functional diversity of voltage-gated calcium channels. *J Cell Physiol* 230: 2019–2031.
- Cassidy JS, Ferron L, Kadurin I, Pratt WS, Dolphin AC (2014). Functional exofacially tagged N-type calcium channels elucidate the interaction with auxiliary $\alpha 2\delta$ -1 subunits. *Proc Natl Acad Sci U S A* 111: 8979–8984.
- Catterall WA (2011). Voltage-gated calcium channels. *Cold Spring Harb Perspect Biol* 3: a003947.
- Chang E, Chen X, Kim M, Gong N, Bhatia S, Luo ZD (2015). Differential effects of voltage-gated calcium channel blockers on calcium channel alpha-2-delta-1 subunit protein-mediated nociception. *Eur J Pain* 19: 639–648.
- Chaparro LE, Wiffen PJ, Moore RA, Gilron I (2012). Combination pharmacotherapy for the treatment of neuropathic pain in adults. *Cochrane Database Syst Rev* 7: CD008943.
- Clarke H, Bonin RP, Orser BA, Englesakis M, Wijeyesundera DN, Katz J (2012). The prevention of chronic postsurgical pain using gabapentin and pregabalin: a combined systematic review and meta-analysis. *Anesth Analg* 115: 428–442.
- Clivatti J, Sakata RK, Issy AM (2009). Review of the use of gabapentin in the control of postoperative pain. *Rev Bras Anesthesiol* 59: 87–98.
- D'Arco M, Margas W, Cassidy JS, Dolphin AC (2015). The upregulation of alpha2delta-1 subunit modulates activity-dependent Ca²⁺ signals in sensory neurons. *J Neurosci* 35: 5891–5903.
- Davies A, Kadurin I, Alvarez-Laviada A, Douglas L, Nieto-Rostro M, Bauer CS *et al.* (2010). The alpha2delta subunits of voltage-gated calcium channels form GPI-anchored proteins, a posttranslational modification essential for function. *Proc Natl Acad Sci U S A* 107: 1654–1659.
- Dickenson AH, Bee LA, Suzuki R (2005). Pains, gains, and midbrains. *Proc Natl Acad Sci U S A* 102: 17885–17886.
- Dolphin AC (2012). Calcium channel auxiliary alpha2delta and beta subunits: trafficking and one step beyond. *Nat Rev Neurosci* 13: 542–555.
- Dolphin AC (2013). The alpha2delta subunits of voltage-gated calcium channels. *Biochim Biophys Acta* 1828: 1541–1549.
- Donato R, Page KM, Koch D, Nieto-Rostro M, Foucault I, Davies A *et al.* (2006). The ducky(2j) mutation in *Cacna2d2* results in reduced spontaneous Purkinje cell activity and altered gene expression. *J Neurosci* 26: 12576–12586.
- Dubel SJ, Altier C, Chaumont S, Lory P, Bourinet E, Nargeot J (2004). Plasma membrane expression of T-type calcium channel alpha(1) subunits is modulated by high voltage-activated auxiliary subunits. *J Biol Chem* 279: 29263–29269.
- Dworkin RH, O'Connor AB, Audette J, Baron R, Gourlay GK, Haanpaa ML *et al.* (2010). Recommendations for the pharmacological management of neuropathic pain: an overview and literature update. *Mayo Clin Proc* 85: S3–14.
- Dworkin RH, O'Connor AB, Backonja M, Farrar JT, Finnerup NB, Jensen TS *et al.* (2007). Pharmacologic management of neuropathic pain: evidence-based recommendations. *Pain* 132: 237–251.
- Edvardson S, Oz S, Abulhijaa FA, Taher FB, Shaag A, Zenvirt S *et al.* (2013). Early infantile epileptic encephalopathy associated with a high voltage gated calcium channelopathy. *J Med Genet* 50: 118–123.
- Eipe N, Penning J, Yazdi F, Mallick R, Turner L, Ahmadzai N *et al.* (2015). Perioperative use of pregabalin for acute pain—a systematic review and meta-analysis. *Pain* 156: 1284–1300.
- Ellis SB, Williams ME, Ways NR, Brenner R, Sharp AH, Leung AT *et al.* (1988). Sequence and expression of mRNAs encoding the alpha 1 and alpha 2 subunits of a DHP-sensitive calcium channel. *Science* 241: 1661–1664.
- Engelman E, Cateloy F (2011). Efficacy and safety of perioperative pregabalin for post-operative pain: a meta-analysis of randomized-controlled trials. *Acta Anaesthesiol Scand* 55: 927–943.
- Eroglu C, Allen NJ, Susman MW, O'Rourke NA, Park CY, Ozkan E *et al.* (2009). Gabapentin receptor alpha2delta-1 is a neuronal thrombospondin receptor responsible for excitatory CNS synaptogenesis. *Cell* 139: 380–392.
- Faria LC, Gu F, Parada I, Barres B, Luo ZD, Prince DA (2017). Epileptiform activity and behavioral arrests in mice overexpressing the calcium channel subunit alpha2delta-1. *Neurobiol Dis* 102: 70–80.
- Field MJ, Cox PJ, Stott E, Melrose H, Offord J, Su TZ *et al.* (2006). Identification of the {alpha}2-{delta}1 subunit of voltage-dependent calcium channels as a molecular target for pain mediating the analgesic actions of pregabalin. *Proc Natl Acad Sci USA* 103: 17537–17542 Epub 12006 Nov 17536.
- Field MJ, Holloman EF, McCleary S, Hughes J, Singh L (1997a). Evaluation of gabapentin and S-(+)-3-isobutylgaba in a rat model of postoperative pain. *J Pharmacol Exp Ther* 282: 1242–1246.
- Field MJ, Oles RJ, Lewis AS, McCleary S, Hughes J, Singh L (1997b). Gabapentin (neurontin) and S-(+)-3-isobutylgaba represent a novel class of selective antihyperalgesic agents. *Brit J Pharmacol* 121: 1513–1522.
- Finnerup NB, Attal N, Haroutounian S, McNicol E, Baron R, Dworkin RH *et al.* (2015). Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis. *Lancet Neurol* 14: 162–173.
- Fletcher D, Martinez V (2015). Should we use gabapentin for postoperative pain control? *Pain* 156: 2402–2403.
- Fuller-Bicer GA, Varadi G, Koch SE, Ishii M, Bodi I, Kadeer N *et al.* (2009). Targeted disruption of the voltage-dependent calcium channel alpha2/delta-1-subunit. *Am J Physiol Heart Circ Physiol* 297: H117–H124.

- Gee NS, Brown JP, Dissanayake VU, Offord J, Thurlow R, Woodruff GN (1996). The novel anticonvulsant drug, gabapentin (neurontin), binds to the alpha2delta subunit of a calcium channel. *J Biol Chem* 271: 5768–5776.
- Gong HC, Hang J, Kohler W, Li L, Su TZ (2001). Tissue-specific expression and gabapentin-binding properties of calcium channel alpha2delta subunit subtypes. *J Membr Biol* 184: 35–43.
- Guan J, Tanaka S, Kawakami K (2016). Anticonvulsants or antidepressants in combination pharmacotherapy for treatment of neuropathic pain in cancer patients: a systematic review and meta-analysis. *Clin J Pain* 32: 719–725.
- Hendrich J, Bauer CS, Dolphin AC (2012). Chronic pregabalin inhibits synaptic transmission between rat dorsal root ganglion and dorsal horn neurons in culture. *Channels (Austin)* 6: 124–132.
- Hendrich J, Van Minh AT, Heblich F, Nieto-Rostro M, Watschinger K, Striessnig J *et al.* (2008). Pharmacological disruption of calcium channel trafficking by the alpha2delta ligand gabapentin. *Proc Natl Acad Sci U S A* 105: 3628–3633.
- Hershman DL, Lacchetti C, Dworkin RH, Lavoie Smith EM, Bleeker J, Cavaletti G *et al.* (2014). Prevention and management of chemotherapy-induced peripheral neuropathy in survivors of adult cancers: American Society of Clinical Oncology clinical practice guideline. *J Clin Oncol* 32: 1941–1967.
- Hino-Fukuyo N, Kikuchi A, Arai-Ichinoi N, Niihori T, Sato R, Suzuki T *et al.* (2015). Genomic analysis identifies candidate pathogenic variants in 9 of 18 patients with unexplained West syndrome. *Hum Genet* 134: 649–658.
- Hoppa MB, Lana B, Margas W, Dolphin AC, Ryan TA (2012). alpha2delta expression sets presynaptic calcium channel abundance and release probability. *Nature* 486: 122–125.
- Hwang JH, Yaksh TL (1997). Effect of subarachnoid gabapentin on tactile-evoked allodynia in a surgically induced neuropathic pain model in the rat. *Reg Anesth* 22: 249–256.
- Hwang SH, Park IJ, Cho YJ, Jeong YM, Kang JM (2015). The efficacy of gabapentin/pregabalin in improving pain after tonsillectomy: a meta-analysis. *Laryngoscope*. <https://doi.org/10.1002/lary.25636>.
- Iossifov I, Ronemus M, Levy D, Wang Z, Hakker I, Rosenbaum J *et al.* (2012). De novo gene disruptions in children on the autistic spectrum. *Neuron* 74: 285–299.
- Ivanov SV, Ward JM, Tessarollo L, McAreavey D, Sachdev V, Fananapazir L *et al.* (2004). Animal model – cerebellar ataxia, seizures, premature death, and cardiac abnormalities in mice with targeted disruption of the *Cacna2d2* gene. *Am J Pathol* 165: 1007–1018.
- Johnson RW, Rice AS (2014). Clinical practice. Postherpetic neuralgia. *N Engl J Med* 371: 1526–1533.
- Jun K, Piedras-Rentería ES, Smith SM, Wheeler DB, Lee SB, Lee TG *et al.* (1999). Ablation of P/Q-type $\text{Ca}(2+)$ channel currents, altered synaptic transmission, and progressive ataxia in mice lacking the alpha(1A)-subunit. *Proc Natl Acad Sci U S A* 96: 15245–15250.
- Kharasch ED, Eisenach JC (2016). wherefore gabapentinoids?: was there rush too soon to judgment? *Anesthesiology* 124: 10–12.
- Kim DS, Li KW, Boroujerdi A, Peter Yu Y, Zhou CY, Deng P *et al.* (2012). Thrombospondin-4 contributes to spinal sensitization and neuropathic pain states. *J Neurosci* 32: 8977–8987.
- Knoflach D, Kerov V, Sartori SB, Obermair GJ, Schmuckermair C, Liu X *et al.* (2013). Cav1.4 IT mouse as model for vision impairment in human congenital stationary night blindness type 2. *Channels* 7: 503–513.
- Korber C, Kuner T (2016). Molecular machines regulating the release probability of synaptic vesicles at the active zone. *Front Synaptic Neurosci* 8: 5.
- Kukkar A, Bali A, Singh N, Jaggi AS (2013). Implications and mechanism of action of gabapentin in neuropathic pain. *Arch Pharm Res* 36: 237–251.
- Lana B, Page KM, Kadurin I, Ho S, Nieto-Rostro M, Dolphin AC (2016). Thrombospondin-4 reduces binding affinity of [(3)H]-gabapentin to calcium-channel $\alpha_2\delta$ -1-subunit but does not interact with $\alpha_2\delta$ -1 on the cell-surface when co-expressed. *Sci Rep* 6: 24531.
- Lana B, Schlick B, Martin S, Pratt WS, Page KM, Goncalves L *et al.* (2014). Differential upregulation in DRG neurons of an alpha2delta-1 splice variant with a lower affinity for gabapentin after peripheral sensory nerve injury. *Pain* 155: 522–533.
- Landmark CJ, Beiske G, Baftiu A, Burns ML, Johannessen SI (2015). Experience from therapeutic drug monitoring and gender aspects of gabapentin and pregabalin in clinical practice. *Seizure-Eur J Epilep* 28: 88–91.
- Lee S (2013). Pharmacological inhibition of voltage-gated $\text{Ca}(2+)$ channels for chronic pain relief. *Curr Neuropharmacol* 11: 606–620.
- Li CY, Song YH, Higuera ES, Luo ZD (2004). Spinal dorsal horn calcium channel alpha2delta-1 subunit upregulation contributes to peripheral nerve injury-induced tactile allodynia. *J Neurosci* 24: 8494–8499.
- Li CY, Zhang XL, Matthews EA, Li KW, Kurwa A, Boroujerdi A *et al.* (2006). Calcium channel alpha2delta1 subunit mediates spinal hyperexcitability in pain modulation. *Pain* 125: 20–34.
- Li KW, Kim DS, Zaucke F, Luo ZD (2014a). Trigeminal nerve injury-induced thrombospondin-4 up-regulation contributes to orofacial neuropathic pain states in a rat model. *Eur J Pain* 18: 489–495.
- Li KW, Yu YP, Zhou C, Kim DS, Lin B, Sharp K *et al.* (2014b). Calcium channel alpha2delta1 proteins mediate trigeminal neuropathic pain states associated with aberrant excitatory synaptogenesis. *J Biol Chem* 289: 7025–7037.
- Li Z, Taylor CP, Weber M, Piechan J, Prior F, Bian F *et al.* (2011). Pregabalin is a potent and selective ligand for alpha(2)delta-1 and alpha(2)delta-2 calcium channel subunits. *Eur J Pharmacol* 667: 80–90.
- Lipscombe D, Andrade A, Allen SE (2013). Alternative splicing: functional diversity among voltage-gated calcium channels and behavioral consequences. *Biochim Biophys Acta* 1828: 1522–1529.
- Liu XG, Zhou LJ (2015). Long-term potentiation at spinal C-fiber synapses: a target for pathological pain. *Curr Pharm Des* 21: 895–905.
- Loh E, Guy SD, Mehta S, Moulin DE, Bryce TN, Middleton JW *et al.* (2016). The CanPain SCI clinical practice guidelines for rehabilitation management of neuropathic pain after spinal cord: introduction, methodology and recommendation overview. *Spinal Cord* 54 (Suppl 1): S1–S6.
- Lotarski S, Hain H, Peterson J, Galvin S, Strenkowski B, Donevan S *et al.* (2014). Anticonvulsant activity of pregabalin in the maximal electroshock-induced seizure assay in alpha2delta1 (R217A) and alpha2delta2 (R279A) mouse mutants. *Epilepsy Res* 108: 833–842.
- Lotarski SM, Donevan S, El-Kattan A, Osgood S, Poe J, Taylor CP *et al.* (2011). Anxiolytic-like activity of pregabalin in the Vogel conflict test in alpha2delta-1 (R217A) and alpha2delta-2 (R279A) mouse mutants. *J Pharmacol Exp Ther* 338: 615–621.
- Lunn TH, Husted H, Laursen MB, Hansen LT, Kehlet H (2015). Analgesic and sedative effects of perioperative gabapentin in total

- knee arthroplasty: a randomized, double-blind, placebo-controlled dose-finding study. *Pain* 156: 2438–2448.
- Luo ZD (2000). Rat dorsal root ganglia express distinctive forms of the $\alpha_2\delta$ calcium channel subunit. *Neuroreport* 11: 3449–3452.
- Luo ZD, Calcutt NA, Higuera ES, Valder CR, Song YH, Svensson CI *et al.* (2002). Injury type-specific calcium channel $\alpha_2\delta$ -1 subunit up-regulation in rat neuropathic pain models correlates with antiallodynic effects of gabapentin. *J Pharmacol Exp Ther* 303: 1199–1205.
- Luo ZD, Chaplan SR, Higuera ES, Sorkin LS, Stauderman KA, Williams ME *et al.* (2001). Upregulation of dorsal root ganglion ($\alpha_2\delta$) calcium channel subunit and its correlation with allodynia in spinal nerve-injured rats. *J Neurosci* 21: 1868–1875.
- Marais E, Klugbauer N, Hofmann F (2001a). Calcium channel $\alpha_2\delta$ subunits-structure and gabapentin binding. *Mol Pharmacol* 59: 1243–1248.
- Marais E, Klugbauer N, Hofmann F (2001b). Calcium channel $\alpha_2\delta$ subunits – Structure and gabapentin binding. *Mol Pharmacol* 59: 1243–1248.
- Mardani-Kivi M, Mobarakeh MK, Keyhani S, Motlagh KH, Ekhtiari KS (2013). Is gabapentin effective on pain management after arthroscopic anterior cruciate ligament reconstruction? A triple blinded randomized controlled trial. *Arch Bone Jt Surg* 1: 18–22.
- Margas W, Ferron L, Nieto-Rostro M, Schwartz A, Dolphin AC (2016). Effect of knockout of $\alpha_2\delta$ -1 on action potentials in mouse sensory neurons. *Philos Trans R Soc Lond B Biol Sci* 371: 20150430.
- Mark MD, Maejima T, Kuckelsberg D, Yoo JW, Hyde RA, Shah V *et al.* (2011). Delayed postnatal loss of P/Q-type calcium channels recapitulates the absence epilepsy, dyskinesia, and ataxia phenotypes of genomic *Cacna1A* mutations. *J Neurosci* 31: 4311–4326.
- Martinez V, Carles M, Marret E, Beloeil H, Regional Anaesthesia PCotFSoAIC (2017a). Perioperative use of gabapentinoids in France. *Anaesth Crit Care Pain Med*. In Press.
- Martinez V, Pichard X, Fletcher D (2017b). Perioperative pregabalin administration does not prevent chronic postoperative pain. Systematic review with a meta-analysis of randomized trials. *Pain* 158: 775–783.
- Mishriky BM, Waldron NH, Habib AS (2015). Impact of pregabalin on acute and persistent postoperative pain: a systematic review and meta-analysis. *Br J Anaesth* 114: 10–31.
- Moore RA, Wiffen PJ, Derry S, Toelle T, Rice AS (2014). Gabapentin for chronic neuropathic pain and fibromyalgia in adults. *Cochrane Database Syst Rev* 4: CD007938.
- Moulin DE, Clark AJ, Gilron I, Ware MA, Watson CP, Sessle BJ *et al.* (2007). Pharmacological management of chronic neuropathic pain - consensus statement and guidelines from the Canadian Pain Society. *Pain Res Manag* 12: 13–21.
- Naka A, Gruber-Schoffnegger D, Sandkuhler J (2013). Non-Hebbian plasticity at C-fiber synapses in rat spinal cord lamina I neurons. *Pain* 154: 1333–1342.
- Neely GG, Hess A, Costigan M, Keene AC, Goulas S, Langeslag M *et al.* (2010). A genome-wide *Drosophila* screen for heat nociception identifies $\alpha_2\delta_3$ as an evolutionarily conserved pain gene. *Cell* 143: 628–638.
- Newton RA, Bingham S, Case PC, Sanger GJ, Lawson SN (2001). Dorsal root ganglion neurons show increased expression of the calcium channel $\alpha_2\delta$ -1 subunit following partial sciatic nerve injury. *Brain Res Mol Brain Res* 95: 1–8.
- Nguyen D, Deng P, Matthews EA, Kim DS, Feng G, Dickenson AH *et al.* (2009). Enhanced pre-synaptic glutamate release in deep-dorsal horn contributes to calcium channel $\alpha_2\delta$ -1 protein-mediated spinal sensitization and behavioral hypersensitivity. *Mol Pain* 5: 6.
- Pan B, Guo Y, Wu HE, Park J, Trinh VN, Luo ZD *et al.* (2016). Thrombospondin-4 divergently regulates voltage-gated Ca^{2+} channel subtypes in sensory neurons after nerve injury. *Pain* 157: 2068–2080.
- Pan B, Yu H, Park J, Yu YP, Luo ZD, Hogan QH (2015). Painful nerve injury upregulates thrombospondin-4 expression in dorsal root ganglia. *J Neurosci Res* 93: 443–453.
- Park J, Luo ZD (2010). Calcium channel functions in pain processing. *Channels (Austin)* 4: 510–517.
- Park J, Yu YP, Zhou CY, Li KW, Wang D, Chang E *et al.* (2016). Central mechanisms mediating thrombospondin-4-induced pain states. *J Biol Chem* 291: 13335–13348.
- Patel R, Bauer CS, Nieto-Rostro M, Margas W, Ferron L, Chaggar K *et al.* (2013). $\alpha_2\delta$ -1 gene deletion affects somatosensory neuron function and delays mechanical hypersensitivity in response to peripheral nerve damage. *J Neurosci* 33: 16412–16426.
- Perez-Reyes E (2003). Molecular physiology of low-voltage-activated t-type calcium channels. *Physiol Rev* 83: 117–161.
- Perret D, Luo ZD (2009). Targeting voltage-gated calcium channels for neuropathic pain management. *Neurotherapeutics* 6: 679–692.
- Pippucci T, Parmeggiani A, Palombo F, Maresca A, Angius A, Crisponi L *et al.* (2013). A novel null homozygous mutation confirms *CACNA2D2* as a gene mutated in epileptic encephalopathy. *PLoS ONE [Electronic Resource]* 8: e82154.
- Ravindran D (2014). Chronic postsurgical pain: prevention and management. *J Pain Palliat Care Pharmacother* 28: 51–53.
- Russo RE, Hounsgaard J (1994). Short-term plasticity in turtle dorsal horn neurons mediated by L-type Ca^{2+} channels. *Neuroscience* 61: 191–197.
- Simms BA, Zamponi GW (2014). Neuronal voltage-gated calcium channels: structure, function, and dysfunction. *Neuron* 82: 24–45.
- Southan C, Sharman JL, Benson HE, Faccenda E, Pawson AJ, Alexander SPH *et al.* (2016). The IUPHAR/BPS guide to PHARMACOLOGY in 2016: towards curated quantitative interactions between 1300 protein targets and 6000 ligands. *Nucl Acids Res* 44: D1054–D1068.
- Stanfa LC, Singh L, Williams RG, Dickenson AH (1997). Gabapentin, ineffective in normal rats, markedly reduces C-fibre evoked responses after inflammation. *Neuroreport* 8: 587–590.
- Tran-Van-Minh A, Dolphin AC (2010). The $\alpha_2\delta$ ligand gabapentin inhibits the Rab11-dependent recycling of the calcium channel subunit $\alpha_2\delta$ -2. *J Neurosci* 30: 12856–12867.
- Turner RW, Zamponi GW (2014). T-type channels buddy up. *Pflugers Arch* 466: 661–675.
- Valder CR, Liu JJ, Song YH, Luo ZD (2003). Coupling gene chip analyses and rat genetic variances in identifying potential target genes that may contribute to neuropathic allodynia development. *J Neurochem* 87: 560–573.
- Vargas-Espinosa ML, Sanmarti-Garcia G, Vazquez-Delgado E, Gay-Escoda C (2012). Antiepileptic drugs for the treatment of neuropathic pain: a systematic review. *Med Oral Patol Oral Cir Bucal* 17: e786–e793.

- Vergult S, Dheedene A, Meurs A, Faes F, Isidor B, Janssens S *et al.* (2015). Genomic aberrations of the CACNA2D1 gene in three patients with epilepsy and intellectual disability. *Eur J Hum Genet* 23: 628–632.
- Verma V, Singh N, Jaggi AS (2014). Pregabalin in neuropathic pain: evidences and possible mechanisms. *Current Neuropharmacol* 12: 44–56.
- Vink S, Alewood PF (2012). Targeting voltage-gated calcium channels: developments in peptide and small-molecule inhibitors for the treatment of neuropathic pain. *Br J Pharmacol* 167: 970–989.
- Wang T, Jones RT, Whippen JM, Davis GW (2016). $\alpha_2\delta_3$ is required for rapid Transsynaptic homeostatic signaling. *Cell Rep* 16: 2875–2888.
- Wheeler DG, Groth RD, Ma H, Barrett CF, Owen SF, Safa P *et al.* (2012). Ca(V)1 and Ca(V)2 channels engage distinct modes of Ca(2+) signaling to control CREB-dependent gene expression. *Cell* 149: 1112–1124.
- Wu J, Yan Z, Li Z, Qian X, Lu S, Dong M *et al.* (2016). Structure of the voltage-gated calcium channel Ca(v)1.1 at 3.6 Å resolution. *Nature* 537: 191–196.
- Wycisk KA, Budde B, Feil S, Skosyrski S, Buzzi F, Neidhardt J *et al.* (2006a). Structural and functional abnormalities of retinal ribbon synapses due to Cacna2d4 mutation. *Invest Ophthalmol Vis Sci* 47: 3523–3530.
- Wycisk KA, Zeitz C, Feil S, Wittmer M, Forster U, Neidhardt J *et al.* (2006b). Mutation in the auxiliary calcium-channel subunit CACNA2D4 causes autosomal recessive cone dystrophy. *Am J Hum Genet* 79: 973–977.
- Youn DH, Gerber G, Sather WA (2013). Ionotropic glutamate receptors and voltage-gated Ca(2)(+) channels in long-term potentiation of spinal dorsal horn synapses and pain hypersensitivity. *Neural Plast* 2013: 654257.
- Yu YP, Kweon TD, Vo B, Gong N, Luo ZD (2017). Thrombospondin 4 acts on pre-synaptic calcium channel alpha-2-delta-1 subunits to induce synaptogenesis between sensory and spinal cord neurons through a T-type calcium channel pathway that is blocked by early, low-dose gabapentin treatments. *in revision*.
- Yusaf SP, Goodman J, Gonzalez IM, Bramwell S, Pinnock RD, Dixon AK *et al.* (2001). Streptozocin-induced neuropathy is associated with altered expression of voltage-gated calcium channel subunit mRNAs in rat dorsal root ganglion neurones. *Biochem Biophys Res Commun* 289: 402–406.
- Zamponi GW (2016). Targeting voltage-gated calcium channels in neurological and psychiatric diseases. *Nat Rev Drug Discov* 15: 19–34.
- Zamponi GW, Striessnig J, Koschak A, Dolphin AC (2015). The physiology, pathology, and pharmacology of voltage-gated calcium channels and their future therapeutic potential. *Pharmacol Rev* 67: 821–870.
- Zhou C, Luo ZD (2013). Electrophysiological characterization of spinal neuron sensitization by elevated calcium channel alpha-2-delta-1 subunit protein. *Eur J Pain* 18: 649–658.
- Zhou C, Luo ZD (2015). Nerve injury-induced calcium channel alpha-2-delta-1 protein dysregulation leads to increased pre-synaptic excitatory input into deep dorsal horn neurons and neuropathic allodynia. *Eur J Pain* 19: 1267–1276.