

UC Davis

UC Davis Previously Published Works

Title

Oleocanthal-rich extra virgin olive oil demonstrates acute anti-platelet effects in healthy men in a randomized trial

Permalink

<https://escholarship.org/uc/item/6vh0h3tp>

Authors

Agrawal, Karan

Melliou, Eleni

Li, Xueqi

et al.

Publication Date

2017-09-01

DOI

10.1016/j.jff.2017.06.046

Peer reviewed



Published in final edited form as:

J Funct Foods. 2017 September ; 36: 84–93. doi:10.1016/j.jff.2017.06.046.

Oleocanthal-rich extra virgin olive oil demonstrates acute anti-platelet effects in healthy men in a randomized trial

Karan Agrawal^{a,b}, Eleni Melliou^c, Xueqi Li^d, Theresa L. Pedersen^{e,1}, Selina C. Wang^{d,f}, Prokopios Magiatis^c, John W. Newman^{a,b,e}, and Roberta R. Holt^a

^aDepartment of Nutrition, University of California-Davis, One Shields Avenue, Davis, CA 95616, USA

^bWest Coast Metabolomics Center, Genome Center, 451 Health Sciences Drive, Davis, CA 95616, USA

^cDepartment of Pharmacognosy and Natural Products Chemistry, Faculty of Pharmacy, National and Kapodistrian University of Athens, Panepistimioupolis Zografou 15 771, Athens, Greece

^dUC Davis Olive Center, University of California-Davis, 392 Old Davis Road, Davis, CA 95616, USA

^eObesity and Metabolism Research Unit, USDA - Agricultural Research Service - Western Human Nutrition Research Center, 430 W Health Sciences Drive, Davis, CA 95616, USA

^fDepartment of Food Science and Technology, University of California-Davis, One Shields Avenue, Davis, CA 95616, USA

Abstract

The phenolic profiles of extra virgin olive oils (EVOOs) may influence their cardiovascular benefits. In a randomized crossover of acute EVOO intake on platelet function, participants (n=9) consumed 40 mL of EVOO weekly. EVOOs were matched for total phenolic content and were either tyrosol-poor with 1:2 oleacein/oleocanthal (D2;0.5), or 2:1 oleacein/oleocanthal (D2;2), or predominantly tyrosol (D2;0). Ibuprofen provided a platelet inhibition control. Blood was collected pre- and 2 hr post-EVOO intake. D2;0.5 and D2;2 reduced 1 µg/mL collagen-stimulated maximum platelet aggregation (P_{max}), with effects best correlated to oleocanthal intake (R=0.56, P=0.002). Total phenolic intake was independently correlated to eicosanoid production inhibition, suggesting that cyclooxygenase blockade was not responsible for the P_{max} inhibition. Five participants exhibited >25% P_{max} declines with D2;0.5 and D2;2 intake and plasma metabolomic profiles discriminated subjects by oil responsiveness. Platelet responses to acute EVOO intake are associated with oil phenolic composition and may be influenced by diet.

Corresponding Author: John W. Newman, PhD, 430 W Health Sciences Drive, Davis, CA 95616, Phone: (530) 752-1009, john.newman@ars.usda.gov.

¹Present Address: Advanced Analytics, 118 First Street, Woodland, CA 95695, USA

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of Interest: The authors declare no conflict of interest

Keywords

platelet aggregation; human; extra virgin olive oil; oleocanthal; oxylipins

1. Introduction

Recent dietary guidelines have moved towards promoting healthy dietary patterns that encourage the intake of a synergy of nutrients for the prevention of chronic disease (Mozaffarian, 2016). While the potential benefit of dietary patterns such as a Mediterranean diet are suggested from epidemiological observations and large controlled dietary interventions (Estruch et al., 2013; Mozaffarian, 2016), adaptation of these results to recommendations for diverse population groups can have several challenges, which include assessing the role of diet coupled to genetic variation and environmental influences. Further refinement of dietary recommendations is also complicated by the knowledge that varietal differences, as well as agricultural and manufacturing practices can affect the final nutrient content of the food product of interest (Mozaffarian, 2016).

Olive oil has long been considered a key component of a Mediterranean dietary pattern. Beyond the potential benefits of its major components such as monounsaturated fatty acids, the minor components of olive oil are also thought to be of benefit (Covas, Fito, & de la Torre, 2015). Extra virgin olive oils (EVOO) contain an array of phenolic antioxidants, from three major chemical classes which include the simple phenolics tyrosol and hydroxytyrosol, the secoiridoids oleuropein aglycon and oleocanthal, and lignans (Bendini et al., 2007). Phenolic content forms the basis of the European Food Safety Authority guidance on olive oil health claims for the prevention of low density lipoprotein oxidation. Specifically, oils must contain at least 5 mg of hydroxytyrosol and its derivatives (e.g. oleuropein complex and tyrosol) per 20 mg of olive oil to make such claims (EFSA Panel on Dietetic Products, Nutrition and Allergies, 2011). It is important to note that oils meeting this definition can have diverse phenolic profiles (Servili et al., 2014). Recent epidemiological evidence has indicated the potential utility of EVOO intake for the primary prevention of cardiovascular disease (Buckland & Gonzalez, 2015). Notably, intake of phenol-rich EVOO for close to five years reduced primary cardiovascular disease events in an elderly Spanish population at risk for cardiovascular disease (Estruch et al., 2013). The underlying mechanism of this cardioprotective effect is as yet unknown, however intervention trials with virgin and extra virgin olive oil have demonstrated anti-platelet and anti-inflammatory effects (Covas et al., 2015).

Atherosclerotic cardiovascular disease is a chronic inflammatory disease, initiated by endothelial damage and promoted by a number of cell types to include platelets (May, Seizer, & Gawaz, 2008). Therapeutic reduction of cardiovascular disease progression and events has focused on limiting platelet activation through inhibition of cyclooxygenase (COX), phosphodiesterase, adenosine diphosphate (ADP) receptors, and platelet-platelet interactions through glycoprotein IIb/IIIa (Yousuf & Bhatt, 2011). Major lipid remodelling occurs upon platelet activation that includes shape change, degranulation and generation of a number of bioactive species that amplify activation while promoting clot stabilization and

inflammation (O'Donnell, Murphy, & Watson, 2014). Oxylipins are a superclass of bioactive lipids produced from polyunsaturated fatty acid oxidation. The best known of these associated with clot formation is thromboxane, whose production is targeted through the inhibition of COX with aspirin. In addition, oxylipins produced from lipoxygenase (LOX) and cytochrome P450 may also have roles in platelet activation and inflammatory modulation (Tourdot, Ahmed, & Holinstat, 2013).

While clear mechanisms have yet to be demonstrated *in vivo*, structural specificity has been demonstrated for EVOO-derived phenolic impacts on platelet function and inflammatory processes *in vitro*. For example, hydroxytyrosol appears to be a more potent inhibitor of ADP- or collagen- induced platelet activation than other EVOO-derived phenolics including oleuropein (Petroni et al., 1995), while oleocanthal (the dialdehydic form of decarboxymethyl ligstroside aglycon) has received considerable interest as a COX inhibitor (Beauchamp et al., 2005), and oleacein (the dialdehydic form of decarboxymethyl oleuropein aglycon) has been reported as a 5-LOX inhibitor (Vougogiannopoulou et al., 2014). Since the characterization of specific phenolic profiles in EVOO interventions remains rare, it remains difficult to discriminate *in vivo* EVOO-associated effects of specific phenolics on the biochemical networks associated with cardiovascular disease initiation and progression.

Therefore, while substantial evidence exists to support the hypothesis that phytochemicals within EVOO are beneficial to health, it is still unclear if the total phenolic content or the specific phenolic profile is the more important aspect. In this pilot study, we seek to directly address this issue by comparing the impact of the acute intake of oleocanthal-rich and oleocanthal-poor EVOO, each containing an equivalent total phenolic content, on platelet reactivity in healthy adults. We hypothesize that the intake of oleocanthal-rich EVOO will reduce platelet aggregation to a greater extent than olive oils containing a lesser amount of oleocanthal, and that these effects will be related to changes in COX-dependent oxygenated lipids but not the 5-LOX-dependent or auto-oxidative polyunsaturated fatty acid metabolites.

2. Materials and Methods

2.1. Study Design

A double-blind, randomized controlled crossover study tested the acute effects of ingesting three unique EVOOs on platelet aggregation. Randomization was performed following a plan formulated via a random number generator and all participants were randomized in one block. All EVOO were provided in coded bottles of similar size, shape and colour. Therefore, the investigators responsible for the conduct of the intervention trial and the study participants were blinded as to the specific EVOO provided for any given study day. This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the institutional review board of the University of California, Davis. Written informed consent was obtained from all subjects before inclusion in the study. This study is also registered with [ClinicalTrials.gov](https://clinicaltrials.gov) as NCT02902913.

Participants were asked to participate in four study days, separated by at least one week (7.4 ± 1.6 days; one of 27 subject visits involving EVOO consumption occurred 14 days after the proceeding visit), at the University of California Davis, Department of Nutrition, Ragle Human Nutrition Center. For the first three study visits, the volunteers were asked to consume 40 mL of their assigned EVOO within 5 minutes. On the fourth and final study visit, participants consumed one 400 mg dose of ibuprofen by mouth. Participants were asked to refrain from taking drugs that are known or may affect platelet function (e.g. acetaminophen, aspirin and non-steroidal anti-inflammatory drugs) or consuming olive oil for at least one week prior to their first study day visit, and throughout the study period. Subjects were also asked to refrain from consumption of foods previously described to affect platelet function for at least 24 hours prior to each study visit. A list of these foods was provided to the subjects during recruitment, and included phenolic- and polyphenol-rich foods, such as, cocoa products, coffee, tea, wine, other grape products and other colourful fruits and vegetables (Holt, Heiss, Kelm, & Keen, 2012; Miller, Rice, Garrett, & Stein, 2014). Dietary compliance was confirmed via questioning of the individual participants upon their arrival to the facility. Prior to each study day, volunteers were asked to participate in an overnight fast (no food or water 12 hr prior to their scheduled visit). See Fig. A.1 for a diagrammatic representation of the study design.

2.2. Participants

Healthy men aged 20–50 years and willing to drink 40 mL of EVOO were recruited by public announcement, and assessed by telephone interview and clinical screening prior to enrolment. Exclusion criteria included current supplement and/or prescription drug use; the daily use of aspirin or other non-steroidal anti-inflammatory drugs; a history of cardiovascular disease; vegetarian, vegan, or non-traditional diets; diets that promoted weight loss or that deviated significantly from the average diet of the general population; elevated lipids beyond levels recommended for lifestyle intervention; any self-reported bleeding or coagulation disorder.

2.3. Olive Oil Selection and Characterization

Olives (*Olea europaea L.*) harvested between November 2014 and January 2015 were used to generate EVOOs discussed in this study. In preparation for this study, 300 commercial samples were screened from Greece and California for chemical composition analysis. Oils were first selected to include extra virgin grades with similar total phenolic content as determined by the Folin-Ciocalteu method (Garcia, Coelho, Costa, Pinto, & Paiva-Martins, 2013), similar lipid profiles (International Olive Oil Council, 2001), and similar peroxide values (AOAC International, 2013). The subset of oils meeting these criteria underwent complete phenolic characterization, with final selection of test oils based on their specific phenolic profile using a recently proposed EVOO bioactive phenolic index ((Karkoula, Skantzari, Melliou, & Magiatis, 2012). This “D” index (D = University of California, Davis) provides the sum of oleocanthal and oleacein (D1 index, D1_i), and the oleacein/oleocanthal ratio (D2 index, D2_i). The D2 index was inconsistently defined by Karkoula et al. in their original report (Karkoula et al., 2012), but was calculated correctly in the manuscript and has been consistently defined since (Karkoula, Skantzari, Melliou, & Magiatis, 2014). Characteristics of the test oils are shown in Table 1. While all three test oils had a similar

total phenolic content, two of the test oils were similar in their $D1_i$, but inverted in the $D2_i$, with both oils containing trace or no detectable levels of other secoiridoid conjugated phenolics. The third oil was chosen as a control and had $D1_i$ of zero. All oils were stored at 4°C immediately after their analysis and during the study period to minimize possible alterations in chemical composition.

2.4. EVOO qNMR Analysis

All commercial oils were analysed by quantitative ^1H NMR (qNMR) to determine their oleocanthal, oleacein and total phenolic content. Sample preparation for qNMR analysis consisted of homogenizing 5 g of EVOO with 20 mL cyclohexane and 25 mL acetonitrile for 30 seconds and then centrifuging the homogenate for 5 min at $4000 \times g$. The acetonitrile phase was subsequently mixed with 1 mL of a syringaldehyde (4-hydroxy-3,5-dimethoxybenzaldehyde) solution (0.5 mg/mL in acetonitrile) and evaporated under reduced pressure using a rotary evaporator (Buchi, Switzerland). Samples were reconstituted in 750 μL of deuterated chloroform (CDCl_3) and 550 μL of the reconstituted solution was transferred to a 5 mm NMR tube.

^1H -NMR spectra were recorded at 600 MHz using an Avance600 spectrometer (Bruker, Billerica, MA). Typically, 50 scans were collected into 32K data points over a spectral width of 0–16 ppm with a relaxation delay of 1 sec and an acquisition time of 1.7 sec. Prior to Fourier transformation, an exponential weighting factor corresponding to a line broadening of 0.3 Hz was applied. The spectra were phased corrected and integrated automatically using Bruker TopSpin software. Manual integrations were performed to enhance accuracy peaks of interest when necessary. Quantitation of oleocanthal and oleacein was performed using calibration curves as previously described (E. Karkoula et al., 2012).

2.5. EVOO UPLC-DAD Analysis

The oils selected above were further subjected to semi-quantitative phenolic profiling using ultra performance liquid chromatography coupled to a diode array detector (UPLC-DAD). Phenolic compounds were extracted from oils using modifications of published solid phase extraction protocols (Gutierrez-Rosales, Rios, & Gomez-Rey, 2003). The pre-weighed oil sample (2.5 ± 0.001 g) and 0.5 mL of an internal standard solution consisting of 67.5 $\mu\text{g}/\text{mL}$ *p*-hydroxyphenyl-acetic acid (Sigma Aldrich, St Louis, MO) was dissolved in 6 mL of hexane. A 1g-6mL diol-bonded phase cartridge (Thermo Scientific, Waltham, MA, USA) was conditioned with 6 mL of methanol and 6 mL of hexane consecutively, by vacuum elution. Cartridges were then loaded with oil, washed twice with 6 mL of hexane, once with 6 mL of hexane/ethyl acetate (90:10, v/v), and eluted with 10 mL of methanol. The eluent was reduced to dryness by rotary evaporation at room temperature and the residue was reconstituted in 1 mL of methanol/water (1:1, v/v).

The UPLC-DAD protocol was adapted from a previously published method (Daskalaki, Kefi, Kotsiou, & Tasioula-Margari, 2009). A 20 μL aliquot of the extract was injected onto a 4.6×250 mm, 5 μm C18 column (Agilent Technologies, Santa Clara, CA, USA) and separated by gradient elution at a flow rate of 1 mL/min using mobile phases of 98:2, water/acetic acid (v/v) (Solvent A) and 1:1, methanol/acetonitrile (v/v) (Solvent B). The mobile

phase gradient was as follows: 0 min – 5% B; 25 min – 30% B; 50 min – 35% B; 65 min – 70% B; 70 min – 100% B; 5 min re-equilibration. Absorbance was recorded at 280 nm and 340 nm. Analytes were quantified by relative response to the internal standard.

2.6. Blood Collection

Blood was collected at 0 and 2 hr after test product ingestion from an antecubital vein in Vacutainer® tubes (Beckton Dickinson, Franklin Lakes, NJ, USA) containing sodium citrate for platelet aggregometry and post-stimulation oxylipin analysis or potassium EDTA for metabolomics analysis. All collection procedures were performed between 08:00 and 13:00 to avoid circadian effects.

2.7. Platelet Aggregometry Measurements

Optical platelet aggregometry was performed in citrated blood using a modified version of the method by Born and Cross (Born & Cross, 1963). Whole blood impedance lumi-aggregometry was measured in duplicate using a 2-channel Chrono-Log 700 (Havertown, PA). Fifteen minutes after the blood draw, platelet rich plasma (PRP) was separated from whole blood by centrifugation ($200 \times g$, 10 min at 25°C). After the top 75% of the PRP layer was collected into a separate tube, platelet counts were manually obtained using a haemocytometer. The final platelet count was adjusted to 250,000 cells/ μL with autologous platelet poor plasma obtained by re-centrifugation of the whole blood tubes at $1500 \times g$ for 10 min at 25°C . After resting the PRP for a minimum of 15 min, platelet aggregation testing commenced. PRP was incubated at 37°C for a minimum of 2 min prior to stimulation with collagen at a final concentration of either 1 or 3 $\mu\text{g}/\text{mL}$ in duplicate at a stirring speed of 1200 rpm and showed intra-assay mean and standard error of $10 \pm 2\%$. After 8 min of data collection, stimulated and unstimulated PRP samples were immediately centrifuged for 3 min at $1500 \times g$, and frozen at -80°C for the oxylipin analysis.

2.8. Oxylipin Analysis

Oxylipins derived from COX, LOX, and cytochrome P450 dependent metabolism of arachidonic acid (AA) were quantified using UPLC with tandem quadrupole mass spectrometry in 100 μL of stimulated or unstimulated PRP plasma collected at 0 and 2 hr post-EVOO or ibuprofen ingestion. Sample preparation generally followed a previously published protocol (La Frano et al., 2017). Briefly, the plasma was enriched with a suite of deuterated surrogates and extracted with 300 μL acetonitrile with 1% formic acid using an Ostro™ Pass-through Sample Preparation Plate (Waters Corp; Milford, MA). Extracts were concentrated and reconstituted in internal standard solution prior to analysis. Analytical targets were separated on a 2.1×150 mm, 1.7 μm BEH C18 column (Waters; Milford, MA) and detected by negative mode electrospray ionization and tandem mass spectrometry on a 4000 QTRAP® (Sciex; Framingham, MA) as previously described (Agrawal et al., 2017). Calibrants and internal standards were purchased from Cayman Chemical (Ann Arbor, MI), Avanti Polar Lipids Inc. (Alabaster, AL), and Larodan Fine Lipids (Malmo, Sweden). Data was processed utilizing MultiQuant™ v. 3.0.2 (Sciex). Inter- and intra-assay variability was less than 20% for all reported analytes.

2.9. Metabolomics Analysis

Small molecules were semi-quantitatively determined in EDTA plasma collected at baseline and 2 hr post-EVOO ingestion by gas chromatography time-of-flight mass spectrometry as previously described (Fahrman et al., 2015). Briefly, 30 μ L aliquots of plasma, were thawed on ice, extracted with methyl tert-butyl ether, derivatized by methoximation/silylation, and analysed. Data, reported as quantitative ion peak heights, were normalized by the sum intensity of all annotated metabolites and used for further statistical analysis. Compounds were identified by processing acquired spectra using the BinBase database (Fiehn, Wohlgemuth, & Scholz, 2005; Scholz & Fiehn, 2007). All analyses were performed in a single run. Inter- and intra-assay variability in normalized metabolomics data is routinely <25% using the described methods.

2.10. Statistical Analysis

All changes () in maximum platelet aggregation (Pmax) and oxylipin concentrations were assessed by repeated measures ANOVA with a threshold of $P < 0.05$ using GraphPad Prism 6 (La Jolla, CA) considering all subjects. Due to high inter-individual variability, Pmax and oxylipin concentration results were range-scaled to each subject's minimum and maximum responses across all four treatments. This practice promotes the identification of biological effects (van den Berg, Hoefsloot, Westerhuis, Smilde, & van der Werf, 2006) and by reducing the influence of intra-individual variability (Krishnan, Newman, Hembrooke, & Keim, 2012).

To identify the main component(s) of the EVOOs that best predicted the observed changes in Pmax or oxylipin concentration, multiple linear regression analyses were conducted using JMP Pro v12 (Cary, NC) considering all subjects. Regression models were built using either Pmax or the sum of selected oxylipin concentrations as the dependent variable, and the mg/kg of body weight dose of various individual EVOO phenolics, total EVOO phenolic content or total EVOO peroxide content as the independent variable. Treatment and subject orders were considered and had no effect on regression results. Oxylipins used in the regression model have been associated with platelet function (Tourdot et al., 2013), and included the sum of the COX products thromboxane B2 (TXB2), prostaglandin E2 (PGE2) and 11-hydroxyeicosatetraenoic acid (11-HETE), and the LOX products 12-HETE, 15-HETE (collectively referred to as [COX + LOX]). Individual phenolics used in the regression model were oleocanthal, oleacein, and tyrosol as the mg/kg of body weight dose of these phenolics were > 0.1 mg/kg for at least one EVOO consumed. Model selection was made using the corrected Akaike Information Criterion (AICc) and Bayesian Information Criterion (BIC) scores, with differences in AICc or BIC scores > 2 considered statistically different (Chatterjee & Hadi, 2015; Ward, 2008).

In addition to the statistical analysis described above sources of variability in the subjects' platelet aggregation response were explored. Specifically, plasma metabolomics profiling pre- and post-EVOO intake was evaluated to determine if these profiles would predictably vary with platelet aggregation responsiveness and may reflect factors not controlled by our study design (e.g. habitual diet, fitness, metabolic variance). Metabolomics data were assessed by partial least squares-discriminant analysis (PLS-DA), with group classification

based on ingested oils and a Pmax decrease of > or <25% in following D2i0.5 or D2i2 consumption. These multivariate analyses were performed in the R-statistical environment (R Foundation for Statistical Computing, Vienna, Austria) using imDEV v1.42, a Microsoft Excel (Microsoft Corporation, Redmond, Washington) Add-In (Grapov & Newman, 2012). Prior to PLS-DA, data were curated such that analytes with < 70% completeness of data were removed from consideration. Curated data were screened for outliers using the Grubb's test (Grubbs, 1950), and missing data were imputed by a two-component probabilistic principle components analysis with univariate scaling (Wang & Wang, 2006). Following normalization of data according to the procedures of Box and Cox (Box & Cox, 1964), PLS-DA was conducted using the orthogonal scores algorithm with univariate scaling and leave-one-out cross-validation. Variables were clustered by Spearman correlation coefficients using the Minkowski distance and Ward agglomeration.

Sample size was estimated for the primary outcome measure from previous data that observed flavonoid-induced changes in collagen-induced platelet aggregation (Hubbard, Wolfram, Lovegrove, & Gibbins, 2004). For a mean difference of 40% and an expected standard deviation of residuals of 25% with a two-tailed α set at 0.05, a minimum of $n = 9$ provides sufficient power at 0.8 to detect a significant difference between three groups with ANOVA.

3. Results

3.1. Subject Recruitment and Retention

Recruitment and interventions occurred between January and September of 2015 and continued until 10 subjects qualified for enrolment. Out of 11 subjects screened for this study, 10 qualified for enrolment and 9 were included in and completed the study. One subject chose not to participate, citing difficulty in giving up olive oil. Baseline physiological characteristics of the enrolled subjects are listed in Table A.1. The average age of the study participants was 26 ± 4 yr with a body mass index of 25.5 ± 4.1 kg/m². In general, the subject population was healthy, with plasma lipids and glucose within the normal reference range, and a mean blood pressure of 124 ± 9 mmHg systolic and 77 ± 4 mmHg diastolic.

3.2. Olive Oil Composition

Of the 300 extra virgin olive oils screened and analysed, three oils were selected: a commercial oil of Mediterranean origin obtained from a California supermarket with undetectable oleocanthal and oleacein (D2i0); an Arbequina monovarietal oil provided by Corto Olive Co. (Lodi CA) harvested in November 2014 in California with 2:1 oleacein/oleocanthal (D2i2); a Koroneiki monovarietal blend of oils harvested in November 2014 from the Akritohori, Leukohora and Lambaina regions in Kalamata, Greece with 1:2 oleacein/oleocanthal (D2i0.5). Each could be classified as good quality oil, having high phenolic content, as well as low peroxide values and free fatty acid content, K232 values < 2.5, and K270 values < 0.22 (Table 1). Concentrations of oleuropein- and ligstroside-derived phenolics quantified by ¹H NMR demonstrate their differences in tyrosol, oleocanthal and oleacein content, while their UPLC-DAD profiles suggest levels of other phenolics were all

low and quite similar except for apparently high levels of 1-acetoxypinoresorsinol in the Arbequina oil (D2;2). The D2;0 oil contained <10 mg/kg of both oleocanthal and oleacein, but was high in tyrosol compared to the other two oils, while both the D2;2 and D2;0.5 oils contained < 10 mg/kg of tyrosol with a D1 index of 460 and 484 mg/kg, respectively. The phenolic content of D2;0.5 and D2;2 was close to the average values reported for Messenia, the main olive oil producing region of Greece (Evangelia Karkoula et al., 2014). Oil total phenolics and oleacein content increased together and were highest in D2;2, whereas oil peroxide values and oleocanthal were inversely related (Table 1). Data for the complete characterization of all screened oil are available upon request.

3.3. Effects on Platelet Aggregation

Between day differences in basal platelet aggregation responses were not observed (Table 2). Ibuprofen treatment (400 mg) served as an assessment of individual platelet response to a known platelet inhibitor. Ibuprofen treatment reduced Pmax by $57.5 \pm 32.9\%$ with 3 $\mu\text{g/mL}$ collagen-stimulation, and by $71.8 \pm 14.3\%$ with 1 $\mu\text{g/mL}$ collagen-stimulation in all subjects. While EVOO intake did not affect 2 hr 3 $\mu\text{g/mL}$ collagen-stimulated Pmax, 1 $\mu\text{g/mL}$ collagen-stimulated Pmax was decreased with a large effect size ($\eta^2 = 0.2$) by both D2;0.5 ($-35 \pm 39\%$) and D2;2 ($-13 \pm 36\%$) intake compared to D2;0 ($7 \pm 24\%$) as seen in Fig. 1A.

Regression analyses were performed to attempt to predict the Pmax in terms of the estimated amount per subject body weight of oleocanthal, oleacein, tyrosol, total EVOO phenolics, or total EVOO peroxides consumed by each subject. Oleocanthal provided the strongest individual Pmax prediction ($R = 0.563$, $P = 0.002$), though a two-factor model of oleocanthal and peroxides ($R = 0.640$, $P = 0.002$), appeared slightly better with an AICc improvement of 1.2. The model was not significantly improved by the addition of tyrosol, oleacein or total EVOO phenolics as factors (Table 3).

3.4. Effects on Oxylipins

This study measured the non-esterified pool of oxylipins produced from the collagen-induced platelet aggregate. Basal oxylipin levels were determined from unstimulated PRP. While a less sensitive measure of lipid autooxidation than inspection of the esterified lipid pool, the presence of high levels of non-esterified 9-HETE and or F2-isoprostanes are still viable measures of autooxidation, if observed. The relative abundance of the measured AA mid-chain alcohols (5-, 8-, 9-, 11- and 15-HETE) were not altered by EVOO consumption, and F2-isoprostanes were not observed. Ibuprofen treatment decreased 1 $\mu\text{g/mL}$ collagen-stimulated oxylipin concentrations compared to the three EVOOs in all subjects (Fig. 1B). EVOO intake did not change either the 1 $\mu\text{g/mL}$ collagen-stimulated individual (Table A.2) or [COX + LOX] product formation (Fig. 1B). Changes in the other detected COX1 product, 6-keto-PGF1a were also not seen between EVOOs (Table A.2).

As changes in eicosanoid production was observed, regression analyses were performed assess the ability of other factors to predict the [COX + LOX]. Specifically, the estimated amount per subject body weight of oleocanthal, oleacein, tyrosol, total EVOO phenolics, or total EVOO peroxides consumed by each subject were evaluated. Total EVOO phenolic

intake provided the strongest prediction of oxylipin production ($R = 0.593$, $P = 0.001$) with the model not significantly improved by the addition of tyrosol, oleacein, oleocanthal, or total EVOO peroxides as factors (Table 4).

3.5. Identification of a Responsive Platelet Aggregation Phenotype

Exploration of individual variability in platelet aggregation response showed five of the nine subjects exhibited >25% reduction in 1 $\mu\text{g/mL}$ collagen-stimulated P_{max} with D2;0.5 or D2;2 EVOO intake. This 25% reduction in P_{max} was used to define a responsive phenotype. Within this group of individuals, 1 $\mu\text{g/mL}$ collagen-stimulated production of TXB₂, 11-HETE and 15-HETE was reduced after D2;2 intake compared to non-responders (Table 5).

Metabolomics analysis of compounds derived from primary metabolism conducted the pre- and post-EVOO intake plasma showed a total of 311 metabolites of which 107 were identified using the BinBase database. Using this data, PLS-DA discriminated EVOO responsive phenotypes (Fig. 2). In general, subjects that “responded” to EVOO intake showed increased plasma concentrations of carbohydrates such as glucose, xylose and pinitol, as well as sugar acids such as glycolic acid, gluconic acid and threonic acid, both prior to, and after EVOO intake. On the other hand, subjects that did not “respond” to EVOO intake had increased fasting plasma non-esterified fatty acid concentrations (particularly oleic acid) that remained elevated relative to “responders” after EVOO intake, as well as increased levels of the citric acid cycle metabolites malic acid, isocitric acid and citric acid. See Fig. A.2 for a partial list of identified metabolites, their metabolic relationships and their impact on the separation between “responders” and “non-responders”.

4. Discussion

Data from epidemiological studies and the “Prevención con Dieta Mediterránea” or PREDIMED trial have demonstrated a reduced risk of cardiovascular disease development with increased intakes of olive oil (Buckland & Gonzalez, 2015; Estruch et al., 2013; Mozaffarian, 2016). Moreover, this trial suggested that increasing “phenolic-rich” virgin olive oil intake might be of particular benefit (Estruch et al., 2013). However, if specific phenolics are important factors in this process uniformity in metabolic and physiological responses to the intake of a specific food type should not be expected, since the content of any plant food can vary by agricultural variety, practices and manufacturing processes. Variation in phytochemical content is a prevalent issue in olive oil products, as the specific phenolics delivered within any oil are dependent on oil grade, olive variety and agricultural region grown, time of harvest, milling practices, and storage conditions (Caporaso et al., 2015; E. Karkoula et al., 2012). In particular, oleocanthal and oleacein concentrations within an EVOO are affected by olive tree varietal and inversely correlated to time of harvest (Evangelia Karkoula et al., 2014). A number of studies have observed benefit of virgin or extra virgin olive oils that are “high” in total phenols compared to low phenol oil (Covas et al., 2015), including a reduction in COX derived thromboxane and LOX derived LTB₄ production (Bogani, Galli, Villa, & Visioli, 2007). To our knowledge, this is the first human

quality EVOOs exhibit greater health benefits. However, this claim would need further verification.

Platelets are metabolically active, utilizing both anaerobic and oxidative phosphorylation for energy production, with the later predominant (Kramer, Ravi, Chacko, Johnson, & Darley-Usmar, 2014). While the current study did not access platelet specific metabolites, analysis of primary metabolism demonstrated that differences that subjects with different plasma metabolomics profiles had distinct platelet response to EVOO intake. Whether these effects were primarily due to dietary or metabolic factors is as of yet unknown. However, “responders” tended to have increased plasma glucose concentrations, along with increased levels of specific carbohydrates and sugar acids suggesting a diet rich in fruits and vegetables. Interestingly, the identification of pinitol as a significant discriminating variable of response regardless of time point suggests that responsive individuals had a higher habitual intake of soy products (Davis et al., 2000). In contrast, the “non-responsive” participants had increased circulating non-esterified fatty acids. In addition, non-responders had greater fasting levels of oleic acid, the primary fatty acid found in olive oil. Since high plasma oleic acid is a reported marker of habitual olive oil consumption that correlates with stroke prevention (Samieri et al., 2011) and EVOO effects on platelet function persist with continuous exposure (Sánchez-Muniz et al., 1998), it is possible that the EVOO effects are saturable and the non-responders in this study were already protected. That being said, the significance of these findings cannot be determined from the current study design, but warrant further exploration.

Platelet aggregation is regulated by a number of mediators, including COX-derived TXB₂, platelet 12-LOX-derived 12-HETE, and 15-LOX-derived 15-HETE (Tourdot et al., 2013). While no changes in these, and other COX and LOX-derived mediators were observed in all subjects following EVOO consumption, “responders” did demonstrate a reduction in plasma TXB₂, 11-HETE and 15-HETE concentrations following consumption of both D₂;0.5 and D₂;2, compared to D₂;0. As each of these mediators can be derived from COX-1 metabolism of arachidonic acid (Caughey, Cleland, Penglis, Gamble, & James, 2001; Tejera, Boeglin, Suzuki, & Schneider, 2012), the anti-platelet effects of the oleocanthal-containing EVOOs may in fact have a COX-1-dependent component. However, further experimentation would be needed to verify the specific mechanism by which oleocanthal affects platelet aggregation.

As platelet activation promotes the inflammatory process associated with atherosclerotic cardiovascular disease, the *in vivo* anti-inflammatory effects and mechanisms of oleocanthal and total phenolics should be explored in future studies. If true, both generally phenolic-rich oils and specifically oleocanthal-rich oils would be predicted to have positive impacts on both platelet aggregation and inflammation, and thus have benefits in cardiovascular risk reduction, as has been previously suggested in a 28-day intervention of post-menopausal women (Sánchez-Muniz et al., 1998). While the above associations are intriguing, care must be taken in their final interpretation as phenols and polyphenols are rapidly and extensively metabolized, with circulating metabolites detected in the plasma in significant levels within minutes and lasting for several hours post intake (Ottaviani et al., 2016; Zamora-Ros et al., 2016). Moreover, some studies evaluating dietary flavonoids on plasma flavanols and ex

vivo platelet function have shown increases in plasma flavonoids with no effects on platelet function (Hubbard et al., 2003). Therefore, whether the effect observed here is the consequence of direct phenolic exposure or a chemical or metabolically responsive covariate of this exposure remains to be seen.

Limitations of the current study include a small sample size, complicated by population stratification (i.e. presence of responders and non-responders), which increased variance in the data. Notably, considerable individual variability in platelet response and oxylipin production was observed with acute EVOO intake, consistent with variability in other platelet responsive factors (e.g. “aspirin resistance”) which have been postulated to result from differences in metabolism, sex, drug interactions and compliance failure (Fitzgerald & Fitzgerald, 2013; Rocca & Patrono, 2005). Dietary factors may also have played a role, and habitual diet assessments may have aided interpretation of this study. However, food frequency questionnaires are designed to provide diet assessment for study populations, not individuals, and our study design limited platelet interacting foods, obviating the need for 24 hr recalls. However, including habitual high olive oil intake as an exclusion criterion may have strengthened the study. The lack of sensitive measures for reactive oxygen species (ROS) generation (i.e. esterified F2-isoprostanes) prevented us from unequivocally ruling out potential post-prandial changes in platelet derived ROS production, such as via NADPH oxidase (Carnevale et al., 2014). The single postprandial time point of two hours post-intake was chosen based on general knowledge on polyphenol pharmacokinetics, and not specifically for oleocanthal and oleacein, which have yet to be determined, and we may have missed the maximum inhibition time, which could differ by oil. Moreover, the anti-platelet effects of the oils may become more apparent for all subjects following short-term rather than acute intake, as has been noted for aspirin.

In conclusion, this study provides evidence that the specific phenolic content within EVOO can influence platelet aggregation responses in healthy male adults. Additionally, findings suggest that the extent of the response may be influenced by individual metabolism and/or diet. Future work will be needed to determine the time course of acute effects, as well as whether or not the observed response changes with prolonged intake periods.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors gratefully acknowledge the assistance of: J. Daniel Flynn from the UC Davis Olive Center during project development; Samson I. Aghedo and Jodi L. Ensunsa from the UC Davis Department of Nutrition during study visits; Michael R. LaFrano, Ira J. Gray, and Oliver Fiehn from the West Coast Metabolomics Center and William R. Keyes from the USDA Western Human Nutrition Research Center during data acquisition; Bret Rust and Janet M. Peerson from the USDA Western Human Nutrition Research Center for statistical support. The authors also thank Corto Olive Co. (Lodi CA) for donation of the D2₂ EVOO used in the study.

Funding: This study was funded by Gaea Products S.A. (United States Department of Agriculture agreement number 58-5306-4-043F, J.W.N.); the Captain Vassilis Foundation (United States Department of Agriculture agreement number 58-5306-4-050F, J.W.N.); the United States Department of Agriculture (Project Number 2032-51530-022-00D, J.W.N.); the National Institutes of Diabetes and Digestive and Kidney Diseases (Grant Number U24DK097154-01, J.W.N.); and the National Institutes of General Medical Sciences (Grant Number T32-

GM008799, K.A.). The contents of this publication are solely the responsibility of the authors and do not necessarily represent the official views of the NIGMS, NIH, or the USDA. The USDA is an equal opportunity provider and employer.

Abbreviations

AA	arachidonic acid
ADP	adenosine diphosphate
AICc	corrected Akaike Information Criterion
BIC	Bayesian Information Criterion
COX	cyclooxygenase
D1i	Sum of an extra virgin olive oil's oleocanthal and oleacein concentrations
D2i	oleocanthal/oleacein ratio for an extra virgin olive oil
D2i0	extra virgin olive oil with undetectable oleocanthal and oleacein
D2i0.5	extra virgin olive oil with 1:2 oleacein/oleocanthal
D2i2	extra virgin olive oil with 2:1 oleacein/oleocanthal
DAD	diode array detector
EVOO	extra virgin olive oil
HETE	hydroxyeicosatetraenoic acid
LOX	lipoxygenase
PG	prostaglandin
PLS-DA	partial least squares-discriminant analysis
Pmax	maximum platelet aggregation
qNMR	quantitative ¹ H nuclear magnetic resonance spectroscopy
TXB2	thromboxane B2
UPLC	ultra-performance liquid chromatography
[COX + LOX]	changes in sum of cyclooxygenase- and lipoxygenase-derived lipid mediators

References

Agrawal K, Hassoun LA, Foolad N, Pedersen TL, Sivamani RK, Newman JW. Sweat lipid mediator profiling: a non-invasive approach for cutaneous research. *Journal of Lipid Research*. 2017; 58:188–195. [PubMed: 27875258]

- AOAC International. Peroxide Value Acetic Acid-Isooctane Method (Official Methods and Recommended Practices of the AOCS, AOCS Official Method Cd 8b-90). Urbana, IL: AOCS Press; 2013.
- Beauchamp GK, Keast RS, Morel D, Lin J, Pika J, Han Q, Breslin PA. Phytochemistry: ibuprofen-like activity in extra-virgin olive oil. *Nature*. 2005; 437:45–46. [PubMed: 16136122]
- Bendini A, Cerretani L, Carrasco-Pancorbo A, Gomez-Caravaca AM, Segura-Carretero A, Fernandez-Gutierrez A, Lercker G. Phenolic molecules in virgin olive oils: a survey of their sensory properties, health effects, antioxidant activity and analytical methods. An overview of the last decade. *Molecules*. 2007; 12:1679–1719. [PubMed: 17960082]
- Born GVR, Cross MJ. The aggregation of blood platelets. *Journal of Physiology*. 1963; 168:178–195. [PubMed: 14056485]
- Box GEP, Cox DR. An analysis of transformations. *Journal of the Royal Statistical Society Series B (Methodological)*. 1964; 26:211–252.
- Buckland G, Gonzalez CA. The role of olive oil in disease prevention: a focus on the recent epidemiological evidence from cohort studies and dietary intervention trials. *British Journal of Nutrition*. 2015; 113:S94–101. [PubMed: 26148926]
- Caporaso N, Savarese M, Paduano A, Guidone G, De Marco E, Sacchi R. Nutritional quality assessment of extra virgin olive oil from the Italian retail market: Do natural antioxidants satisfy EFSA health claims? *Journal of Food Composition and Analysis*. 2015; 40:154–162.
- Carnevale R, Pignatelli P, Nocella C, Loffredo L, Pastori D, Vicario T, Violi F. Extra virgin olive oil blunt post-prandial oxidative stress via NOX2 down-regulation. *Atherosclerosis*. 2014; 235:649–658. [PubMed: 24980290]
- Caughey GE, Cleland LG, Penglis PS, Gamble JR, James MJ. Roles of cyclooxygenase (COX)-1 and COX-2 in prostanoid production by human endothelial cells: selective up-regulation of prostacyclin synthesis by COX-2. *The Journal of Immunology*. 2001; 167:2831–2838. [PubMed: 11509629]
- Chatterjee, S., Hadi, AS. Variable selection procedures. In: Chatterjee, S., Hadi, AS., editors. *Regression analysis by example*. 5. Hoboken: John Wiley & Sons; 2015. p. 299-334.
- Covas, M-I., Fito, M., de la Torre, R. Minor bioactive olive oil components and health: Key data for their role in providing health benefits in humans. In: Bosku, D., editor. *Olive and Olive Oil Bioactive Constituents*. Urbana: AOCS Press; 2015. p. 31-52.
- Daskalaki D, Kefi G, Kotsiou K, Tasioula-Margari M. Evaluation of phenolic compounds degradation in virgin olive oil during storage and heating. *Journal of Food and Nutrition Research*. 2009; 48:31–41.
- Davis A, Christiansen M, Horowitz JF, Klein S, Hellerstein MK, Ostlund RE. Effect of pinitol treatment on insulin action in subjects with insulin resistance. *Diabetes Care*. 2000; 23:1000–1005. [PubMed: 10895854]
- EFSA Panel on Dietetic Products, Nutrition and Allergies. Scientific opinion on the substantiation of health claims related to polyphenols in olive and protection of LDL particles from oxidative damage (ID 1333, 1638, 1639, 1696, 2865), maintenance of normal blood HDL-cholesterol concentrations (ID1639), maintenance of normal blood pressure (ID 3781),"anti-inflammatory properties" (ID 1882),"contributes to the upper respiratory tract health" (ID 3468),"can help to maintain a normal function of gastrointestinal tract" (3779) and "contributes to body defences against external agents" (ID 3467) pursuant to Article 13 (1) of Regulation (EC) No 1924/2006. *EFSA Journal*. 2011; 9:2033.
- Estruch R, Ros E, Salas-Salvado J, Covas MI, Corella D, Aros F, Martinez-Gonzalez MA. Primary prevention of cardiovascular disease with a Mediterranean diet. *New England Journal of Medicine*. 2013; 368:1279–1290. [PubMed: 23432189]
- Fahrman J, Grapov D, Yang J, Hammock B, Fiehn O, Bell GI, Hara M. Systemic alterations in the metabolome of diabetic NOD mice delineate increased oxidative stress accompanied by reduced inflammation and hypertriglyceremia. *American Journal of Physiology - Endocrinology and Metabolism*. 2015; 308:E978–E989. [PubMed: 25852003]
- Fiehn, O., Wohlgemuth, G., Scholz, M. Setup and annotation of metabolomic experiments by integrating biological and mass spectrometric metadata. In: Ludäscher, B., Raschid, L., editors.

- Data Integration in the Life Sciences: Second International Workshop. Berlin, Heidelberg: Springer Berlin Heidelberg; 2005. p. 224-239.
- Fitzgerald DJ, Fitzgerald GA. Historical lessons in translational medicine: cyclooxygenase inhibition and P2Y₁₂ antagonism. *Circulation Research*. 2013; 112:174–194. [PubMed: 23287454]
- Garcia B, Coelho J, Costa M, Pinto J, Paiva-Martins F. A simple method for the determination of bioactive antioxidants in virgin olive oils. *Journal of the Science of Food and Agriculture*. 2013; 93:1727–1732. [PubMed: 23197371]
- Grapov D, Newman JW. imDEV: a graphical user interface to R multivariate analysis tools in Microsoft Excel. *Bioinformatics*. 2012; 28:2288–2290. [PubMed: 22815358]
- Grubbs FE. Sample criteria for testing outlying observations. *The Annals of Mathematical Statistics*. 1950; 21:27–58.
- Gutierrez-Rosales F, Rios JJ, Gomez-Rey ML. Main polyphenols in the bitter taste of virgin olive oil. Structural confirmation by on-line high-performance liquid chromatography electrospray ionization mass spectrometry. *Journal of Agricultural and Food Chemistry*. 2003; 51:6021–6025. [PubMed: 13129311]
- Holt RR, Heiss C, Kelm M, Keen CL. The potential of flavanol and procyanidin intake to influence age-related vascular disease. *Journal of Nutrition in Gerontology and Geriatrics*. 2012; 31:290–323. [PubMed: 22888843]
- Hubbard GP, Wolfram S, Lovegrove JA, Gibbins JM. The role of polyphenolic compounds in the diet as inhibitors of platelet function. *Proceedings of the Nutrition Society*. 2003; 62:469–478. [PubMed: 14506895]
- Hubbard GP, Wolfram S, Lovegrove JA, Gibbins JM. Ingestion of quercetin inhibits platelet aggregation and essential components of the collagen-stimulated platelet activation pathway in humans. *Journal of Thrombosis and Haemostasis*. 2004; 2:2138–2145. [PubMed: 15613018]
- Hulten LM, Olson FJ, Aberg H, Carlsson J, Karlstrom L, Boren J, Wiklund O. 15-Lipoxygenase-2 is expressed in macrophages in human carotid plaques and regulated by hypoxia-inducible factor-1 α . *European Journal of Clinical Investigation*. 2010; 40:11–17. [PubMed: 19912316]
- International Olive Oil Council. Preparation of The Fatty Acid Methyl Esters from Olive Oil and Olive-Pomace Oil (International Olive Council Testing Methods, COI/T.20/Doc. No. 24). Madrid: IOOC; 2001.
- Karkoula E, Skantzari A, Melliou E, Magiatis P. Direct measurement of oleocanthal and oleacein levels in olive oil by quantitative (1)H NMR. Establishment of a new index for the characterization of extra virgin olive oils. *Journal of Agricultural and Food Chemistry*. 2012; 60:11696–11703. [PubMed: 23116297]
- Karkoula E, Skantzari A, Melliou E, Magiatis P. Quantitative measurement of major secoiridoid derivatives in olive oil using qNMR. Proof of the artificial formation of aldehydic oleuropein and ligstroside aglycon isomers. *Journal of Agricultural and Food Chemistry*. 2014; 62:600–607. [PubMed: 24384036]
- Kramer PA, Ravi S, Chacko B, Johnson MS, Darley-Usmar VM. A review of the mitochondrial and glycolytic metabolism in human platelets and leukocytes: implications for their use as bioenergetic biomarkers. *Redox Biology*. 2014; 2:206–210. [PubMed: 24494194]
- Krishnan S, Newman JW, Hembrooke TA, Keim NL. Variation in metabolic responses to meal challenges differing in glycemic index in healthy women: Is it meaningful? *Nutrition & Metabolism*. 2012; 9:26. [PubMed: 22458475]
- La Frano MR, Fahrman JF, Grapov D, Fiehn O, Pedersen TL, Newman JW, Wedgwood S. Metabolic perturbations of postnatal growth restriction and hyperoxia-induced pulmonary hypertension in a bronchopulmonary dysplasia model. *Metabolomics*. 2017; 13:32.
- Li Z, Delaney MK, O'Brien KA, Du X. Signaling During Platelet Adhesion and Activation. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2010; 30:2341–2349.
- May AE, Seizer P, Gawaz M. Platelets: inflammatory firebugs of vascular walls. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2008; 28:S5–S10.
- Miller CH, Rice AS, Garrett K, Stein SF. Gender, race and diet affect platelet function tests in normal subjects, contributing to a high rate of abnormal results. *British Journal of Haematology*. 2014; 165:842–853. [PubMed: 24617520]

- Mozaffarian D. Dietary and policy priorities for cardiovascular disease, diabetes, and obesity: A comprehensive review. *Circulation*. 2016; 133:187–225. [PubMed: 26746178]
- Nording HM, Seizer P, Langer HF. Platelets in inflammation and atherogenesis. *Frontiers in Immunology*. 2015; 6:98. [PubMed: 25798138]
- O'Donnell VB, Murphy RC, Watson SP. Platelet lipidomics. Modern day perspective on lipid discovery and characterization in platelets. *Circulation Research*. 2014; 114:1185–1203. [PubMed: 24677238]
- Ottaviani JI, Borges G, Momma TY, Spencer JP, Keen CL, Crozier A, Schroeter H. The metabolome of [2- (14)C] (–)-epicatechin in humans: implications for the assessment of efficacy, safety, and mechanisms of action of polyphenolic bioactives. *Scientific Reports*. 2016; 6:29034. [PubMed: 27363516]
- Patrono C, Baigent C. Nonsteroidal anti-inflammatory drugs and the heart. *Circulation*. 2014; 129:907–916. [PubMed: 24566065]
- Petroni A, Blasevich M, Salami M, Papini N, Montedoro GF, Galli C. Inhibition of platelet aggregation and eicosanoid production by phenolic components of olive oil. *Thrombosis Research*. 1995; 78:151–160. [PubMed: 7482432]
- Pignatelli P, Pulcinelli FM, Lenti L, Gazzaniga PP, Violi F. Hydrogen peroxide is involved in collagen-induced platelet activation. *Blood*. 1998; 91:484–490. [PubMed: 9427701]
- Priora R, Summa D, Frosali S, Margaritis A, Di Giuseppe D, Lapucci C, Di Simplicio P. Administration of minor polar compound-enriched extra virgin olive oil decreases platelet aggregation and the plasma concentration of reduced homocysteine in rats. *Journal of Nutrition*. 2008; 138:36–41. [PubMed: 18156401]
- Rocca B, Patrono C. Determinants of the interindividual variability in response to antiplatelet drugs. *Journal of Thrombosis and Haemostasis*. 2005; 3:1597–1602. [PubMed: 16102024]
- Rondina MT, Weyrich AS, Zimmerman GA. Platelets as cellular effectors of inflammation in vascular diseases. *Circulation Research*. 2013; 112:1506–1519. [PubMed: 23704217]
- Samieri C, Féart C, Proust-Lima C, Peuchant E, Tzourio C, Stapf C, Barberger-Gateau P. Olive oil consumption, plasma oleic acid, and stroke incidence: The Three-City Study. *Neurology*. 2011; 77:418–425. [PubMed: 21676914]
- Sánchez-Muniz FJ, Oubiña P, Benedí J, Ródenas S, Cuesta C. A preliminary study on platelet aggregation in postmenopausal women consuming extra-virgin olive oil and high-oleic acid sunflower oil. *Journal of the American Oil Chemists' Society*. 1998; 75:217–233.
- Scholz M, Fiehn O. SetupX—a public study design database for metabolomic projects. *Pacific Symposium on Biocomputing*. 2007; 12:169–180.
- Servili M, Esposto S, Taticchi A, Urbani S, Di Maio I, Sordini B, Selvaggini R. Effect of diverse agricultural and technological factors on olive oil quality and yield. *Acta Horticulturae*. 2014; doi: 10.17660/ActaHortic.2014.1057.77
- Smith AB, Han Q, Breslin PAS, Beauchamp GK. Synthesis and assignment of absolute configuration of (–)-oleocanthal: a potent, naturally occurring non-steroidal anti-inflammatory and anti-oxidant agent derived from extra virgin olive oils. *Organic Letters*. 2005; 7:5075–5078. [PubMed: 16235961]
- Tejera N, Boeglin WE, Suzuki T, Schneider C. COX-2-dependent and - independent biosynthesis of dihydroxy-arachidonic acids in activated human leukocytes. *Journal of Lipid Research*. 2012; 53:87–94. [PubMed: 22068350]
- Tourdot BE, Ahmed I, Holinstat M. The emerging role of oxylipins in thrombosis and diabetes. *Frontiers in Pharmacology*. 2013; 4:176.
- van den Berg RA, Hoefsloot HC, Westerhuis JA, Smilde AK, van der Werf MJ. Centering, scaling, and transformations: improving the biological information content of metabolomics data. *BMC Genomics*. 2006; 7:142. [PubMed: 16762068]
- Vougogiannopoulou K, Lemus C, Halabalaki M, Pergola C, Werz O, Smith AB, Deguin B. One-step semisynthesis of oleacein and the determination as a 5-lipoxygenase inhibitor. *Journal of Natural Products*. 2014; 77:441–445. [PubMed: 24568174]
- Wang C, Wang W. Links between PPCA and subspace methods for complete Gaussian density estimation. *IEEE Transactions on Neural Networks*. 2006; 17:789–792. [PubMed: 16722180]

- Ward EJ. A review and comparison of four commonly used Bayesian and maximum likelihood model selection tools. *Ecological Modelling*. 2008; 211:1–10.
- Yousuf O, Bhatt DL. The evolution of antiplatelet therapy in cardiovascular disease. *Nature Reviews Cardiology*. 2011; 8:547–559. [PubMed: 21750497]
- Zamora-Ros R, Achaintre D, Rothwell JA, Rinaldi S, Assi N, Ferrari P, Scalbert A. Urinary excretions of 34 dietary polyphenols and their associations with lifestyle factors in the EPIC cohort study. *Scientific Reports*. 2016; 6:26905. [PubMed: 27273479]

Highlights

- EVOO varietal differences influence platelet aggregation and oxylipin production
- Inhibition of platelet aggregation correlated to oleocanthal content
- Inhibition of eicosanoids correlated to EVOO total phenolic content
- Basal diet may influence platelet inhibition responses to EVOOs




Fig. 1. Effects of tested EVOOs on (A) maximum platelet aggregation and (B) oxylipins associated with platelet function in healthy male subjects. D2_i2, D2_i0.5 and Ibuprofen all decreased maximum platelet aggregation compared to D2_i0, and Ibuprofen decreased oxylipin concentrations compared to all oils. Data shown are presented as means ($n = 9$) with their standard deviations. Data points with unlike letters were significantly different at $P < 0.05$ (repeated measures ANOVA).




Fig. 2. Partial Least Squares Discriminant Analysis (PLS-DA) Scores Plots showing discrimination between “responders” and “non-responders” at (A) baseline and (B) two-hours post-EVOO consumption. “Responders” are defined at subjects exhibiting >25% reduction in 1 $\mu\text{g}/\text{mL}$ collagen-stimulated P_{max} with D2i0.5 or D2i2 EVOO intake.

Table 1

Experimental EVOO characteristics. Where shown averages are means \pm SD.

	Units	Assay	D2:0	D2:2	D2:0.5
D1 Index	mg/kg	¹ H NMR	0	484	460
D2 Index	---	¹ H NMR	0	1.8	0.48
Oleocanthal	mg/kg	¹ H NMR	< 10	172 \pm 8	310 \pm 15
Oleacein	mg/kg	¹ H NMR	< 10	312 \pm 15	150 \pm 8
Tyrosol	mg/kg	¹ H NMR	189 \pm 10	< 10	< 10
DAFOA	mg/kg	¹ H NMR	< 10	< 10	< 10
DAFLA	mg/kg	¹ H NMR	< 10	< 10	< 10
AFOA	mg/kg	¹ H NMR	25 \pm 2	5 \pm 1	23 \pm 2
AFLA	mg/kg	¹ H NMR	21 \pm 2	5 \pm 1	10 \pm 1
Peroxide Values	meq/kg	PV-AcA-Iso	10.2	7.78	4.63
Total Phenols	mg caffeic /kg	F-C	212	295	214
K232 index	---	HPLC-DAD	< 2.5	< 2.5	< 2.5
K270 index	---	HPLC-DAD	< 0.22	< 0.22	< 0.22
Tyrosol ^a	% UV Detected	HPLC-DAD	19	1	3
Hydroxytyrosol ^a	% UV Detected	HPLC-DAD	13	3	5
Vanillic Acid ^a	% UV Detected	HPLC-DAD	0	3	1
p-Coumaric Acid ^a	% UV Detected	HPLC-DAD	1	5	0
Ferullic Acid ^a	% UV Detected	HPLC-DAD	0	0	0
1-Acetoxypinoreosinol ^a	% UV Detected	HPLC-DAD	5	22	8
Luteolin ^a	% UV Detected	HPLC-DAD	2	4	2
Apigenin ^a	% UV Detected	HPLC-DAD	1	1	1
C14:0	mol%	GC-FID	< 0.1	< 0.1	< 0.1
C16:0	mol%	GC-FID	12.2	14.5	15.5
C16:1	mol%	GC-FID	0.76	1.12	1.36

	Units	Assay	D2 ₀	D2 ₂	D2 _{0.5}
C17:0	mol%	GC-FID	<0.3	<0.3	<0.3
C17:1	mol%	GC-FID	<0.3	<0.3	<0.3
C18:0	mol%	GC-FID	2.65	2.48	1.97
C18:1	mol%	GC-FID	73.9	72.6	69.7
C18:2	mol%	GC-FID	9.26	6.63	7.53
C18:3	mol%	GC-FID	0.13	0.18	0.19
C20:0	mol%	GC-FID	0.46	0.43	0.4
C20:1	mol%	GC-FID	0.33	0.28	0.3
C22:0	mol%	GC-FID	<0.2	<0.2	<0.2
C24:0	mol%	GC-FID	0.15	0.12	0.11

Abbreviations: AFLA, Aldehydic Form of Ligstroside Aglycone; AFOA, Aldehydic Form of Oleuropein Aglycone; D1 Index, oleocanthal + oleacein; D2 Index, oleocanthal/oleacein ratio; D2₀, undetectable oleocanthal and oleacein; D2_{0.5}, 1:2 oleacein/oleocanthal; D2₂, 2:1 oleacein/oleocanthal; DAFLA, Dialdehydic Form of Ligstroside Aglycone.; DAFOA, Dialdehydic Form of Oleuropein Aglycone; F-C, Folin-Ciocalteu assay; PV-AcA-Iso, peroxide value acetic acid-isooctane.

^aThe internal standard normalized absorbance of the individual compound divided by the sum of normalized phenolics in all three oils (i.e. sum of %UV Detected across all oils = 100).

Table 2Baseline platelet aggregation (mean \pm SD) in collagen-stimulated plasma by study visit

Study Visit	n	1 μ g/mL collagen	3 μ g/mL collagen	<i>pa</i>
1	9	76.4 \pm 31.5	91.1 \pm 11.7	0.22
2	9	69.2 \pm 23.9	88.8 \pm 7.7	0.04
3	9	65.9 \pm 32.5	85.9 \pm 12.9	0.12
4	9	72.0 \pm 21.7	85.5 \pm 7.7	0.11
<i>pb</i>		0.35	0.49	
Inter-day variance ^c		6.31%	3.01%	

^a*P* value determined by two-tailed heteroscedastic Student's *t*-test.

^b*P* value determined by repeated measures ANOVA with Tukey's post-hoc HSD.

^cVariance determined using the average of daily average platelet aggregation measures.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3

Phenolic-dependent Pmax inhibition regression models. Models were built iteratively using all EVOO treatments; $n = 9$ participants \times 3 treatments = 27. The regression equation takes the form $P_{\max} = b_0 + b_1 x_1 + b_2 x_2 + b_3 x_3 + b_4 x_4 + b_5 x_5$. All values reported as mean \pm SE.

b_0 (Intercept)	b_1 (Oleocanthal)	b_2 (Peroxides)	b_3 (Tyrosol)	b_4 (Oleacein)	b_5 (Total Phenols)	R	AICc	BIC	P
0.13 \pm 0.10	-3.41 \pm 1.00	-	-	-	-	0.57	18.1	20.9	0.002
0.82 \pm 0.37	-5.64 \pm 1.49	-148 \pm 76	-	-	-	0.64	16.9	20.3	0.002
0.85 \pm 0.38	-6.21 \pm 1.93	-130 \pm 84.8	-0.79 \pm 1.64	-	-	0.65	19.7	23.3	0.006
1.21 \pm 0.42	-9.47 \pm 2.61	224 \pm 217	-13.4 \pm 7.4	-8.94 \pm 5.07	-	0.70	19.5	23.1	0.004
1.25 \pm 0.43	-40.3 \pm 42.9	197 \pm 223	-74.3 \pm 84.8	-85.9 \pm 107	102 \pm 141	0.71	22.5	25.7	0.008

Abbreviations: AICc, Corrected Akaike information criterion score; BIC, Bayesian information criterion score.

Table 4

Phenolic-dependent [COX + LOX] inhibition regression models. Models were built iteratively using all EVOO treatments; $n = 9$ participants $\times 3$ treatments = 27. The regression equation takes the form [COX + LOX]: $y = b_0 + b_1 x_1 + b_2 x_2 + b_3 x_3 + b_4 x_4 + b_5 x_5$. All values reported as mean \pm SE.

b_0 (Intercept)	b_1 (Total Phenols)	b_2 (Oleacein)	b_3 (Peroxides)	b_4 (Tyrosol)	b_5 (Oleocanthal)	R	AICc	BIC	P
0.93 ± 0.27	-8.71 ± 2.4	-	-	-	-	0.59	15.8	18.6	0.001
1.08 ± 0.36	-10.7 ± 3.9	1.00 ± 1.56	-	-	-	0.60	18.1	21.5	0.005
1.44 ± 0.42	-22.6 ± 8.7	5.91 ± 3.57	179 ± 118	-	-	0.65	18.6	22.2	0.005
1.39 ± 0.43	-23.7 ± 8.8	3.03 ± 4.74	355 ± 224	-4.36 ± 4.72	-	0.66	20.9	24.5	0.009
1.31 ± 0.43	-169 ± 143	113 ± 108	386 ± 226	82.8 ± 86.0	44.1 ± 43.5	0.69	23.3	26.5	0.015

Abbreviations: AICc, Corrected Akaike information criterion score; BIC, Bayesian information criterion score.

Table 5

Effect of EVOOs on COX and LOX-derived oxylipins associated with platelet function. Responders and non-responders demonstrated differential effects on oxylipin formation following consumption of EVOOs, with TXB2, 11-HETE and 15-HETE formation significantly changing following EVOO consumption in the responders. Data points with unlike letters were significantly different at $P < 0.05$ (repeated measures ANOVA). All values reported as mean \pm SD.

Oxylipin	Responders ($n = 5$)				Non-responders ($n = 4$)			
	D2i0	D2i2	D2i0.5	P^a	D2i0	D2i2	D2i0.5	P^a
TXB2	-0.04 ± 0.21^A	-0.37 ± 0.15^B	$-0.14 \pm 0.22^{A,B}$	0.02	0.10 ± 0.41	0.08 ± 0.40	0.22 ± 0.08	0.77
PGE2	-0.02 ± 0.24	-0.32 ± 0.26	-0.10 ± 0.22	0.09	0.05 ± 0.31	0.08 ± 0.40	0.28 ± 0.09	0.51
PGF2a	0.04 ± 0.19	-0.31 ± 0.45	0.13 ± 0.54	0.18	0.05 ± 0.21	-0.19 ± 0.51	0.12 ± 0.25	0.31
11-HETE	-0.02 ± 0.20^A	-0.40 ± 0.15^B	$-0.11 \pm 0.22^{A,B}$	0.02	0.11 ± 0.38	0.13 ± 0.39	0.25 ± 0.10	0.65
15-HETE	-0.02 ± 0.23^A	-0.40 ± 0.19^B	-0.10 ± 0.23^A	0.01	0.05 ± 0.44	0.11 ± 0.34	0.23 ± 0.06	0.67
12-HETE	0.01 ± 0.23	-0.47 ± 0.41	-0.20 ± 0.18	0.15	0.09 ± 0.58	0.02 ± 0.50	0.25 ± 0.05	0.68

Abbreviations: D2i0, EVOO with undetectable oleoacanthal and oleacein; D2i2, EVOO with 2:1 oleacein/oleoacanthal; D2i0.5, EVOO with 1:2 oleacein/oleoacanthal

^a Calculated using repeated measures ANOVA with Tukey's post-hoc HSD