

UCLA

UCLA Previously Published Works

Title

ImmunoChip Meta-Analysis of Inflammatory Bowel Disease Identifies Three Novel Loci and Four Novel Associations in Previously Reported Loci

Permalink

<https://escholarship.org/uc/item/7jg1n8n0>

Journal

Journal of Crohn's and Colitis, 12(6)

ISSN

1197-4982

Authors

Hong, Myunghee
Ye, Byong Duk
Yang, Suk-Kyun
[et al.](#)

Publication Date

2018-05-25

DOI

10.1093/ecco-jcc/jjy002

Peer reviewed



Original Article

ImmunoChip Meta-Analysis of Inflammatory Bowel Disease Identifies Three Novel Loci and Four Novel Associations in Previously Reported Loci

Myunghee Hong^{a,†}, Byong Duk Ye^{b,†}, Suk-Kyun Yang^{b,†}, Seulgi Jung^a,
Ho-Su Lee^a, Byoung Mok Kim^a, Soo Bin Lee^a, Jeonghoon Hong^a,
Jiwon Baek^a, Sang Hyoung Park^b, Buhm Han^c, Yi Li^d, Wenting Liu^d,
Talin Haritunians^e, Kent D Taylor^f, Jerome I Rotter^f, So-Young Bang^g,
Tae-Hwan Kim^g, Dermot P. B. McGovern^e, Jianjun Liu^d, Kyuyoung Song^a

^aDepartment of Biochemistry and Molecular Biology, University of Ulsan College of Medicine, Seoul, Korea ^bDepartment of Gastroenterology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea ^cDepartment of Convergence Medicine, University of Ulsan College of Medicine & Asan Institute for Life Sciences, Asan Medical Center, Seoul, Korea ^dHuman Genetics Group, Genome Institute of Singapore, Singapore ^eF. Widjaja Foundation Inflammatory Bowel and Immunobiology Research Institute; Cedars-Sinai Medical Center, Los Angeles, California, USA ^fInstitute for Translational Genomics and Population Sciences, Los Angeles Biomedical Research Institute and Department of Pediatrics, Harbor-UCLA Medical Center, Torrance, California, USA ^gDepartment of Rheumatology, Hanyang University Hospital for Rheumatic Diseases, Seoul, Korea

[†]Myunghee Hong, Byong Duk Ye and Suk-Kyun Yang contributed equally to this article.

Corresponding author: Kyuyoung Song, Department of Biochemistry and Molecular Biology, University of Ulsan College of Medicine, 88, Olympic-ro, 43-gil, Songpa-Gu, Seoul 05505, Korea. Tel: +82-2-3010-4277; Fax: +82-2-3010-4248; E-mail: kysong@amc.seoul.kr

Abstract

Background and Aims: Recent meta-analysis of genome-wide association studies have identified over 241 inflammatory bowel disease susceptibility loci. However, the known variants only account for a fraction of inflammatory bowel disease heritability. To identify additional susceptibility loci, we performed a trans-ethnic meta-analysis as well as an Asian-specific meta-analysis, using all published ImmunoChip association results of inflammatory bowel disease.

Methods: An inverse-variance fixed-effects meta-analysis was carried out across Korean and East Asian ImmunoChip datasets of 4156 cases and 4904 controls [Asian ancestry]. A trans-ethnic meta-analysis of inflammatory bowel disease was performed together with the European datasets of 38 155 cases and 48 485 controls genotyped on the immunoChip using a Bayesian approach, Meta-Analysis of Trans-ethnic Association studies [MANTRA].

Results: We identified seven novel associations, including three novel susceptibility loci at *MYO10-BASP1*, *PPP2R3C/KIAA0391/PSMA6/NFKB1A* and *LRRK1* as well as four novel secondary associations within previously known loci at *NCF4*, *TSPAN32*, *CIITA* and *VANGL2*. The new loci further implicate

Abbreviations: CD, Crohn's disease; CI, confidence interval; GWAS, genome-wide association study; IBD, inflammatory bowel disease; LD, linkage disequilibrium; MAF, minor-allele frequency; OR, odds ratio; SNP, single nucleotide polymorphism; UC, ulcerative colitis

alterations in B cell biology in Crohn's disease pathogenesis. The effects of five loci were universal across European and Asian ancestries, whereas the *NCF4* and *CIITA* loci showed significant heterogeneity between European and East Asian populations. In addition, 103 previously known IBD loci showed supporting evidence of association with nominal significance [$p < 0.05$] in Asians.

Conclusions: Our findings of new loci not previously associated with IBD support the importance of studying inflammatory bowel disease genetics in diverse populations.

Key Words: Inflammatory bowel disease; meta-analysis; trans-ethnic

1. Introduction

Inflammatory bowel disease [IBD] is a chronic inflammatory disorder of the gastrointestinal tract. Crohn's disease [CD] and ulcerative colitis [UC] are the two major subtypes of IBD. IBD is believed to develop due to dysregulated mucosal immune responses to gut flora in genetically susceptible individuals.¹

Although the incidence of IBD is lower in Asian populations than in Western populations, the incidence of IBD is rapidly increasing throughout Asia due to environmental changes.²⁻⁵ Epidemiological and clinical studies indicate that the phenotype and clinical course of IBD differs between Asians and Europeans.²⁻⁷ First, there is a male predominance of CD in Asia, with male to female ratios ranging from 1.67:1 to 2.9:1. Second, ileocolonic disease is predominant in Asian cases, comprising about two-thirds of CD cases, while colonic disease is only noted in ~10% of cases. It is unclear whether these Asian-specific clinical characteristics of IBD are solely due to the difference in the environments between Asia and the West, which underscores the necessity of genetic studies of IBD in Asian populations.

Recent large-scale studies on populations of European ancestry have markedly advanced our understanding of IBD-related genetics. A meta-analysis by the International IBD Genetics Consortium [IIBDGC] of genome-wide association studies [GWASs] and Immunochip data from 96 486 individuals with multiple ancestries [including Asian samples] identified >200 susceptibility loci for IBD, and reported an overlap in the directionality of the odds ratios [ORs] between the cohorts with European and Asian ancestry.⁸ The latest genome-wide meta-analysis performed on populations with European ancestry reported 241 susceptibility loci for IBD.⁹ However, it is clear that additional studies are needed to expand our understanding of the genetic architecture of IBD. Despite the differences in the clinical characteristics of IBD between different ethnicities, the number of studies in non-European populations is limited. Several GWASs on CD or UC have been performed in Asian populations.¹⁰⁻¹⁵ These studies provided insights into the genetic architecture of each disease, but not a broad view of the genetic basis of IBD in Asians. In the recent largest-to-date Asian-specific GWAS of IBD, we identified two new susceptibility loci to IBD and confirmed associations with 28 established IBD loci in Koreans.¹⁶ In the present study, we aimed to identify [i] novel genetic variants associated with IBD in Asians through a meta-analysis of Korean and East Asian studies and [ii] additional IBD susceptibility loci through a trans-ethnic meta-analysis of Korean, East Asian, and European studies. Here we present the findings of the meta-analyses, including seven novel associations identified at the level of genome-wide significance.

2. Methods

2.1. Study population

The samples of the three Immunochip datasets are described in [Supplementary Table 1](#). Samples in the Korean dataset were used

in our previous published Immunochip studies of CD and UC,^{12,13} respectively. The summary statistics of Korean IBD data can be downloaded from a webpage [<https://drive.google.com/drive/folders/1L1Zu4G0yzVuB0Ea11HkF-XQqr9pX15-0>]. Their clinical characteristics are described in [Supplementary Table 2](#). Previously published summary statistics of the East Asian and European IBD Immunochip datasets available at the IIBDGC [<https://www.ibdgenetics.org/downloads.html>] were used. The East Asian study comprised 2054 cases from Japan, 453 cases from Korea, 317 cases from China, and 3719 controls. The European dataset comprised 38 155 cases and 48 485 controls.

2.2. Quality control

The quality control [QC] procedures of the Korean, East Asian and European Immunochip datasets are described in previous publications.^{8,12,13} As Korean Immunochip studies were performed on CD and UC separately, standard QC procedures were applied using the PLINK v1.9 software [<https://www.cog-genomics.org/plink2>] as described previously for the Korean IBD dataset.^{12,13} Briefly, single nucleotide polymorphisms [SNPs] with missingness rates of >2%, a minor allele frequency [MAF] of <1%, or failing the Hardy-Weinberg equilibrium [$p < 1 \times 10^{-5}$] test were excluded. Samples with a high proportion of missing genotypes [>4%] and identity by descent [PI_HAT >0.2] were removed. QC was conducted on each dataset separately and the combined set of samples using a common approach. To control for population stratification, genotypes that passed QC filters were merged with HapMap Phase III data [194 individuals] from three populations: European [CEU], Asian [CHB + JPT], and African [YRI]. A principal-components analysis was subsequently performed on the merged dataset, and population outliers were excluded [[Supplementary Figure 1](#)]. For the meta-analysis, 87 overlapping cases [79 CD and 8 UC cases that were present in both Korean and East Asian datasets] were excluded from the Korean dataset. Finally, the Korean dataset consisted of 1332 cases and 1185 controls, with 89 051 SNPs [[Supplementary Table 3](#)].

2.3. Statistical analysis

Association tests were performed on the Korean dataset of IBD, CD and UC using a logistic regression analysis in PLINK v1.9. A quantile-quantile [Q-Q] plot was generated using R [3.2.0] [<http://www.r-project.org/>] to evaluate the overall significance of the genome-wide associations and the potential impact of population stratification. The impact of population stratification was also evaluated by calculating the genomic control inflation factor [λ_{GC}] and the genomic inflation factor for 1000 cases and 1000 controls [λ_{GC1000}] using a set of 3120 "null" SNPs that are not associated with autoimmune diseases [[Supplementary Table 1](#)]. After QC, 2117 SNPs were used as null markers to generate the Q-Q plot shown in [Supplementary Figure 2](#) [$\lambda_{GC} = 0.98$]. Both the Q-Q plots and genomic inflation factor [λ_{GC}] of Immunochip test statistics showed

that the three Immuchip association analyses had negligible inflation due to population stratification. A Manhattan plot was generated with $-\log_{10} p$ by using R [3.2.0]. For regional plots of novel associations that were identified in the Asian meta-analysis, imputation was performed using ImpG version 1.0 [https://github.com/huwenboshi/ImpG]¹⁷ and the Asian reference data [JPT + CHB] from the 1000 Genomes Project [February 2012 release] [http://www.1000genomes.org/]. To assess whether candidate novel signals were due to long-range linkage disequilibrium [LD] with variants in previously reported loci, a conditional analysis in the regions identified in trans-ethnic meta-analysis was performed. For all variants in candidate loci that were <3 Mb away from a known locus, conditional analysis was performed on each of the three datasets separately (East Asian/European dataset, GCTA [cns.genomics.com/software/gcta/]; Korean dataset, PLINK v1.9) followed by a meta-analysis, or on the combined dataset. Secondary SNPs with conditional $p < 5 \times 10^{-8}$ or $\text{Log}_{10}\text{BF} > 6$ were assumed to be independent from the reported lead SNP in the region.

2.3.1. Asian-specific meta-analysis

For the Asian-specific meta-analyses, Korean and East Asian association results were combined using the inverse-variance method under the assumption of a fixed effect as implemented in METAL.¹⁸ Between-study heterogeneity was quantified using the I^2 heterogeneity score, and statistical significance was assessed using the Q test statistic. For the fixed-effects model, significance was defined as $p_{\text{meta}} < 5 \times 10^{-8}$.

2.3.2. Trans-ethnic meta-analysis

For the trans-ethnic meta-analyses, the three independent datasets were combined using two approaches. First, a meta-analysis was performed using the METAL software assuming fixed effects across studies and using inverse-variance weighting. Second, the ethnic-specific Immuchip summary statistics were combined using the MANTRA [Meta-Analysis of Trans-ethnic Association Studies] package, a meta-analysis software tool allowing for heterogeneity in allelic effects caused by differences in LD structure in diverse populations.¹⁹ MANTRA results are reported as Log_{10} Bayes' factors [Log_{10}BF]. A $\text{Log}_{10}\text{BF} > 6$ was considered to be a genome-wide significant threshold value, and SNPs with posterior probability of heterogeneity [p_{het}] > 0.5 were interpreted as having significant heterogeneity. Both fixed effects results and MANTRA results are reported.

2.4. eQTL and bioinformatics analysis

To gain more insight into the potential functional roles of the novel IBD loci, a *cis*-eQTL analysis was performed by searching publicly available expression data generated from eQTL Blood Browser,²⁰ the Genotype-Tissue Expression [GTEx] database²¹ and Geuvadis/1000 Genomes resources.²² Whole blood, small intestine, transverse colon, and sigmoid colon were selected in the GTEx browser because they are the most important tissues in mucosal immunity. To explore epigenetic profiles of genomic locations associated with IBD, ENCODE histone modification data, HaploReg and Regulome DB were used to examine whether any of the SNPs or their proxies [$r^2 \geq 0.8$ in the 1000 genomes of JPT + CHB reference panel] were annotated as transcription factor binding or enhancer elements. Evidence of prior association signals with autoimmune diseases or other immune-related phenotypes was searched for in the Ensembl, UCSC Genome Bioinformatics, and GeneCards databases. When the SNP was not directly typed, a proxy SNP was used [$r^2 \geq 0.8$].

3. Results

3.1. Asian-specific meta-analysis

To identify additional IBD risk loci in Asians, we first performed a meta-analysis using the Korean IBD Immuchip dataset and the summary statistics from the East Asian Immuchip dataset.^{8,12,13} The combined dataset consisted of 4156 IBD cases and 4904 controls, with 1332 cases and 1185 controls from the Korean population and 2824 cases and 3719 controls from the East Asian population. Association analysis of the Korean dataset alone did not show any novel IBD loci with genome-wide significance [Supplementary Figure 3]. In a meta-analysis of the combined dataset, 12 of the previously reported regions exceeded statistical significance [$p_{\text{meta}} < 5 \times 10^{-8}$], including two novel associations in Asians at rs2072711 in *NCF4-CSF2RB* at 22q13 [OR = 1.21; 95% confidence interval [CI], 1.13 – 1.29; $p_{\text{Asian-meta}} = 5.07 \times 10^{-9}$] and at rs12928665 in *CIITA* at 16p13 [OR = 1.19; 95% CI, 1.13 – 1.25; $p_{\text{Asian-meta}} = 3.93 \times 10^{-9}$] [Supplementary Table 4, Supplementary Figure 4]. We also examined the 245 previously established European IBD-associated loci [276 independent SNPs] in the Asian meta-analysis.^{8,9} Of 193 SNPs in 174 loci [including 36 proxy SNPs with $r^2 \geq 0.8$] available, 111 SNPs from 102 loci were replicated at nominal $p < 0.05$ in association analyses for IBD, CD or UC [excluding rs864745 and rs10995235 with effects in opposite directions; Supplementary Table 5]. An additional locus [JAK2], not in LD with the European SNP, reached a genome-wide significant association in the Asian meta-analysis [Supplementary Table 4], resulting 112 SNPs being replicated at 103 loci in Asians.

3.2. Trans-ethnic meta-analysis

We performed a trans-ethnic meta-analysis of IBD using three independent Immuchip datasets. The combined dataset consisted of 42 311 IBD cases and 53 389 controls, with 1332 cases and 1185 controls from Korean population, 2824 cases and 3719 controls from East Asian population, and 38 155 cases and 48 485 controls from the European population. After applying stringent quality controls, we tested 89 051, 106 681, 126 098 and 80 291 SNPs for association in the Korean, East Asian, European and combined cohort, respectively [Supplementary Table 1]. We performed a meta-analysis using [i] inverse-variance based meta-analysis in METAL software and [ii] the MANTRA package [Supplementary Figure 5]. Following meta-analyses, the SNPs within 241 known loci⁹ in the MHC region [25–34 Mb, hg19] and known genes were removed.^{9,23} Twenty-two loci showed significant associations with IBD in the MANTRA analyses [$\text{Log}_{10}\text{BF} > 6$], of which 19 loci also showed significant associations in the fixed-effects trans-ethnic METAL analyses [$p < 5 \times 10^{-8}$, Supplementary Table 6]. As these loci were <3 Mb away from known loci except for rs2624435 near *MYO10-BASP1*, we performed a conditional analysis for 21 variants [Supplementary Table 7]. Only three loci [*CIITA*, *TSPAN32* and *NCF4*] showed genome-wide significant associations following conditional analyses on the reported European lead SNPs, suggesting the presence of three novel secondary associations within previously reported loci [Table 1]. The MANTRA analyses of CD showed eight loci with $\text{Log}_{10}\text{BF} > 6$ [Supplementary Table 6]. Only two of these loci, rs57275892 [*PSMA6-NFKBIA*] and rs7170683 [*LRRK1*], were novel [Table 1]. Of the three loci with $\text{Log}_{10}\text{BF} > 6$ in the MANTRA analyses of UC [Supplementary Table 6], rs17371986 [*VANGL2*] showed an independent association within a previously known locus following conditional analysis [Supplementary Table 7]. Of seven novel associations, only rs12928665 [*CIITA*] and rs2072711

Table 1. Seven novel associations identified through Asian and trans-ethnic meta-analysis on inflammatory bowel disease.

Chr	SNP	Position [hg19]	Effect allele	Study	Effect allele frequency	Inflammatory bowel disease		p	p _{random}	pQ			
						Controls [N]	Cases [N]				OR	[95% CI]	
22 [NCF4, CSF2RB]	rs2072711	37 268 555	A	Korean	0.32	1 185	1 332	1.21		0.002			
				East Asian	0.26	3 719	2 824	1.21	[1.07-1.36]		6.55 × 10 ⁻⁷		
				European	0.17	48 485	38 155	1.00	[1.13-1.28]		0.954		
				Combined1[†]		4 904	4 156	1.21	[0.97-1.03]		5.07 × 10^{-9**}	5.07 × 10^{-9**}	0.97
				Combined2[‡]		53 389	42 311	1.03	[1.13-1.29]		0.027	0.121	4.42 × 10⁻⁷
				Korean	0.43	1 185	1 332	0.86	[1.00-1.05]		0.008		
16 [CIITA]	rs12928665	10 971 474	A	East Asian	0.46	3 719	2 824	0.83	[0.75-0.97]		1.29 × 10 ⁻⁷		
				European	0.76	48 485	38 155	0.94	[0.76-0.90]		1.29 × 10 ⁻⁸		
				Combined1[†]		4 904	4 156	0.84	[0.91-0.96]		3.93 × 10^{-9**}	3.93 × 10^{-9**}	0.61
				Combined2[‡]		53 389	42 311	0.92	[0.79-0.89]		1.23 × 10^{-13**}	0.006	0.002
				Korean	0.20	1 185	1 332	0.88	[0.90-0.94]		0.061		
				East Asian	0.25	3 719	2 824	0.87	[0.74-1.01]		8.45 × 10 ⁻⁴		
5 [MYO10, BASP1]	rs2624435	17 095 269	A	European	0.40	48 485	38 155	0.95	[0.79-0.95]		1.71 × 10 ⁻⁷		
				Combined1[†]		4 904	4 156	0.87	[0.93-0.97]		1.29 × 10⁻⁴	1.29 × 10⁻⁴	0.92
				Combined2[‡]		53 389	42 311	0.94	[0.81-0.94]		1.05 × 10^{-9**}	0.005	0.09
				Korean	0.14	1 185	1 332	0.92	[0.92-0.96]		0.287		
				East Asian	0.14	3 719	2 824	0.89	[0.76-1.08]		0.035		
				European	0.23	48 485	38 155	0.94	[0.79-1.00]		1.93 × 10 ⁻⁷		
Combined1[†]		4 904	4 156	0.90	[0.92-0.96]		0.019	0.019	0.79				
Combined2[‡]		53 389	42 311	0.94	[0.83-0.98]		1.80 × 10^{-8**}	1.80 × 10^{-8**}	0.62				

Table 1. Continued

Chr	SNP	Position [hg19]	Effect allele	Study	Effect allele frequency	Inflammatory bowel disease			<i>p</i>	<i>p</i> _{random}	<i>p</i> <i>Q</i>
						Controls [N]	Cases [N]	OR [95% CI]			
15	rs7170683 [LRRK1]	101 488 582	A	Korean	0.48	1 185	1 332	0.98	0.748		
				East Asian	0.47	3 719	2 824	[0.87–1.09]	0.673		
				European	0.52	48 485	38 155	[0.91–1.06]	1.98 × 10 ⁻⁴		
				Combined1 [†]		4 904	4 156	[0.94–0.98]	0.597	0.597	0.97
				Combined2 [‡]		53 389	42 311	[0.93–1.05]	2.11 × 10 ⁻⁴	2.11 × 10 ⁻⁴	0.81
						1 185	1 332	[0.95–0.98]	1.98 × 10 ⁻⁴		
14	rs57275892 [PPP2R3C, KIAA0391, PSM A6, NFKBIA]	35 747 271	A	Korean	0.35	1 185	1 332	0.80	0.025		
				East Asian	0.33	3 719	2 824	[0.69–0.92]	1.50 × 10 ⁻⁴		
				European	0.17	48 485	38 155	[0.84–0.99]	9.25 × 10 ⁻⁵	0.029	0.06
				Combined1 [†]		4 904	4 156	[0.92–0.98]	5.55 × 10 ⁻⁷	0.013	0.02
				Combined2 [‡]		53 389	42 311	[0.83–0.94]			
						1 185	1 332	[0.92–0.96]			
1	rs17371986 [VANG L2]	160 402 259	A	Korean	0.83	1 185	1 332	0.88	0.080		
				East Asian	0.79	3 719	2 824	[0.73–1.02]	0.205		
				European	0.79	48 485	38 155	[0.86–1.03]	5.18 × 10 ⁻⁵		
				Combined1 [†]		4 904	4 156	[0.93–0.98]	0.047	0.047	0.39
				Combined2 [‡]		53 389	42 311	[0.86–1.00]	8.27 × 10 ⁻⁶	8.27 × 10 ⁻⁶	0.55
						1 185	1 332	[0.93–0.97]			

Table 1. Continued

Chr	Crohn's disease						Ulcerative colitis					
	Cases	OR	<i>p</i>	<i>P</i> _{random}	<i>pQ</i>	<i>pQ</i>	Cases	OR	<i>p</i>	<i>P</i> _{random}	<i>pQ</i>	
	(N)	(95% CI)					[N]	[95% CI]				
22 [NCF4, CSF2RB]	637	1.29 [1.11-1.49]	6.95 × 10 ⁻⁴				695	1.14 [0.98-1.31]	0.082			
	1 690	1.26 [1.17-1.35]	2.17 × 10 ⁻⁷				1 134	1.15 [1.04-1.26]	0.009			
	20 550	0.96 [0.92-0.99]	0.008				17 647	1.05 [1.01-1.08]	0.008			
	2 327	1.27 [1.18-1.37]	6.01 × 10 ^{-10**}	6.01 × 10 ^{-10**}	0.83		1 829	1.15 [1.05-1.25]	0.002	0.002	0.89	
	22 877	1.00 [0.97-1.03]	0.986	0.218	1.31 × 10 ⁻¹⁰		19 476	1.06 [1.03-1.09]	3.20 × 10 ⁻⁴	0.017	0.15	
	637	0.90 [0.76-1.04]	0.140				695	0.82 [0.69-0.96]	0.004			
16 [CIITA]	1 690	0.81 [0.73-0.90]	1.07 × 10 ⁻⁶				1 134	0.85 [0.76-0.95]	8.92 × 10 ⁻⁴			
	20 550	0.94 [0.91-0.97]	6.61 × 10 ⁻⁶				17 647	0.94 [0.91-0.97]	2.36 × 10 ⁻⁵			
	2 327	0.84 [0.78-0.90]	7.53 × 10 ⁻⁷	3.48 × 10 ⁻⁴	0.22		1 829	0.84 [0.78-0.91]	1.38 × 10 ⁻⁵	1.38 × 10 ⁻⁵	0.64	
	22 877	0.93 [0.90-0.95]	2.22 × 10 ^{-9**}	0.020	0.005		19 476	0.93 [0.90-0.95]	4.50 × 10 ^{-8**}	0.007	0.03	
	637	0.82 [0.65-0.99]	0.026				695	0.93 [0.77-1.09]	0.361			
	1 690	0.87 [0.77-0.96]	0.004				1 134	0.88 [0.77-0.98]	0.017			
11 [TSPAN32]	20 550	0.95 [0.92-0.97]	1.04 × 10 ⁻⁵				17 647	0.94 [0.92-0.97]	1.73 × 10 ⁻⁵			
	2 327	0.86 [0.79-0.93]	2.94 × 10 ⁻⁴	2.94 × 10 ⁻⁴	0.58		1 829	0.89 [0.81-0.98]	0.013	0.013	0.56	
	22 877	0.94 [0.92-0.96]	4.18 × 10 ⁻⁷	0.012	0.07		19 476	0.94 [0.92-0.96]	1.44 × 10 ⁻⁶	1.44 × 10 ⁻⁶	0.41	
	637	1.05 [0.86-1.24]	0.628				695	0.80 [0.61-1.00]	0.028			
	1 690	0.90 [0.78-1.02]	0.099				1 134	0.89 [0.74-1.03]	0.093			
	20 550	0.96 [0.93-0.99]	0.003				17 647	0.93 [0.91-0.96]	3.64 × 10 ⁻⁶			
5 [MYO10, BASP1]	2 327	0.94 [0.85-1.04]	0.262	0.525	0.19		1 829	0.86 [0.76-0.96]	0.008	0.008	0.42	
	22 877	0.96 [0.93-0.98]	0.001	0.001	0.41		19 476	0.93 [0.90-0.96]	2.80 × 10 ⁻⁷	0.005	0.26	

Table 1. Continued

Chr	Crohn's disease						Ulcerative colitis					
	Cases	OR	<i>p</i>	<i>P</i> _{random}	<i>pQ</i>	<i>pQ</i>	Cases	OR	<i>p</i>	<i>P</i> _{random}	<i>pQ</i>	
	(N)	(95% CI)					[N]	[95% CI]				
15	637	0.86	0.027				695	1.11	0.117			
[<i>LRKK1</i>]	1 690	[0.72-0.99]	0.073				1 134	[0.98-1.25]	0.097			
	20 550	[0.84-1.01]	4.97×10^{-7}				17 647	[0.99-1.18]	0.212			
	2 327	[0.92-0.96]	0.007	0.007	0.34		1 829	[0.96-1.01]	0.024	0.024	0.76	
	22 877	[0.84-0.97]	$1.99 \times 10^{-8**}$	$1.99 \times 10^{-8**}$	0.39		19 476	[1.01-1.18]	0.608	0.608	0.04	
		0.94						[0.97-1.02]				
14	637	0.82	0.006				695	0.79	7.59×10^{-4}			
[<i>PPP2R3C</i> , <i>KIAA0391</i> , <i>P5MA6</i> , <i>NFKB1A</i>]	1 690	[0.67-0.96]	0.014				1 134	[0.65-0.93]	0.399			
	20 550	[0.81-0.98]	4.89×10^{-6}				17 647	[0.86-1.06]	0.355			
	2 327	0.93						0.98				
	22 877	[0.90-0.96]	4.34×10^{-4}	7.98×10^{-4}	0.30		1 829	[0.95-1.02]	0.008	0.167	0.03	
		0.87						0.90				
		[0.81-0.94]	$2.38 \times 10^{-8**}$	3.91×10^{-4}	0.20		19 476	[0.82-0.97]	0.064	0.145	0.01	
		0.92						0.97				
1	637	[0.89-0.95]	0.046				695	[0.94-1.00]	0.341			
[<i>VANGL2</i>]	1 690	0.83					1 134	0.92	0.037			
	20 550	[0.66-1.01]	0.913				17 647	[0.74-1.10]	1.95×10^{-8}			
	2 327	[0.89-1.10]	0.576				1 829	[0.77-1.00]	0.023	0.023	0.74	
	22 877	0.99						0.92				
		[0.96-1.02]	0.276	0.368	0.09			[0.89-0.95]				
		0.95						0.89				
		[0.87-1.04]	0.384	0.414	0.17		19 476	[0.81-0.98]	$1.58 \times 10^{-9**}$	$1.58 \times 10^{-9**}$	0.83	
		0.99						0.92				
		[0.96-1.02]						[0.89-0.94]				

Chr, chromosome; CI, confidence interval; OR, odds ratio; Position, chromosome position. Genome-wide significant results from the combined analysis are indicated in bold.

**Association results at the genome-wide significance level [$p < 5.0 \times 10^{-8}$].

[†]Combined1: Korean+East Asian dataset.

[‡]Combined2: Korean+East Asian+European dataset.

[NCF4] showed evidence of heterogeneity between Asian and European populations [$p_{\text{het}} \geq 0.91$; Table 1, Supplementary Table 6].

3.3. Novel loci

All three newly discovered SNPs are non-coding variants, rs2624435 in *MYO10-BASP1* at 5p15 for IBD, rs57275892 in *PPP2R3C/KIAA0391/PSMA6/NFKB1A* at 14q13, and rs7170683 in *LRRK1*

at 15q26 for CD. The SNP densities across the *MYO10-BASP1* and *LRRK1* loci were sparse [Figure 1A, 1B]; however, upon examining the eQTL database, rs2624435 showed an association with *BASP1* expression levels in blood, and rs7170683 showed an association with both *LRRK1* and *ALDH1A3* expression levels in lymphoblastoid cell lines of European ancestry [Supplementary Table 8]. A novel IBD susceptibility locus at rs2624435 is located between *MYO10*




Figure 1. Signal plots for the three novel IBD loci: [A] rs2624435 at 5p15, [B] rs7170683 at 15q26 and [C] rs57275892 at 14q13. The left panels represent signal plots from the fixed-effects meta-analyses of Korean and East Asian-ancestry individuals. SNPs are plotted according to their chromosomal positions [NCBI Build 37] with $-\log_{10} p$ values from the Asian meta-analysis in the region flanking 750 kb on either side of the marker SNP. The circles indicate the genotyped SNPs, and the squares indicate the imputed SNPs. The most strongly associated SNP in the discovery stage is shown as a small purple circle. Linkage disequilibrium [LD; r^2 values] between the lead SNP and the other SNPs are indicated using colours. The relative location of the annotated genes and the direction of transcription are shown in the lower portion of the figure. The estimated recombination rates of the Asian samples from the 1000 Genomes Project [Nov 2014] are plotted to reflect the local LD structure. Plots are generated using LocusZoom. The right panel presents signal plots for the MANTRA association signal after trans-ethnic meta-analysis of Korean, East-Asian, and European Immunochip data. Each point represents an SNP that passed the QC in the MANTRA analysis, plotted with their BF [on a \log_{10} scale] as a function of genomic position [NCBI Build 37]. Plots were generated using R.




Figure 1. Continued

and *BASP1*. The biological function of the Brain Acid Soluble Protein 1 [*BASP1*] gene is not well known, except for its involvement in nephrogenesis as a transcriptional co-suppressor,²⁴ whereas an unconventional myosin, Myosin X [*MYO10*], is implicated in phagocytosis via its role in filopodia induction. Macrophages lacking Myo10 showed markedly reduced filopodia formation.²⁵ A novel CD susceptibility locus at rs7170683 is located in the Leucine-rich repeat kinase 1 [*LRRK1*]. *LRRK1* belongs to a member of the ROCO family of proteins with multiple functional domains including ankyrin-like repeats, leucine-rich repeats [LRRs], a Ras-like GTPase domain [ROC] and an adjacent C-terminal domain [COR], and a serine-threonine kinase domain. Its homolog *LRRK2* was shown to be associated with CD²³ and Parkinson's disease.²⁶ *LRRK1* is involved in a variety of functions including autophagy and osteoclast differentiation. A recent knock-out mice study reported that *LRRK1* plays a critical role in B cell development and antibody production by regulating NF- κ B signaling.²⁷ The other novel CD susceptibility SNP [rs57275892] at 14q13 is located ~500 bp upstream of *PSMA6* [Figure 1C] in a LD region of 645 kb [35.212–35.857 Mb] that includes *BAZ1A*, *SRP54*, *FAM177A1*, *PPP2R3C*, *KIAA0391*, *PSMA6* and *NFKB1A*. Although rs57275892 is located ~500 bp upstream of *PSMA6*, eQTL analyses showed that it was associated most significantly with the expression of a nearby gene, protein phosphatase 2 regulatory subunit B' gamma [*PPP2R3C*] [$p = 2.00 \times 10^{-22}$] in whole blood [Supplementary Table 8]. *PPP2R3C* is a regulatory subunit of a serine/threonine phosphatase, protein phosphatase 2. Knock-out and transgenic mice studies show that *PPP2R3C* is involved in both the survival of germinal center B cells and the differentiation of peritoneal B cells into autoantibody-producing plasma cells, suggesting possible roles in inflammatory diseases.²⁸

3.4. Novel secondary associations within previously known loci

The association signal [rs2074023] at chromosome 11p15 is located within intron 1 of Tetraspanin32 [*TSPAN32*, $\text{Log}_{10}\text{BF} = 7.49$, $p_{\text{meta}} = 1.05 \times 10^{-9}$] in a LD region of 117.8 kb that includes

C11orf21, *TSPAN32*, *CD81* and *TSSC4* [Table 1, Supplementary Figure 6]. rs2074023 is located ~452 kb away from the previously reported rs907611^{8,9} [$r^2 < 0.2$]; however, conditional logistic regression analysis on rs907611 supported the independent effects of the two SNPs [Supplementary Table 7]. Examination of the eQTL database for rs2074023 showed that it is associated with the expression levels of both *C11orf21* and *TSPAN32* in lymphoblastoid cell lines and small intestine, whereas previously reported rs907611 is associated with the expression levels of *CTSD* in blood [Supplementary Table 8], supporting their independent effects. A search of HaploReg v4 for rs2074023 showed a Regulome DB score 1b, indicating that it is likely to affect binding and to be linked to expression of a gene target [Supplementary Table 9]. *TSPAN32* is a member of the tetraspanins of integral membrane proteins with functional roles in cell motility, membrane fusion, proliferation and immunity. Tetraspanins, of which over 30 have been identified in humans, can associate with one another and with other molecules such as integrins or proteins of the immunoglobulin superfamily to form a network on the surfaces of many different cell types.²⁹ Knock-out mice experiments have shown that *TSPAN32* may play a role in the negative regulation of peripheral T-lymphocyte proliferation,³⁰ suggesting that it may have roles in inflammatory diseases. The biological function of the other gene, *C11orf21*, is unknown; however, rs7944004, located in 5.7 kb 3' of *C11orf21* [14.4 kb away from rs2074023 with $r^2 = 0.75$], is associated with chronic lymphocytic leukemia.³¹ The association signal [rs17371986] at chromosome 1q23 is located at 3.8 kb from the 3' end of *VANGL* planar cell polarity protein 2 gene [*VANGL2*] in a LD region of 110.7 kb that includes *NCSN*, *NHLH1* and *VANGL2* [Table 1, Supplementary Figure 6]. The previously reported rs4656958 [~2 kb 5' of *ITLN1*] is located ~454 kb away from rs17371986 [$r^2 < 0.2$]; however, conditional logistic regression analysis of rs4656958 supported the independent effects of the two SNPs [Supplementary Table 7]. Examination of the eQTL database for rs17371986 showed that it is associated with *VANGL2* expression levels in the sigmoid colon in the GTEx database [Supplementary Table 8]. As *VANGL2* is

involved in the regulation of planar cell polarity, its involvement in UC susceptibility is not obvious.

3.5. Loci showing heterogeneity between Asian and European populations

Of the two novel IBD associations that showed significant heterogeneity between Asian and European populations, association effects of *NCF4* were mapped to two independent SNPs in two populations [$p_{\text{het}} = 1.0$, [Supplementary Table 6](#) and [Supplementary Figure 6](#)]. The top SNP rs2072711 identified in the Asian-specific meta-analyses showed much stronger association with CD than with UC, whereas the recently reported European top SNP rs4821544 [10 kb away from rs2072711, $r^2 < 0.2$ in ASN] showed a significant association with CD [OR = 1.07, $p = 6.75 \times 10^{-8}$] only [[Supplementary Table 10](#)]. rs2072711 showed weak associations with CD and UC in the European dataset with different effect directions, but rs4821544 failed to show associations with IBD in the Asian samples. The conditional analysis on rs4821544 did not abolish the association at rs2072711 in East Asian [$p = 6.55 \times 10^{-7}$, $p_{\text{condition}} = 6.15 \times 10^{-7}$] and Korean samples [$p = 2.16 \times 10^{-3}$, $p_{\text{condition}} = 1.95 \times 10^{-3}$], suggesting two independent associations in the *NCF4* locus. rs2072711 is located on chromosome 22q12 in a LD region of ~60.3 kb, which includes two genes, *NCF4* [neutrophil cytosolic factor 4] and *CSF2RB* [colony stimulating factor 2 receptor beta common subunit]. Examination of eQTL databases for rs2072711 showed that it had a much stronger association with the mRNA expression levels of *NCF4* [$p = 6.20 \times 10^{-15}$ in lymphoblastoid, $p = 4.35 \times 10^{-55}$ in blood] than with those of *CSF2RB* [$p = 1.10 \times 10^{-5}$ in lymphoblastoid] [[Supplementary Table 8](#)].^{20,32,33} *NCF4* encodes a cytosolic regulatory component of superoxide-producing phagocyte NADPH oxidase, a multicomponent enzyme system important for host defense. The risk alleles for both rs4821544 and rs2072711 were associated with decreased expression of *NCF4* in the blood eQTL database, $p = 3.24 \times 10^{-28}$ and $p = 4.35 \times 10^{-55}$, respectively.²⁰

rs12928665 in intron 1 of the MHC class II transactivator gene [*CIITA*] at chromosome 16p13 showed significant heterogeneity between Asian and European populations [$\text{Log}_{10} \text{BF} = 7.76$, $p_{\text{het}} = 1.00$; $p_{\text{trans-ethnic meta}} = 1.23 \times 10^{-13}$, $p_{\text{Q}} = 2.14 \times 10^{-3}$] [[Supplementary Table 6](#) and [Supplementary Figure 6](#)]. Of note is the significant difference in frequency of rs12928665 between Asian and European populations [[Supplementary Table 11](#)]. Previous studies have reported two independent association signals, rs529866 in the *SOCS1* locus^{8,9} [401.8 kb away from rs12928665] for CD and rs11641184 in the *LITAF* locus⁸ [733.1 kb from rs12928665] for IBD. The conditional analysis on rs529866, rs11641184 and both did not abolish the association at rs12928665, suggesting three independent associations in the *CIITA-SOCS1-LITAF* locus [[Supplementary Table 11](#)]. Previously, rs4781011 of *CIITA* [intron 2, 3838 bp away from rs12928665] was reported to be associated with UC in Europeans, but not to be present at 200 IBD susceptibility loci or present at the level of genome-wide significance.³⁴ LD between rs12928665 and rs4781011 in Asians [JPT + CHB] was low [$r^2 < 0.2$], whereas it was almost complete in Europeans [$r^2 = 0.95$]. Indeed, association of rs12928665 with IBD [$p = 1.29 \times 10^{-8}$] was stronger than that of rs4781011 in the European dataset [$p = 4.21 \times 10^{-8}$], suggesting that the association previously reported might have been due to an indirect association caused by the LD between the two SNPs. Examination of the eQTL database for the three SNPs showed that they were associated with the expression levels of different genes in blood, indicating that their effects are independent [[Supplementary Table 8](#)]. *CIITA* is an important transcription factor for the expression of HLA class II molecules and is involved in the expression of HLA class I molecules.^{35,36}

4. Discussion

Here we present the largest ImmunoChip meta-analysis of IBD, using all published ImmunoChip association results of IBD. Despite the




Figure 2. Comparison of odds ratios [ORs] for the 193 reported and 7 newly identified Crohn's disease and ulcerative colitis risk variants in Europeans and East Asians [based on the minor allele in Europeans]. The 193 reported SNPs are listed in [Supplementary Table 5](#) and 7 newly identified SNPs in [Table 1](#). Each dot represents the ORs [on a log scale] for each SNP in [A] Crohn's disease and [B] ulcerative colitis. Colour denotes the range of association p values for Crohn's disease or ulcerative colitis in the discovery dataset. The red line refers to linear regression, which was weighted by the inverse of the variance of the log[ORs] in the discovery dataset. The correlation coefficients and p values of the linear model are shown at the bottom-right corner.

limitation of using only 80 291 common SNPs for the analyses, we identified seven novel associations, including three novel susceptibility loci at *MYO10-BASP1* for IBD, *PPP2R3C/KIAA0391/PSMA6/NFKB1A* and *LRRK1* for CD, and four novel secondary associations within previously known loci at *NCF4*, *TSPAN32*, *CIITA* and *VANGL2*. In addition, we were able to replicate 103 loci of known IBD susceptibility loci in Asian samples [Supplementary Table 5]. The comparison of the effect sizes for all 193 SNPs from known loci and seven novel associations identified in the present study between Asians and Europeans revealed a positive correlation in the direction of effects for both CD and UC [Figure 2] [$r^2 = 0.40$ and $p = 1.54 \times 10^{-23}$ for CD; $r^2 = 0.57$ and $p = 1.84 \times 10^{-37}$ for UC]. This observation is consistent with a previous large-scale study that reported substantial genetic overlap between Europeans and Asians.⁸

Of the two novel IBD associations showing significant heterogeneity between Asian and European populations, the association effects of *NCF4* were mapped to two independent SNPs in two populations. The fact that the risk alleles for both SNPs were associated with decreased expression of *NCF4* in blood eQTL database suggests that the heterogeneity of the most significant signals might be due to the LD difference between Asians and Europeans [Supplementary Figure 7]. The *NCF4-CSF2RB* region showed genome-wide significant associations with both IBD and CD in Asians, but with CD only in Europeans. Previous studies have shown association of rs4821544 with ileal CD in Europeans^{37–39} and its effects on reactive oxygen species production following stimulation with GM-CSF.⁴⁰ Stronger association of *NCF4* in Asians could be due to the fact that ileocolonic disease is the most common type of CD in Asian populations, whereas ileal, colonic and ileocolonic disease occur in equal proportions in the CD of western populations.

In this study, we conducted the largest Immunochip trans-ethnic meta-analysis of IBD and discovered seven novel associations, including three novel susceptibility loci and four novel independent associations within previously known loci. The new loci suggest that additional factors are involved in the T cell biology and B cell immunity of IBD. In conclusion, this trans-ethnic study advances our understanding of the genetic architecture of IBD susceptibility by discovering novel associations and revealing allelic heterogeneity between Asian and European populations.

Web Resources

The URLs for data presented herein are as follows:

METAL, <http://csg.sph.umich.edu/abecasis/metal/>

The 1000 Genome Project, <http://www.1000genomes.org/>

UCSC Genome Browser, <http://genome.ucsc.edu/>

GCTA, <http://cns.genomics.com/software/gcta/>

IIBDGC, <https://www.ibdgenetics.org/>

RegulomeDB v2, <http://www.broadinstitute.org/mammals/haploreg/haploreg.php>

Genotype–Tissue Expression [GTEx] project, <http://www.gtexportal.org/home>

eQTL Blood Browser, <http://www.genenetwork.nl/loodeqtlbrowser/>

Geuvadis/1000 Genomes resources, <http://www.ebi.ac.uk/Tools/geuvadis-das/>

Funding

This work was supported by a Korean Health Technology R&D Project grant through the Korea Health Industry Development Institute to S-KY [A120176], funded by the Ministry of Health & Welfare, as well as a Mid-career Researcher Program grant through the National Research Foundation of Korea to KS [2014R1A2A1A09005824, 2017R1A2A1A05001119], funded by the Ministry of Science, Information & Communication Technology and Future Planning, the Republic of Korea. MH was supported by Basic Science

Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education [NRF-2016R1A6A3A01013069].

Conflict of Interest

None declared.

Acknowledgments

We would like to thank all the participating patients and healthy donors who provided the DNA and clinical information for this study. This work was supported by the PLSI supercomputing resources of the Korea Institute of Science and Technology Information. Ethics approval was given by the Institutional Review Board of the Asan Medical Center.

Author Contributions

KS and S-KY obtained financial support. KS conceived and designed the study and supervised the data analysis and interpretation. BDY and BH participated in the study design and supervised the data analysis and interpretation. S-KY, BDY, SHP and HSL recruited subjects and participated in the diagnostic evaluation. TH, KDT, JIR, JL and DPBM, participated in genotyping and supervised the data analysis. SYB and T-HK provided additional control data. MH, SJ, JH, JB, WL, YL, BMK and SBL performed data analyses. KS and MH drafted the manuscript. KS revised the manuscript.

Supplementary Data

Supplementary data are available at *ECCO-JCC* online.

References

1. Khor B, Gardet A, Xavier RJ. Genetics and pathogenesis of inflammatory bowel disease. *Nature* 2011;474:307–17.
2. Thia KT, Loftus EV Jr, Sandborn WJ, Yang SK. An update on the epidemiology of inflammatory bowel disease in Asia. *Am J Gastroenterol* 2008;103:3167–82.
3. Yang SK, Yun S, Kim JH, et al. Epidemiology of inflammatory bowel disease in the Songpa–Kangdong district, Seoul, Korea, 1986–2005: a KASID study. *Inflamm Bowel Dis* 2008;14:542–9.
4. Ng SC, Tang W, Ching JY, et al. Incidence and phenotype of inflammatory bowel disease based on results from the Asia–Pacific Crohn’s and colitis epidemiology study. *Gastroenterology* 2013;145:158–62.e2.
5. Park SJ, Kim WH, Cheon JH. Clinical characteristics and treatment of inflammatory bowel disease: a comparison of Eastern and Western perspectives. *World J Gastroenterol* 2014;20:11525–37.
6. Burisch J, Pedersen N, Čuković-Čavka S, et al.; EpiCom-group. East–West gradient in the incidence of inflammatory bowel disease in Europe: the ECCO-EpiCom inception cohort. *Gut* 2014;63:588–97.
7. Cosnes J, Gower-Rousseau C, Seksik P, Cortot A. Epidemiology and natural history of inflammatory bowel diseases. *Gastroenterology* 2011;140:1785–94.
8. Liu JZ, van Sommeren S, Huang H, et al.; International Multiple Sclerosis Genetics Consortium; International IBD Genetics Consortium. Association analyses identify 38 susceptibility loci for inflammatory bowel disease and highlight shared genetic risk across populations. *Nat Genet* 2015;47:979–86.
9. de Lange KM, Moutsianas L, Lee JC, et al. Genome-wide association study implicates immune activation of multiple integrin genes in inflammatory bowel disease. *Nat Genet* 2017;49:256–61.
10. Yang SK, Hong M, Zhao W, et al. Genome-wide association study of ulcerative colitis in Koreans suggests extensive overlapping of genetic susceptibility with Caucasians. *Inflamm Bowel Dis* 2013;19:954–66.
11. Yang SK, Hong M, Zhao W, et al. Genome-wide association study of Crohn’s disease in Koreans revealed three new susceptibility loci and common attributes of genetic susceptibility across ethnic populations. *Gut* 2014;63:80–7.

12. Yang SK, Hong M, Choi H, *et al.* Immunochip analysis identification of 6 additional susceptibility loci for Crohn's disease in Koreans. *Inflamm Bowel Dis* 2015;21:1–7.
13. Ye BD, Choi H, Hong M, *et al.* Identification of ten additional susceptibility loci for ulcerative colitis through immunochip analysis in Koreans. *Inflamm Bowel Dis* 2016;22:13–9.
14. Asano K, Matsushita T, Umeno J, *et al.* A genome-wide association study identifies three new susceptibility loci for ulcerative colitis in the Japanese population. *Nat Genet* 2009;41:1325–9.
15. Yamazaki K, Umeno J, Takahashi A, *et al.* A genome-wide association study identifies 2 susceptibility loci for Crohn's disease in a Japanese population. *Gastroenterology* 2013;144:781–8.
16. Yang SK, Hong M, Oh H, *et al.* Identification of loci at 1q21 and 16q23 that affect susceptibility to inflammatory bowel disease in Koreans. *Gastroenterology* 2016;151:1096–9.e4.
17. Pasaniuc B, Zaitlen N, Shi H, *et al.* Fast and accurate imputation of summary statistics enhances evidence of functional enrichment. *Bioinformatics* 2014;30:2906–14.
18. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 2010;26:2190–1.
19. Morris AP. Transethnic meta-analysis of genomewide association studies. *Genet Epidemiol* 2011;35:809–22.
20. Westra HJ, Peters MJ, Esko T, *et al.* Systematic identification of trans eQTLs as putative drivers of known disease associations. *Nat Genet* 2013;45:1238–43.
21. GTEx Consortium. Human Genomics. The Genotype–Tissue Expression [GTEx] pilot analysis: multitissue gene regulation in humans. *Science* 2015;348: 648–60.
22. Lappalainen T, Sammeth M, Friedländer MR, *et al.*; Geuvadis Consortium. Transcriptome and genome sequencing uncovers functional variation in humans. *Nature* 2013;501:506–11.
23. Jostins L, Ripke S, Weersma RK, *et al.*; International IBD Genetics Consortium [IIBDGC]. Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease. *Nature* 2012;491:119–24.
24. Green LM, Wagner KJ, Campbell HA, Addison K, Roberts SG. Dynamic interaction between WT1 and BASP1 in transcriptional regulation during differentiation. *Nucleic Acids Res* 2009;37:431–40.
25. Horsthemke M, Bachg AC, Groll K, *et al.* Multiple roles of filopodial dynamics in particle capture and phagocytosis and phenotypes of Cdc42 and Myo10 deletion. *J Biol Chem* 2017;292:7258–73.
26. Zimprich A, Biskup S, Leitner P, *et al.* Mutations in LRRK2 cause autosomal-dominant Parkinsonism with pleomorphic pathology. *Neuron* 2004;44:601–7.
27. Morimoto K, Baba Y, Shinohara H, *et al.* LRRK1 is critical in the regulation of B-cell responses and CARMA1-dependent NF- κ B activation. *Sci Rep* 2016;6:25738.
28. Kitabatake M, Soma M, Zhang T, *et al.* JNK regulatory molecule GSPR induces IgG autoantibody-producing plasmablasts from peritoneal B1a cells. *J Immunol* 2015;194:1480–8.
29. Jones EL, Demaria MC, Wright MD. Tetraspanins in cellular immunity. *Biochem Soc Trans* 2011;39:506–11.
30. Gartlan KH, Belz GT, Tarrant JM, *et al.* A complementary role for the tetraspanins CD37 and Tssc6 in cellular immunity. *J Immunol* 2010;185:3158–66.
31. Berndt SI, Skibola CF, Joseph V, *et al.* Genome-wide association study identifies multiple risk loci for chronic lymphocytic leukemia. *Nat Genet* 2013;45:868–76.
32. Dimas AS, Deutsch S, Stranger BE, *et al.* Common regulatory variation impacts gene expression in a cell type-dependent manner. *Science* 2009;325:1246–50.
33. Fehrmann RS, Jansen RC, Veldink JH, *et al.* Trans-eQTLs reveal that independent genetic variants associated with a complex phenotype converge on intermediate genes, with a major role for the HLA. *PLoS Genet* 2011;7:e1002197.
34. McGovern DP, Gardet A, Törkviist L, *et al.*; NIDDK IBD Genetics Consortium. Genome-wide association identifies multiple ulcerative colitis susceptibility loci. *Nat Genet* 2010;42:332–7.
35. Chang CH, Flavell RA. Class II transactivator regulates the expression of multiple genes involved in antigen presentation. *J Exp Med* 1995;181:765–7.
36. Steimle V, Otten LA, Zufferey M, Mach B. Complementation cloning of an MHC class II transactivator mutated in hereditary MHC class II deficiency [or bare lymphocyte syndrome]. *Cell* 1993;75:135–46.
37. Rioux JD, Xavier RJ, Taylor KD, *et al.* Genome-wide association study identifies new susceptibility loci for Crohn disease and implicates autophagy in disease pathogenesis. *Nat Genet* 2007;39:596–604.
38. Roberts RL, Hollis-Moffatt JE, Geary RB, Kennedy MA, Barclay ML, Merriman TR. Confirmation of association of IRGM and NCF4 with ileal Crohn's disease in a population-based cohort. *Genes Immun* 2008;9:561–5.
39. Muise AM, Xu W, Guo CH, *et al.*; NEOPICS. NADPH oxidase complex and IBD candidate gene studies: identification of a rare variant in NCF2 that results in reduced binding to RAC2. *Gut* 2012;61:1028–35.
40. Somasundaram R, Deuring JJ, van der Woude CJ, Peppelenbosch MP, Fuhler GM. Linking risk conferring mutations in NCF4 to functional consequences in Crohn's disease. *Gut* 2012;61:1097; author reply 1097–8.