

# UC Davis

## UC Davis Previously Published Works

### Title

Implications of salinity normalization of seawater total alkalinity in coral reef metabolism studies

### Permalink

<https://escholarship.org/uc/item/7r19z7n7>

### Journal

PLOS ONE, 16(12)

### ISSN

1932-6203

### Authors

Courtney, Travis A  
Cyronak, Tyler  
Griffin, Alyssa J  
[et al.](#)

### Publication Date

2021

### DOI

10.1371/journal.pone.0261210

Peer reviewed

## RESEARCH ARTICLE

# Implications of salinity normalization of seawater total alkalinity in coral reef metabolism studies

Travis A. Courtney<sup>1,2\*</sup>, Tyler Cyronak<sup>3</sup>, Alyssa J. Griffin<sup>4</sup>, Andreas J. Andersson<sup>1</sup>

**1** Scripps Institution of Oceanography, University of California San Diego, La Jolla, California, United States of America, **2** Department of Marine Sciences, University of Puerto Rico Mayagüez, Mayagüez, Puerto Rico, United States of America, **3** Department of Marine and Environmental Sciences, Nova Southeastern University, Fort Lauderdale, Florida, United States of America, **4** Bodega Marine Laboratory, University of California Davis, Davis, CA, United States of America

\* [traviscourtney@gmail.com](mailto:traviscourtney@gmail.com)**OPEN ACCESS**

**Citation:** Courtney TA, Cyronak T, Griffin AJ, Andersson AJ (2021) Implications of salinity normalization of seawater total alkalinity in coral reef metabolism studies. PLoS ONE 16(12): e0261210. <https://doi.org/10.1371/journal.pone.0261210>

**Editor:** Goulven G. Laruelle, Universite Libre de Bruxelles, BELGIUM

**Received:** April 29, 2021

**Accepted:** November 28, 2021

**Published:** December 29, 2021

**Copyright:** © 2021 Courtney et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** Computer code and all data used to generate the analyses and figures in this manuscript are publicly available via GitHub at <https://github.com/traviscourtney/SalinityNormalization> and the final version will be archived with a permanent DOI via Zenodo following peer review.

**Funding:** This work was supported by funding from the National Oceanographic and Atmospheric Administration Ocean Acidification Program Award# NA15OAR4320071 (AJA, TAC). The

## Abstract

Salinity normalization of total alkalinity (TA) and dissolved inorganic carbon (DIC) data is commonly used to account for conservative mixing processes when inferring net metabolic modification of seawater by coral reefs. Salinity (S), TA, and DIC can be accurately and precisely measured, but salinity normalization of TA (nTA) and DIC (nDIC) can generate considerable and unrecognized uncertainties in coral reef metabolic rate estimates. While salinity normalization errors apply to nTA, nDIC, and other ions of interest in coral reefs, here, we focus on nTA due to its application as a proxy for net coral reef calcification and the importance for reefs to maintain calcium carbonate production under environmental change. We used global datasets of coral reef TA, S, and modeled groundwater discharge to assess the effect of different volumetric ratios of multiple freshwater TA inputs (i.e., groundwater, river, surface runoff, and precipitation) on nTA. Coral reef freshwater endmember TA ranged from -2 up to 3032  $\mu\text{mol/kg}$  in hypothetical reef locations with freshwater inputs dominated by riverine, surface runoff, or precipitation mixing with groundwater. The upper bound of freshwater TA in these scenarios can result in an uncertainty in reef TA of up to 90  $\mu\text{mol/kg}$  per unit S normalization if the freshwater endmember is erroneously assumed to have 0  $\mu\text{mol/kg}$  alkalinity. The uncertainty associated with S normalization can, under some circumstances, even shift the interpretation of whether reefs are net calcifying to net dissolving, or vice versa. Moreover, the choice of reference salinity for normalization implicitly makes assumptions about whether biogeochemical processes occur before or after mixing between different water masses, which can add uncertainties of  $\pm 1.4\%$  nTA per unit S normalization. Additional considerations in identifying potential freshwater sources of TA and their relative volumetric impact on seawater are required to reduce uncertainties associated with S normalization of coral reef carbonate chemistry data in some environments. However, at a minimum, researchers should minimize the range of salinities over which the normalization is applied, precisely measure salinity, and normalize TA values to a carefully selected reference salinity that takes local factors into account.

funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

## Introduction

Coral reef metabolic measurements are important tools used to quantify a reef's carbon cycle, health and function, and responses to ongoing environmental change. Net reef metabolism (e.g. primary production, respiration, calcification, and  $\text{CaCO}_3$  dissolution) is typically quantified through changes in seawater dissolved inorganic carbon (DIC) and total alkalinity (TA) concentrations to determine net ecosystem production ( $\text{NEP} = \text{primary production} - \text{total respiration}$ ) and net ecosystem calcification ( $\text{NEC} = \text{calcification} - \text{CaCO}_3 \text{ dissolution}$ ) [1, 2]. Notably, positive coral reef net calcification (i.e., +NEC) is critical for maintaining the calcium carbonate structures and the associated ecosystem services that coral reefs provide [3]. Herein, we evaluate the salinity normalization of TA data and its impact on using alkalinity as a proxy for evaluating coral reef net calcification.


The total alkalinity anomaly technique is pragmatic for determining NEC in coral reefs because TA changes by a factor of two for every unit of  $\text{CaCO}_3$  formed or dissolved [4–6] with negligible to little influence from other processes (e.g., uptake and release of nutrients) in most coral reef environments [7]. Furthermore, TA is conservative with respect to mixing and changes in seawater temperature and pressure [8]. As a result of reef calcification, TA can vary by upwards of hundreds of  $\mu\text{mol/kg}$  across coral reef environments [2]. Consequently, the alkalinity anomaly method, which calculates the difference between initial (e.g., offshore, upstream, or proximal) and final (e.g., coral reef, downstream, or distal) TA can be used to determine whether a reef is net calcifying and combined with measurements of depth and water residence time or transit times to quantify rates of coral reef net ecosystem calcification [5, 9]. However, to account for any changes in TA due to freshwater dilution, evaporation, and mixing between water masses, TA data are commonly normalized with respect to changes in salinity, resulting in a salinity normalized alkalinity anomaly (i.e.,  $\Delta n\text{TA}$ ). Throughout this manuscript, we will use the notation  $\Delta n\text{TA} = n\text{TA}_{\text{offshore}} - n\text{TA}_{\text{reef}}$  to maintain consistency, but it is important to note that  $\Delta n\text{TA}$  could represent any  $n\text{TA}_{\text{initial}} - n\text{TA}_{\text{final}}$ . Salinity normalization of TA is performed using a reference salinity ( $S_{\text{ref}}$ ), freshwater TA endmember ( $\text{TA}_{S=0}$ ), and salinity ( $S$ ) data per the following equation [10]:

$$n\text{TA} = \frac{\text{TA} - \text{TA}_{S=0}}{S} \times S_{\text{ref}} + \text{TA}_{S=0} \quad (1)$$

Salinity normalization of seawater TA in coral reef metabolism studies has traditionally used simple dilution concentration (SDC) mechanisms (i.e., freshwater dilution and evaporation) between  $\text{TA}_{\text{offshore}}$  and  $\text{TA}_{S=0} = 0$  (Eq 1) [10]. However, assumptions regarding SDC can be invalidated by processes that modify freshwater alkalinity such as the weathering of carbonate and silicate minerals [11, 12], anaerobic redox processes, and the anthropogenic deposition of sulfuric and nitric oxides as acid rain [13, 14]. Coral reefs are complex coastal ecosystems with many potential sources of freshwater TA (i.e.,  $\text{TA}_{S=0}$ ) including groundwater (95–13,000  $\mu\text{mol/L}$  [15]), tropical rivers (224–2,156  $\mu\text{mol/L}$  [12]), and precipitation (-2.7–18  $\mu\text{mol/L}$  [14]) such that:

$$\text{TA}_{S=0} = \chi_{\text{gw}} \text{TA}_{\text{gw}} + \chi_{\text{rw}} \text{TA}_{\text{rw}} + \chi_{\text{p}} \text{TA}_{\text{p}} \quad (2)$$

where  $\chi$  represents the relative volumetric freshwater proportion of groundwater ( $\chi_{\text{gw}}$ ), riverine water ( $\chi_{\text{rw}}$ ), or precipitation ( $\chi_{\text{p}}$ ), and their TA concentrations, respectively. If freshwater sources with significant  $\text{TA}_{S=0}$  are volumetrically important, but their concentrations and relative proportions are unknown (Eq 2), the system is underdetermined and could result in errors in both the magnitude and direction of coral reef  $\Delta n\text{TA}$  (Fig 1A–1C). Moreover, in



**Fig 1. Conceptual diagram of salinity normalization of coral reef total alkalinity data.** (a,b,c) Conceptual diagram demonstrates how conservative mixing of a representative surface TA-S of open ocean (i.e., designated by O on plot;  $\text{TA}_{\text{offshore}} = 2300 \mu\text{mol/kg}$ ,  $S_{\text{offshore}} = 35 \text{ g/kg}$ ) with a representative freshwater TA endmember of  $\text{TA}_{S=0} = 1500$  (e.g., see simulations below) or  $\text{TA}_{S=0} = 0$  can create considerable uncertainties in the (b) sign and (c) magnitude of  $\Delta n\text{TA}$  calculated with respect to the coral reef seawater TA (i.e.,  $\text{TA}_{\text{reef}}$  designated by R on plot). The yellow regions indicate the uncertainty of  $\Delta n\text{TA}$  if the actual  $\text{TA}_{S=0}$  is unknown but within two values in this region. The  $\Delta n\text{TA}$  box (b, c) shows the magnitude of uncertainty for  $\Delta n\text{TA}$  between the ocean (O) and reef (R) owing to an unknown  $\text{TA}_{S=0}$  where  $\Delta n\text{TA}$  is not equal ( $\neq$ ) in direction in (b) and  $\Delta n\text{TA}$  is much greater ( $\gg$ ) when assuming  $\text{TA}_{S=0} = 1500 \mu\text{mol/kg}$  compared to  $\text{TA}_{S=0} = 0 \mu\text{mol/kg}$  in (c). Conceptual diagram demonstrates how assuming mixing occurs 1<sup>st</sup> ( $S_{\text{ref}} = S_{\text{reef}}$ ) or calcification occurs 1<sup>st</sup> ( $S_{\text{ref}} = S_{\text{offshore}}$ ) affects the magnitude of  $\Delta n\text{TA}$  inferred from salinity normalized changes in coral reef TA.  $\Delta n\text{TA}$  box shows the magnitude of uncertainty for  $\Delta n\text{TA}$  owing to biogeochemical processes at point R relative to point O such that  $\Delta n\text{TA} = R' - R$  is less than ( $<$ )  $\Delta n\text{TA} = O - O'$ . The conceptual framework is expanded from Fig 9 in [16] to show how unknown  $\text{TA}_{S=0}$  (a,b,c) and unknown order of processes (d,e) can affect interpretation of the magnitude and direction of coral reef NEC determined from changes in seawater total alkalinity.

<https://doi.org/10.1371/journal.pone.0261210.g001>

coastal ecosystems such as coral reefs, it is challenging to accurately estimate freshwater alkalinity endmembers ( $\text{TA}_{S=0}$ ), and consequently nTA, from TA-S relationships because evaporation and net calcification further modify TA-S mixing lines [16]. The presence of multiple potential freshwater TA sources and variable mixing ratios co-occurring with biogeochemical modifications of TA raise concerns that unknown  $\text{TA}_{S=0}$  could be a source of significant uncertainty in coral reef  $\Delta n\text{TA}$  estimates and, consequently, estimates of net coral reef calcification rates.

In addition to the effects of unknown  $\text{TA}_{S=0}$  on  $\Delta n\text{TA}$ , the application of salinity normalization and the choice of a reference salinity implicitly introduce assumptions about whether conservative mixing occurs before biogeochemical modification (i.e., net ecosystem calcification) or vice versa [16]. For example, Fig 1D and 1E illustrate mixing and biogeochemical modification between offshore TA-S (point O) and reef TA-S (point R). By salinity normalizing coral reef TA (point R) to open ocean endmember salinity (point O) before calculating  $\Delta n\text{TA}$ , R is normalized to O' and  $\Delta n\text{TA}$  is calculated as the difference in TA between O and O' (Fig 1D and 1E). This scenario ( $S_{\text{ref}} = S_{\text{offshore}}$ ) implicitly assumes that calcification occurs prior to mixing with the freshwater endmember. Conversely, by salinity normalizing open ocean TA endmember (point O) to the coral reef salinity (point R) before calculating  $\Delta n\text{TA}$ , O

is normalized to  $R'$  and  $\Delta nTA$  is calculated as the difference in TA between  $R'$  and  $R$  (Fig 1E). This scenario ( $S_{ref} = S_{reef}$ ) implicitly assumes that mixing with the freshwater endmember occurs prior to calcification. The  $\Delta nTA$  inferred from  $O$  and  $O'$  is slightly greater than the  $\Delta nTA$  inferred from  $R'$  and  $R$  (Fig 1E). Therefore, the choice of  $S_{ref}$  in the salinity normalization of coral reef TA data via Eq 1 implicitly assumes whether calcification occurs before mixing (i.e.,  $S_{ref} = S_{offshore}$ ), mixing occurs before calcification (i.e.,  $S_{ref} = S_{reef}$ ), or some order of mixing between these two extremes (i.e.,  $S_{reef} < S_{ref} < S_{offshore}$ ) and generates an additional source of uncertainty in coral reef  $\Delta nTA$ .

In a previous study, we suggested that uncertainties in seawater depth and residence time can potentially drive large uncertainties in estimates of coral reef NEC rates while assuming that the accurate determination of TA ( $\pm 2 \mu\text{mol/kg}$ ) was unlikely to be a significant source of error [17]. Here, we quantify the potential uncertainties in  $\Delta nTA$  associated with the salinity normalization of coral reef total alkalinity to unknown freshwater endmembers ( $TA_{S=0}$ ) and reference salinities ( $S_{ref}$ ) using global seawater TA and  $S$ , fresh submarine groundwater discharge, a range of different  $TA_{S=0}$  datasets, and different mixing proportions of freshwater endmembers. We provide a series of suggestions to reduce the uncertainties associated with salinity normalization of coral reef TA data with implications for other carbonate chemistry parameters and ions of interest to coral reef metabolism studies (e.g., DIC,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ , etc.).


## Methods

### Estimating potential coral reef $TA_{S=0}$

In the absence of detailed freshwater budgets and relative volumetric proportions of different freshwater TA endmembers in coral reefs, we simulated coral reef  $TA_{S=0}$  using literature derived data in a Monte Carlo approach ( $n = 10,000$ ) assuming three distinct scenarios: (i) riverine + groundwater freshwater inputs (Fig 2A); (ii) terrestrial surface runoff + groundwater freshwater inputs (Fig 2B); and (iii) precipitation + groundwater freshwater inputs (Fig 2C). For each scenario, a random selection was made from the ratio of fresh submarine groundwater discharge to surface runoff for coral reefs ( $n = 4,241$ ; Fig 2D) calculated by [18]. We then combined this selected ratio of groundwater inputs value with randomly generated TA concentrations from within the range of the typical coral reef groundwaters (95–13,000  $\mu\text{mol/L}$  [15]), tropical rivers (224–2,156  $\mu\text{mol/L}$  [12]), and precipitation (-2.7–18  $\mu\text{mol/L}$  [14]). This analysis allowed us to simulate the potential TA concentration of mixed freshwater endmembers for each of the three scenarios described above (Fig 2A). The simulated coral reef  $TA_{S=0}$  were positively skewed so the 50<sup>th</sup>, 2.5<sup>th</sup>, and 97.5<sup>th</sup> percentiles were determined for each of three scenarios to approximate the median and range of potential coral reef  $TA_{S=0}$ .

### Global coral reef total alkalinity data

We used a global assessment of coral reef seawater carbonate chemistry and salinity data [2] to quantify coral reef TA-S relationships, salinity ranges, and the effect of salinity normalization on  $TA_{reef}$ . To assess TA-S relationships, type II ordinary least squares linear regression of TA vs.  $S$  were constructed for each of the 27 coral reefs using the function *lmodel2* [19] in *R* [20]. To evaluate the range and distribution of salinity changes, we subtracted every salinity value from the maximum salinity value for each of the 27 coral reefs.  $nTA_{reef}$  was calculated via Eq (1) using the mean salinity for each site (i.e.,  $S_{ref} = S_{mean}$ ) and assuming  $TA_{S=0} = 0$ . We then calculated the difference between  $TA_{reef}$  and  $nTA_{reef}$  for each of the 27 coral reefs to estimate the magnitude to which salinity normalization changes  $TA_{reef}$  data.



**Fig 2. Simulations of freshwater total alkalinity endmembers.** (a,b,c) Conceptual diagram shows the relative contributions of freshwater TA endmembers (i.e.,  $TA_{S=0}$ ) for simulations of (a) riverine dominated, (b) surface runoff dominated, and (c) precipitation dominated reef systems where  $r_{gw:sw}$  = ratio of groundwater to surface water,  $r_{p:rw}$  = ratio of precipitation to riverine water,  $TA_{gw}$  = groundwater TA,  $TA_{rw}$  = riverine TA,  $TA_p$  = precipitation TA, and  $TA_{sw}$  = surface water TA. (d) Map shows the coral reef locations with fresh submarine groundwater discharge in [18] used with the equations in (a, b, c) to calculate (e) coral reef  $TA_{S=0}$  box plots representing 2.5<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, and 97.5<sup>th</sup> percentiles of estimated coral reef  $TA_{S=0}$  from the Monte-Carlo simulations for the precipitation, riverine, and surface runoff dominated reef systems.

<https://doi.org/10.1371/journal.pone.0261210.g002>

### Effects of unknown $TA_{S=0}$ on $\Delta nTA$

We quantified  $nTA_{reef}$  via Eq (1) using the maximum salinity for each site as the reference salinity (i.e.,  $S_{ref} = S_{max}$ ) and  $TA_{S=0}$  values of 0, 1000, 2000, and 3000  $\mu\text{mol/kg}$ , which span the full range of potential simulated freshwater TA endmembers in coral reef ecosystems from this study (Fig 2E). In the absence of reported  $S_{offshore}$  for every coral reef location, we assumed that  $S_{max}$  represents  $S_{offshore}$  to calculate  $\Delta nTA$  as the mean of  $TA_{offshore} - nTA_{reef}$  and  $\Delta S$  as the mean of  $S_{max} - S_{ref}$  for all samples from each reef location in [2]. Furthermore, because the true  $TA_{S=0}$  for each reef location were unknown, we quantified the uncertainties in  $\Delta nTA$  owing to mixing within the range of freshwater TA endmembers explored in this study. To accomplish this, we subtracted the  $\Delta nTA$  values calculated assuming a zero freshwater TA endmember from the  $\Delta nTA$  values calculated assuming a positive freshwater TA endmember (i.e.,  $\Delta nTA_{TA>0@S=0} - \Delta nTA_{TA=0@S=0}$ ). Linear regression of these  $\Delta nTA$  uncertainty vs.  $\Delta S$  relationships for each  $TA_{S=0}$  represent the potential range (i.e., uncertainty) in the actual  $\Delta nTA$  values for a given change in S assuming the true  $TA_{S=0}$  for each reef site is between the 0 and 3000  $\mu\text{mol/kg}$  values explored in this analysis. To further explore how the precision of S can affect  $nTA$ , the minimum difference in reported salinity data for each reef was used to estimate the precision of the salinity measurement.



## Effects of unknown order of processes on $\Delta nTA$

To quantify how the implicit order of mixing and biogeochemical processes impact coral reef  $\Delta nTA$  data, we quantified  $\Delta nTA$  as either  $\Delta nTA_{\text{offshore}} = TA_{\text{offshore}} - nTA_{\text{reef}}$  (i.e.,  $S_{\text{ref}} = S_{\text{offshore}}$  assuming calcification occurs before mixing; O to O' to R sensu Fig 1E) or  $\Delta nTA_{\text{reef}} = nTA_{\text{offshore}} - TA_{\text{reef}}$  (i.e.,  $S_{\text{ref}} = \text{mean}(S_{\text{reef}})$  assuming mixing occurs before calcification; O to R' to R sensu Fig 1E) for each reef location in [2]. We then quantified the mean percent difference between  $\Delta nTA_{\text{offshore}}$  and  $\Delta nTA_{\text{reef}}$  for each site to evaluate the type II ordinary least squares linear regression between %  $\Delta nTA$  uncertainty and  $\Delta S$  for each reef site using the function *lmodel2* [19] in R [20] to see how uncertainty in  $\Delta nTA$  scales with  $\Delta S$  owing to the unknown order of mixing and biogeochemical processes.


## Results and discussion

### Coral reef $TA_{S=0}$

The global ratio of coral reef fresh submarine groundwater discharge to surface water runoff database [18] provided an opportunity to simulate  $TA_{S=0}$  for coral reef locations where precipitation, riverine, and freshwater runoff occur in different volumetric proportions (Fig 2). Coral reef  $TA_{S=0}$  estimates were positively skewed with median (2.5<sup>th</sup> to 97.5<sup>th</sup> percentile) values of 15  $\mu\text{mol/L}$  (-2 to 1997  $\mu\text{mol/L}$ ) for groundwater mixing with precipitation, 1306  $\mu\text{mol/L}$  (293 to 2979  $\mu\text{mol/L}$ ) for groundwater mixing with river water, and 598  $\mu\text{mol/L}$  (43 to 2507  $\mu\text{mol/L}$ ) for groundwater mixing with surface runoff (Fig 2E). The range of simulations in this study suggest that these uncertainties vary between coral reefs owing to proximity to terrestrially derived freshwater inputs and over time and space within coral reefs owing to temporally and geographically variable freshwater inputs (Fig 2). In particular, assessing the proximity of the reef system to land masses and/or notable surface freshwater inputs could inform which of the three simulated coral reef  $TA_{S=0}$  ( $\pm$  uncertainty) scenarios explored in this study may best represent uncertainties associated with unknown  $TA_{S=0}$  in future studies where  $TA_{S=0}$  and their respective volumetric proportions are unknown. However, we nonetheless caution that these simulations are hypothetical (Fig 2D) and that  $TA_{S=0}$  may exceed these estimates in select coral reefs with elevated volumetric proportions of groundwater or within individual coral reefs where the assumption of complete mixing between multiple  $TA_{S=0}$  is invalid. Submarine groundwater discharge can be patchy and heterogeneous across small spatial scales, meaning biogeochemical processes and hydrodynamics should be assessed locally [21, 22]. Conversely, the influx of low  $TA_{S=0}$  owing to precipitation falling directly on the reef and/or adjacent lagoon waters could drive mean freshwater  $TA_{S=0}$  lower than the simulated values in this study. Moreover, land-based precipitation is likely to reach the reef in the form of runoff, riverine inputs, and/or groundwater at some time lag following the initial precipitation event, suggesting that the timing of precipitation events is an important consideration in estimating the relative volumetric contributions of multiple  $TA_{S=0}$  sources.

### Global coral reef TA-S relationships

We used the global dataset of TA and S data to explicitly test TA-S relationships in coral reef metabolism studies conducted over a broad range of spatial and temporal scales [2]. The majority of coral reefs exhibited detectable linear relationships between TA and S (i.e., 19 of 25 reefs have 95% confidence intervals that do not overlap the dashed 0 line) (Fig 3A and S1 Table), which indicates the need for salinity normalization of  $TA_{\text{reef}}$  data to account for mixing of water with different properties and the influence of freshwater inputs or evaporation on TA [10]. Notably, data from two reefs exhibited no reported changes in S so the TA-S slopes were



**Fig 3. Global coral reef total alkalinity and salinity relationships.** Relationships between TA, S, and nTA are explored across global coral reef environments from [2]. (a) The slope ( $\pm$  95% confidence interval) of linear regressions between TA and S for each reef. Slopes that do not overlap the dashed zero line are detectably different from zero. (b) The range of salinities defined as the  $S_{max}-S$  for each reef datapoint with all data outlined by the probability density for each site to visualize the distribution. (c) The difference between TA and nTA calculated using  $S_{mean}$  as the  $S_{ref}$  and  $TA_{S=0} = 0$  via Eq (1) for each reef with all data outlined by the probability density for each site to visualize the distribution. Sources of coral reef data are 01 = Makapu’u, 03 = Palmyra, 04 = Red Sea 1997, 05 = Red Sea 1998, 06 = Red Sea 2013, 07 = Ofu, 08 = Heron Island, 09 = Palau 1, 11 = Lizard Island, 12 = Ishigaki, 13 = One Tree 1, 14 = Kaiona, 15 = Lady Elliot, 16 = One Tree 2, 17 = Davies, 18 = Palau 2, 19 = Cook Islands, 20 = Cheeca Rocks, 21 = St. John, 22 = West Panama, 23 = Florida Keys, 24 = Bermuda, 25 = Majuro, 26 = Maldives, and 27 = Puerto Rico sensu [2]. See references and discussion in [2] for further details on each coral reef location.

<https://doi.org/10.1371/journal.pone.0261210.g003>


undefined for these reef sites and removed from this analysis (i.e., 02 = Mo’orea and 10 = Yonge Reef). Mean ranges in  $S_{reef}$  (i.e.,  $S_{max}-S_{reef}$ ) for each of the reefs varied from 0.05 to 2.1 (mean  $\pm$  SD for all reefs =  $0.5\pm 0.6$ ) (Fig 3B). Mean ( $\pm$ SD) differences between non-normalized and normalized TA (i.e., assuming  $TA_{S=0} = 0$  and  $S_{ref} = S_{max} = S_{offshore}$  via Eq 1) were  $-0.2\pm 30.4$   $\mu\text{mol/kg}$  (range =  $-311.9$  to  $+138.2$   $\mu\text{mol/kg}$ ) (Fig 3C). This mean difference of  $-0.2$   $\mu\text{mol/kg}$  supports previous findings that salinity normalization had a negligible impact on mean  $\Delta nTA$  across the reef sites presented here [2], especially for reef sites with minimal S ranges (Fig 3B and 3C). However, the range of  $-311.9$  to  $+138.2$   $\mu\text{mol/kg}$  values is the same magnitude as the estimated changes in TA strictly owing to coral reef NEC [17] suggesting



that salinity normalization of coral reef TA can nonetheless have a significant impact on individual  $\Delta nTA$  estimates with greater deviations in salinity from  $S_{ref}$ .

### Estimated effects of unknown $TA_{S=0}$ on $nTA_{reef}$

The alkalinity anomalies for each reef location normalized to different  $TA_{S=0}$  show the potential for non-negligible uncertainties in  $\Delta nTA$  if a positive  $TA_{S=0}$  is not accounted for (Fig 4A). Most importantly, the magnitude of the difference between  $\Delta nTA$  calculated assuming a positive freshwater TA endmember and  $\Delta nTA$  calculated assuming a zero freshwater TA endmember (i.e.,  $\Delta nTA_{TA>0@S=0} - \Delta nTA_{TA=0@S=0}$ ) scaled with the magnitude of the difference between  $S_{max}$  and  $S_{ref}$  (Eq 1 and Fig 4A). Maximum uncertainties associated with freshwater ( $TA_{S=0}$ ) equal to 1000, 2000, and 3000  $\mu\text{mol/kg}$  compared to  $TA_{S=0} = 0 \mu\text{mol/kg}$  were 30, 60, and 90  $\mu\text{mol/kg}$  per unit difference in salinity, respectively (Fig 4A). Assuming that  $TA_{S=0} = 0$  in the salinity normalization of coral reef TA via Eq 1 can therefore yield uncertainties in  $\Delta nTA$  that increase linearly with increasing  $TA_{S=0}$ . Similarly, if  $TA_{S=0} > 0$  is assumed during salinity normalization, then TA uncertainties increase with increasing offset between  $TA_{S=0}$  and the assumed  $TA_{S=0}$ . Establishing potential upper and lower bounds on  $TA_{S=0}$  for different freshwater endmembers on coral reefs and their relative volumetric contributions to the reef therefore provide an effective means to reduce the potential range of uncertainties associated with the salinity normalization of  $TA_{reef}$  (Fig 2E). The canonical assumption that  $TA_{S=0} = 0$  for salinity normalization [10] is close to the median  $TA_{S=0} = 15 \mu\text{mol/L}$  for the precipitation mixing with groundwater scenario in this study, which suggests that  $TA_{S=0} = 0$  may remain a reasonable approximation at least for precipitation dominated reef systems and habitats. However, it is important to note that this is from simulated freshwater endmembers in coral reef ecosystems and freshwater TA concentrations can vary considerably in reef ecosystems. Calculating  $nTA$  using the full range of  $TA_{S=0}$  for precipitation dominated systems (i.e.,



**Fig 4. Salinity normalized alkalinity anomaly uncertainties scale with salinity changes.** (a) The mean alkalinity anomaly for each reef evaluated for  $TA_{S=0}$  ranging from 0, 1000, 2000, 3000  $\mu\text{mol/kg}$  using  $S_{ref} = S_{max}$  in Eq (1) minus the mean alkalinity anomaly for  $TA_{S=0} = 0$  are plotted against the mean  $S_{max} - S_{ref}$  for each reef location in [2] to quantify how uncertainty in  $\Delta nTA$  scales with unknown  $TA_{S=0}$ . Values to the right of each line plotted on the graph are the uncertainty in  $\Delta nTA$  per unit salinity for the respective  $TA_{S=0}$  line. (b) The mean percent difference between  $\Delta nTA_{offshore}$  and  $\Delta nTA_{reef}$  owing to the choice of  $S_{ref}$  of  $S_{max} = S_{offshore}$  or  $S_{reef}$  via Eq (1) are plotted against the mean  $S_{max} - S_{ref}$  for each reef location in [2] to evaluate how uncertainty associated with the order of mixing processes scale with salinity changes across the reef. The vertical dashed gray line indicates uncertainty per unit salinity change in each panel.

<https://doi.org/10.1371/journal.pone.0261210.g004>

-2 to 2038  $\mu\text{mol/L}$ ) may nonetheless provide a more honest assessment of  $n\text{TA} \pm$  uncertainties for cases where  $\text{TA}_{S=0}$  and the relative volumetric freshwater fluxes are unknown.

We recommend viewing coral reefs from an ecosystem-scale perspective to consider the volumetric input of freshwater from different sources (i.e., groundwater, river input, and/or runoff account for a substantial fraction of freshwater input) when using  $\Delta\text{TA}$  as a proxy for coral reef calcification and, if warranted, locate and directly measure  $\text{TA}_{S=0}$  of these freshwater endmembers. However, the multitude of endmembers and biogeochemical processes increasingly complicate estimates of the mixing derived component of coral reef  $\text{TA}_{\text{reef}}:S$  variability. While the estimated freshwater TA endmembers for precipitation, riverine, and runoff dominated systems from this study (Fig 2E) may provide alternative  $\text{TA}_{S=0}$  estimates  $\pm$  uncertainties wherever the precise quantification of  $\text{TA}_{S=0}$  values and relative volumetric inputs is not feasible, any assumptions regarding the salinity normalization of TA data should nonetheless be carefully considered (e.g., see below list of considerations). Moreover, utilizing other tracers to normalize seawater TA data such as silicate, Ra, Rn, Li, Mg, Sr, Ca, Ba,  $\delta^{18}\text{O}$ , and other tracers for submarine groundwater discharge [22–25] warrant further consideration as additional means to parse out the relative contributions of mixing of multiple freshwater endmembers and biogeochemical fluxes on coral reef seawater carbonate chemistry data.

### Effects of unknown order of processes on $\Delta n\text{TA}$

The  $\Delta n\text{TA}$  uncertainties owing to the implicitly assumed order of mixing and biogeochemical processes in the salinity normalization of TA were much smaller than the uncertainties owing to the unknown  $\text{TA}_{S=0}$  (Fig 4). For example, there was a 2.8% range of uncertainty in  $\Delta n\text{TA}$  per unit salinity owing to normalizing ocean and coral reef TA data to  $S_{\text{offshore}}$  compared to  $S_{\text{reef}}$  (Fig 4B). Assuming an actual 100  $\mu\text{mol/kg}$  alkalinity anomaly solely owing to net calcification, this 2.8% uncertainty per unit salinity translates to a 2.8  $\mu\text{mol/kg}$  uncertainty range, which is similar to the typical precision of  $\pm 2$   $\mu\text{mol/kg}$  for coral reef TA measurements [17]. The choice of normalizing to  $S_{\text{offshore}}$  or  $S_{\text{reef}}$  represent extreme scenarios where either all calcification is assumed to occur before all mixing ( $S_{\text{ref}} = S_{\text{offshore}}$ ) or all mixing is assumed to occur before all calcification ( $S_{\text{ref}} = S_{\text{reef}}$ ) (Fig 1D and 1E). Unless there is information to rigorously evaluate which order of processes is occurring, calculating both  $\Delta n\text{TA}_{\text{offshore}}$  using  $S_{\text{ref}} = S_{\text{offshore}}$  (i.e., calcification then mixing) and  $\Delta n\text{TA}_{\text{reef}}$  using  $S_{\text{ref}} = S_{\text{reef}}$  (i.e., mixing then calcification) quantifies the maximum uncertainty associated with the unknown order of mixing and calcification processes as follows:

$$\Delta n\text{TA}_{\text{offshore}} = \text{TA}_{\text{offshore}} - \left[ (\text{TA}_{\text{reef}} - \text{TA}_{S=0}) \frac{S_{\text{offshore}}}{S_{\text{reef}}} + \text{TA}_{S=0} \right] \tag{3}$$

$$\Delta n\text{TA}_{\text{reef}} = \left[ (\text{TA}_{\text{offshore}} - \text{TA}_{S=0}) \frac{S_{\text{reef}}}{S_{\text{offshore}}} + \text{TA}_{S=0} \right] - \text{TA}_{\text{reef}} \tag{4}$$

Importantly, the average of  $\Delta n\text{TA}_{\text{offshore}}$  and  $\Delta n\text{TA}_{\text{reef}}$  results in a  $\Delta n\text{TA}$  that assumes net calcification and mixing occur simultaneously and results in a centered, quantifiable  $\pm 1.4\%$  uncertainty in  $\Delta n\text{TA}$  per unit salinity change.

### Salinity and $\text{TA}_{\text{reef}}$ variability

In studies aimed at assessing net reef calcification, the simplest way to reduce uncertainties in coral reef  $n\text{TA}$  is by selecting sampling sites and time periods with minimal S variability (Figs 3 and 4). Selecting sites with known hydrodynamics could further reduce mixing related

uncertainties. Any deviations between  $S_{\text{offshore}}$  and  $S_{\text{reef}}$  have the potential to generate non-negligible uncertainties in coral reef salinity normalized total alkalinity anomalies (i.e.,  $\Delta n\text{TA} = n\text{TA}_{\text{offshore}} - n\text{TA}_{\text{reef}}$ ). While we have assumed that  $S_{\text{max}} = S_{\text{offshore}}$  to fill in missing data from [2],  $S_{\text{offshore}}$  should ideally be carefully quantified alongside  $\text{TA}_{\text{offshore}}$  to reduce actual  $\Delta n\text{TA}$  uncertainties in subsequent studies. Critically, the choice of sampling location for  $\text{TA}_{\text{offshore}}$  represents an important consideration depending on the aims of the study and characteristics of the coral reef being investigated. For example, choosing an offshore reference that is more proximal to the reef environment can minimize the difference between  $S_{\text{offshore}}$  and  $S_{\text{reef}}$  to reduce uncertainties owing to a large range in salinity. Depending on the specifics of individual reef systems and nearshore oceanographic processes,  $\text{TA}_{\text{offshore}}$  of these more proximal offshore waters may be more strongly influenced by other nearshore processes such as upwelling and coastal freshwater inputs [16]. As a result, the choice of sampling location for offshore reference samples is likely context dependent with the capacity to impact both the salinity normalization of TA and the relative contributions of non-reef processes on  $\Delta n\text{TA}$ .

While TA is often evaluated with a high degree of precision and accuracy (i.e., many labs were within  $\pm 2 \mu\text{mol/kg}$  and the majority of labs were within  $\pm 10 \mu\text{mol/kg}$  in a recent inter-comparison study; [26]), the estimated precision of reported salinity measurements were  $\leq 0.01$  for 18 reefs, 0.03 for 1 reef, 0.1 for 6 reefs, and undefined for the two reef sites which reported no changes in  $S$ . Even small salinity changes of  $\leq 0.1$  across samples can drive significant uncertainties in  $\Delta n\text{TA}$  suggesting that ideally the most precise and accurate  $S$  measurement should be used in the evaluation of coral reef  $\Delta n\text{TA}$ . For example, the rounding of 34.44 to 34.4 and 34.45 to 34.5 by a salinity precision of  $\pm 0.1$  shows how a real 0.01 salinity change can result in an apparent 0.1 salinity change. In this case, normalizing TA to an apparent salinity change of  $\pm 0.1$  could generate uncertainties up to  $9 \mu\text{mol/kg}$  in  $n\text{TA}$  values based on the  $\text{TA}_{S=0} = 3000 \mu\text{mol/kg}$  relative to  $\text{TA}_{S=0} = 0 \mu\text{mol/kg}$  scenario. With current salinity sensing technology, uncertainties in salinity and, consequently,  $n\text{TA}$  could be much smaller. Therefore, any such studies that salinity normalize their TA data may be underreporting the true uncertainties in their measurements by simply stating the precision of their TA measurements and not including the uncertainties owing to their  $S$  measurements and subsequent salinity normalization. Furthermore, if changes in salinity are within the reported uncertainties of the salinity measurement, then there is not sufficient evidence that salinity has detectably changed across the reef environment. In these cases, salinity normalization may introduce unnecessary error into TA changes, and it may be more appropriate to avoid salinity normalization if the  $S_{\text{range}}$  is within the reported precision and/or accuracy of the salinity measurement.

### Implications for coral reef net ecosystem calcification

Here we have shown that the uncertainty of  $\Delta n\text{TA}$  measurements is likely greater than the reported  $\pm 2$  to  $10 \mu\text{mol/kg}$  precision of TA owing to uncertainties associated with the normalization of  $\text{TA}_{\text{reef}}$  data to  $S_{\text{ref}}$  and especially if  $\text{TA}_{S=0}$  is unknown. We have further provided quantitative simulations of coral reef freshwater TA endmembers ( $\text{TA}_{S=0} \pm$  uncertainties) to approximate the potential uncertainties associated with the salinity normalization of coral reef TA data. We conclude that salinity normalization can change both the sign and magnitude of  $\Delta n\text{TA}$  across coral reef environments suggesting that future studies should exercise caution when evaluating and interpreting  $\Delta n\text{TA}$  from coral reef TA and  $S$  data. This is especially true for studies with larger ranges in salinity because the uncertainties associated with  $\Delta n\text{TA}$  positively scale with changes in  $S$ . We offer several considerations for salinity normalizing coral reef carbonate chemistry data in the future to reduce potential  $\Delta n\text{TA}$  uncertainties when using the alkalinity anomaly technique as a proxy for net coral reef calcification in any reef system:

1. Do not automatically assume salinity normalization is necessary, as it could introduce more error into your measurements than expected.
2. Examine data for any detectable correlations between salinity and TA as these relationships indicate the potential influence of salinity dilution or concentration mechanisms on TA. Even if there are no detectable correlations between salinity and TA, evaluate the potential role that simple dilution or concentration mechanisms may have on coral reef TA data.
3. Precisely measure both salinity and TA. Consider the range and precision of your salinity measurements and the potential impact that S normalization has on TA values. If the range in salinity is within the precision of the salinity measurement, consider not salinity normalizing your data.
4. Take a holistic approach to examine possible sources and relative contributions of freshwater and submarine groundwater discharge to the reef system. Consider taking direct freshwater samples to better determine potential endmembers (i.e.,  $TA_{S=0}$ ) that are mixing with the reef water mass to reduce  $\Delta nTA$  uncertainties and consider avoiding reef metabolism studies in regions with large salinity changes that may confound conclusions regarding  $\Delta nTA$ .
5. When the relative volumetric contributions of multiple  $TA_{S=0}$  are unknown, consider calculating  $\Delta nTA$  for a range of  $TA_{S=0}$  values to propagate the potential  $\Delta nTA \pm$  uncertainties owing to unknown  $TA_{S=0}$ .
6. Carefully select  $S_{ref}$  while considering the implicit order of mixing and biogeochemical modification. The mean and range of  $\Delta nTA_{offshore}$  (i.e.,  $S_{ref} = S_{offshore}$ ) and  $\Delta nTA_{reef}$  (i.e.,  $S_{ref} = S_{reef}$ ) can be used to estimate the  $\Delta nTA \pm$  uncertainties owing to the implicitly assumed order of mixing in the salinity normalization of TA.
7. Propagate the potential uncertainties associated with the salinity normalization of coral reef TA data on calculated  $\Delta nTA$  to evaluate the relative importance of the salinity normalization process to the resulting estimate of  $\Delta nTA$  and calculated reef metabolism.

These findings build upon our previous assessment that depth and residence time are the primary sources of NEC uncertainty [17] to include the potentially large uncertainties associated with salinity normalization of TA and DIC data. While this study has focused on coral reef TA for the purpose of brevity, many of these same findings also apply to the salinity normalization of dissolved inorganic carbon (DIC) and other ions of interest (e.g.,  $Ca^{2+}$ ,  $Mg^{2+}$ ) for coral reef metabolism studies, which could be exacerbated depending on their background concentrations in seawater. Reducing uncertainties in the evaluation of coral reef metabolism (e.g. calcification, dissolution, respiration, and photosynthesis) remains critical for observing and quantifying coral reef function and health under unprecedented environmental change.

## Supporting information

**S1 Table. Summary table of coral reef total alkalinity and salinity.** *Source* refers to the following reef locations: 01 = Makapu'u, 03 = Palmyra, 04 = Red Sea 1997, 05 = Red Sea 1998, 06 = Red Sea 2013, 07 = Ofu, 08 = Heron Island, 09 = Palau 1, 11 = Lizard Island, 12 = Ishigaki, 13 = One Tree 1, 14 = Kaiona, 15 = Lady Elliot, 16 = One Tree 2, 17 = Davies, 18 = Palau 2, 19 = Cook Islands, 20 = Cheeca Rocks, 21 = St. John, 22 = West Panama, 23 = Florida Keys, 24 = Bermuda, 25 = Majuro, 26 = Maldives, and 27 = Puerto Rico sensu [2], *method* refers to OLS = ordinary least squares regression, *term* represents the Slope of the relationship, *estimate* represents the value of the slope, *conf.low* represents the lower 95% confidence interval of the

slope, *conf.high* represents the upper 95% confidence interval of the slope, and *p.value* represents the p-value of the slope.  
(CSV)

## Acknowledgments

All coral reef TA-S data [2] and modeled coral reef fresh submarine groundwater discharge data [18] are available from their respective publications and references therein. Computer code and all data used to generate the analyses and figures in this manuscript are publicly available via *GitHub* at <https://github.com/traviscourtney/SalinityNormalization> and *Zenodo* at <https://doi.org/10.5281/zenodo.5752957>.

## Author Contributions

**Conceptualization:** Travis A. Courtney, Tyler Cyronak, Alyssa J. Griffin, Andreas J. Andersson.

**Data curation:** Tyler Cyronak.

**Formal analysis:** Travis A. Courtney.

**Investigation:** Travis A. Courtney, Tyler Cyronak, Alyssa J. Griffin, Andreas J. Andersson.

**Methodology:** Travis A. Courtney, Tyler Cyronak, Alyssa J. Griffin, Andreas J. Andersson.

**Project administration:** Travis A. Courtney, Andreas J. Andersson.

**Resources:** Travis A. Courtney, Andreas J. Andersson.

**Software:** Travis A. Courtney.

**Supervision:** Andreas J. Andersson.

**Visualization:** Travis A. Courtney, Tyler Cyronak, Alyssa J. Griffin.

**Writing – original draft:** Travis A. Courtney.

**Writing – review & editing:** Travis A. Courtney, Tyler Cyronak, Alyssa J. Griffin, Andreas J. Andersson.

## References

1. Smith S. V., Kinsey DW. Calcification and organic carbon metabolism as indicated by carbon dioxide. *Coral reefs: research methods*. 1978. pp. 469–484.
2. Cyronak T, Andersson AJ, Langdon C, Albright R, Bates NR, Caldeira K, et al. Taking the metabolic pulse of the world's coral reefs. *PLOS ONE*. 2018; 13: e0190872. <https://doi.org/10.1371/journal.pone.0190872> PMID: 29315312
3. Brandl SJ, Rasher DB, Côté IM, Casey JM, Darling ES, Lefcheck JS, et al. Coral reef ecosystem functioning: eight core processes and the role of biodiversity. *Frontiers in Ecology and the Environment*. 2019; 17: 445–454. <https://doi.org/10.1002/fee.2088>
4. Broecker WS, Takahashi T. Calcium carbonate precipitation on the Bahama Banks. *Journal of Geophysical Research*. 1966; 71: 1575. <https://doi.org/10.1029/JZ071i006p01575>
5. Smith SV, Key GS. Carbon Dioxide and Metabolism in Marine Environments. *Limnology and Oceanography*. 1975; 20: 493–495. <https://doi.org/10.4319/lo.1975.20.3.0493>
6. Kinsey DW. Alkalinity changes and coral reef calcification1. *Limnology and Oceanography*. 1978; 23: 989–991. <https://doi.org/10.4319/lo.1978.23.5.0989>
7. Chisholm JRM, Gattuso J. Validation of the alkalinity anomaly technique for investigating calcification of photosynthesis in coral reef communities. *Limnology and Oceanography*. 1991; 36: 1232–1239. <https://doi.org/10.4319/lo.1991.36.6.1232>

8. Wolf-Gladrow DA, Zeebe RE, Klaas C, Körtzinger A, Dickson AG. Total alkalinity: The explicit conservative expression and its application to biogeochemical processes. *Marine Chemistry*. 2007; 106: 287–300. <https://doi.org/10.1016/j.marchem.2007.01.006>
9. Langdon C, Gattuso J-P, Andersson A. Measurements of calcification and dissolution of benthic organisms and communities. Riebesell U, Fabry V, Hansson L, Gattuso J, editors. *Guide to best practices for ocean acidification research and data reporting*. 2010; 213–232.
10. Friis K, Körtzinger A, Wallace DWR. The salinity normalization of marine inorganic carbon chemistry data. *Geophysical Research Letters*. 2003; 30: 1–4. <https://doi.org/10.1029/2002GL015898>
11. Berner RA, Lasaga AC, Garrels RM. Carbonate-silicate geochemical cycle and its effect on atmospheric carbon dioxide over the past 100 million years. *Am J Sci*; (United States). 1983; 283:7. <https://doi.org/10.2475/ajs.283.7.641>
12. Cai W-J, Guo X, Chen C-TA, Dai M, Zhang L, Zhai W, et al. A comparative overview of weathering intensity and HCO<sub>3</sub><sup>-</sup> flux in the world's major rivers with emphasis on the Changjiang, Huanghe, Zhujiang (Pearl) and Mississippi Rivers. *Continental Shelf Research*. 2008; 28: 1538–1549. <https://doi.org/10.1016/j.csr.2007.10.014>
13. Likens GE, Bormann FH, Johnson NM. Acid Rain. *Environment: Science and Policy for Sustainable Development*. 1972; 14: 33–40. <https://doi.org/10.1080/00139157.1972.9933001>
14. Avila A, Rodà F. Assessing decadal changes in rainwater alkalinity at a rural Mediterranean site in the Montseny Mountains (NE Spain). *Atmospheric Environment*. 2002; 36: 2881–2890. [https://doi.org/10.1016/S1352-2310\(02\)00098-5](https://doi.org/10.1016/S1352-2310(02)00098-5)
15. Cyronak T, Santos IR, Erler DV, Eyre BD. Groundwater and porewater as major sources of alkalinity to a fringing coral reef lagoon (Muri Lagoon, Cook Islands). *Biogeosciences*. 2013; 10: 2467–2480. <https://doi.org/10.5194/bg-10-2467-2013>
16. Jiang Z-P, Tyrrell T, Hydes DJ, Dai M, Hartman SE. Variability of alkalinity and the alkalinity-salinity relationship in the tropical and subtropical surface ocean. *Global Biogeochemical Cycles*. 2014; 28: 729–742. <https://doi.org/10.1002/2013GB004678>
17. Courtney TA, Andersson AJ. Evaluating measurements of coral reef net ecosystem calcification rates. *Coral Reefs*. 2019; 38: 997–1006. <https://doi.org/10.1007/s00338-019-01828-2>
18. Luijendijk E, Gleeson T, Moosdorf N. Fresh groundwater discharge insignificant for the world's oceans but important for coastal ecosystems. *Nature Communications*. 2020; 11: 1260. <https://doi.org/10.1038/s41467-020-15064-8> PMID: 32152309
19. Legendre P. *Imodel2: Model II Regression*. R package version 1.7–3. <https://CRAN.R-project.org/package=Imodel2>. 2018.
20. R Core Team. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. 2020; [www.R-project.org/](http://www.R-project.org/).
21. Santos IR, Eyre BD, Huettel M. The driving forces of porewater and groundwater flow in permeable coastal sediments: A review. *Estuarine, Coastal and Shelf Science*. 2012; 98: 1–15. <https://doi.org/10.1016/j.ecss.2011.10.024>
22. Taniguchi M, Dulai H, Burnett KM, Santos IR, Sugimoto R, Stieglitz T, et al. Submarine Groundwater Discharge: Updates on Its Measurement Techniques, Geophysical Drivers, Magnitudes, and Effects. *Front Environ Sci*. 2019;7. <https://doi.org/10.3389/fenvs.2019.00141>
23. Shen C-C, Liu K-K, Lee M-Y, Lee T, Wang C-H, Lee H-J. A novel method for tracing coastal water masses using Sr/Ca ratios and salinity in Nanwan Bay, southern Taiwan. *Estuarine, Coastal and Shelf Science*. 2005; 65: 135–142. <https://doi.org/10.1016/j.ecss.2005.05.010>
24. Silbiger NJ, Donahue MJ, Lubarsky K. Submarine groundwater discharge alters coral reef ecosystem metabolism. *Proceedings of the Royal Society B: Biological Sciences*. 2020; 287: 20202743. <https://doi.org/10.1098/rspb.2020.2743> PMID: 33323091
25. Mayfield KK, Eisenhauer A, Santiago Ramos DP, Higgins JA, Horner TJ, Auro M, et al. Groundwater discharge impacts marine isotope budgets of Li, Mg, Ca, Sr, and Ba. *Nature Communications*. 2021; 12: 148. <https://doi.org/10.1038/s41467-020-20248-3> PMID: 33420047
26. Bockmon EE, Dickson AG. An inter-laboratory comparison assessing the quality of seawater carbon dioxide measurements. *Marine Chemistry*. 2015; 171: 36–43. <https://doi.org/10.1016/j.marchem.2015.02.002>